



## **Response of Organic Manure and Azotobacter on Quality and Leaf Nutrient Status of Strawberry (*Fragaria x ananassa* Duch.) Cv. Winter Dawn**

**G. Chandramohan Reddy <sup>a\*</sup>, R. K. Goyal <sup>a</sup> and A. K. Godara <sup>a</sup>**

<sup>a</sup> Department of Horticulture, College of Agriculture, CCS Haryana Agricultural University, Hissar, Haryana, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJPSS/2021/v33i2430785

#### Editor(s):

(1) Dr. Hon H. Ho, State University of New York, USA.

#### Reviewers:

(1) Renisson Neponuceno de Araújo Filho, Universidade Federal do Tocantins, Brazil.

(2) Jane Alexander Ruley, University of Juba, South Sudan.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:  
<https://www.sdiarticle5.com/review-history/75266>

**Original Research Article**

**Received 19 August 2021**  
**Accepted 23 October 2021**  
**Published 18 December 2021**

### **ABSTRACT**

The field experiment was conducted during two subsequent years, *i.e.*, 2017-18 and 2018-19 at the Department of Horticulture, Hisar with aim to study the effect of organic manures and *Azotobacter* on quality and leaf nutrient status of strawberry cv. Winter Dawn under hi-tech greenhouse conditions of Haryana. In this experiment, organic manures in combination with biofertilizers comprised of nine treatments, *viz.*, T<sub>1</sub>: Sand: FYM (3:1), T<sub>2</sub>: Sand: FYM (2:1), T<sub>3</sub>: Sand: Vermicompost (3:1), T<sub>4</sub>: Sand: Vermicompost (2:1), T<sub>5</sub>: Sand: FYM (3:1) + *Azotobacter*, T<sub>6</sub>: Sand: FYM (2:1) + *Azotobacter*, T<sub>7</sub>: Sand: Vermicompost (3:1) + *Azotobacter*, T<sub>8</sub>: Sand: Vermicompost (2:1) + *Azotobacter* and T<sub>9</sub>: Control (Sand); were tested as potting media in complete randomized block design. Maximum TSS, ascorbic acid, anthocyanin content and minimum acidity content and leaf with maximum N, P and K content were observed under sand: vermicompost (2:1) + *Azotobacter* in comparison to the other treatments. The results of this experiment revealed that, the combination of sand: vermicompost (2:1) with *Azotobacter* showed significant influence on leaf nutrient content of strawberry and produced better results with respect to quality of strawberry fruits.

**Keywords:** Strawberry; organic manure; Azotobacter; Quality.

## 1. INTRODUCTION

The modern cultivated strawberry (*Fragaria x ananassa* Duch.), is an inter-specific hybrid of two new world species, *Fragaria virginiana* Duch., a wild strawberry native to Virginia and *Fragaria chilonensis* (L.) Duch., a South American strawberry, and belongs to the family Rosaceae. Fruits are popular for their distinctive succulent nature, bright red colour, juicy texture, aroma and are a rich source of vitamins, minerals and phenolics. Strawberry plants having shallow root system need effective and balanced nutrient management, thus, nutrient status of soil is most important factor affecting the production of strawberry. The modern-day intensive crop cultivation involves bulk application of inorganic fertilizers, which are not only in short supply but also costly. They also pollute the environment, soil and water. Farmyard manure and vermicompost are the main organic components for horticultural crop production [1,2].

Biofertilization is the new concept that is being adopted in numerous agricultural and horticultural crop production systems, reducing cost of production and minimizing environmental pollution [3]. Nitrogen fixing bacteria (*Azospirillum*, *Azotobacter*), phosphate solubilizers (*Pseudomonas*, *Bacillus*) and phosphate absorbers (*Mycorrhizal* fungi) are the most important biofertilizers applied for horticultural crop cultivation [4]. *Azotobacter* is a free-living heterotrophic nitrogen fixing bacteria that enhances plant growth by stimulating growth promoting substances such as auxins, gibberellins and vitamins etc. [5,6]. Vermicompost supplies all essential macro elements such as N, P, K, Ca, Mg and microelements such as Fe, Zn, Mo and Co to fulfil plant requirements during the crop growth period. The application of bio-fertilizer along with organic or inorganic fertilizers significantly increases fertilizer use efficiency, nutrient uptake, growth and yield of strawberry [7]. Hence, the application of vermicompost and *Azotobacter* should help to produce quality berries with higher yields.

In India, strawberry cultivation is primarily confined to temperate zones of Uttarakhand, Himachal Pradesh, Jammu and Kashmir, hills of West Bengal (Darjeeling), Tamil Nadu (Ooty) and the tropical and subtropical zones of Haryana,

Uttar Pradesh, Punjab, Maharashtra (Mahabaleshwar) and Karnataka (Bengaluru) with an area of 1000 hectare and production of 5000 tonne [8].

Keeping the above facts in view, a field trial was conducted to .....???? And so what was the problem to be addressed by this study? The authors should ensure that the aim of the study and the problem to be addressed are clearly shown at this point before delving into the materials and methods section.

## 2. MATERIALS AND METHODS

The present investigations were carried out at Hitech- greenhouse, CCS Haryana Agricultural University, Hisar, (Haryana) in which country? during the year 2017-18 and 2018-19, to study the effect of organic manures with biofertilizers on quality and leaf NPK status of strawberry cv. Winter Dawn. Uniform runners were selected for planting and single healthy uniform runners were planted in each pot after treating with copper oxychloride (0.1%) for 10 minutes. The planting was done in the 2<sup>nd</sup> week of October in both consecutive years (2017-18 and 2018-19). There were nine treatments, viz., T<sub>1</sub>: Sand: FYM (3:1), T<sub>2</sub>: Sand: FYM (2:1), T<sub>3</sub>: Sand: Vermicompost (3:1), T<sub>4</sub>: Sand: Vermicompost (2:1), T<sub>5</sub>: Sand: FYM (3:1) + *Azotobacter*, T<sub>6</sub>: Sand: FYM (2:1) + *Azotobacter*, T<sub>7</sub>: Sand: Vermicompost (3:1) + *Azotobacter*, T<sub>8</sub>: Sand: Vermicompost (2:1) + *Azotobacter* and T<sub>9</sub>: Control (Sand), replicated five times in complete randomized block design. Each twelve inches' pot was filled with sand, FYM and vermicompost; volume by volume on basis of treatment imposition. *Azotobacter* were inoculated @10 ml per plant in the form of liquid at 20 and 40 days after transplanting in the pots.

The recommended dose of fertilizers (NPK 1.95: 0.8:2.75 g plant<sup>-1</sup>) was applied in the form of water-soluble fertilizers (Urea, 19:19:19 and KNO<sub>3</sub>). Fertigation was done at weekly intervals after trans planting. Strawberry plants were subjected to uniform application of recommended dose of fertilizers, plant protection measures and other cultural practices. Observations on various quality parameters were recorded by using standard methods. Twenty fruits from each treatment were randomly selected to record the data on quality parameters. TSS was measured with the help of ERMA hand refractometer (0-32°Bx), Titratable acidity (%), anthocyanin and

vitamin C (mg/100 gm) were estimated by using the method suggested by [9]. Total N was determined by micro-Kjeldahl method [10]. Diacid mixture was used for wet digestion of samples to determine the P and K. Phosphorus was determined by Vanadomolybo-phosphoric acid yellow colour method and potassium content in leaf samples was estimated by flame photometric method [11]. The data was analysed with the help of a window-based computer package OPSTAT [12], to calculate standard error of means SE(m), standard error of difference in mean SE(d), and critical difference between treatments mean CD.

### 3. RESULTS AND DISCUSSION

#### 3.1 Total Soluble Solids (<sup>o</sup>Brix)

TSS of strawberry fruits was significantly affected by the application of different ratios of organic manures and *Azotobacter* with respect to various treatments as revealed by Table 1. The maximum TSS were observed under Sand: Vermicompost (2:1) + *Azotobacter* (T<sub>8</sub>), whereas the minimum TSS was recorded from the control (T<sub>9</sub>) plants during both years of investigation. An increase in TSS with plant growth promoting bacteria might be due to the quick metabolic transformation of starch and pectin into soluble compounds and rapid translocation of sugars from leaves to the developing fruits [13]. The

results of present study are in harmony with the findings of [14] who recorded maximum TSS in fruits harvested from the plants grown under organic manures with biofertilizers, which might be due to the reason that biofertilizers in combination with *Azotobacter* increased the accumulation of carbohydrates and metabolites which were converted into disaccharides leading to higher TSS in strawberry fruits. Similar results were also obtained by authors [15,16,17,18] in strawberry.

#### 3.2 Titratable Acidity (%)

The effect of various treatments on acidity of strawberry fruits is shown in Table 1. The minimum titratable acidity was recorded in sand: vermicompost (2:1) + *Azotobacter* (T<sub>8</sub>), which was at par with the application of sand: vermicompost (3:1) + *Azotobacter* (T<sub>7</sub>) and maximum was recorded in T<sub>9</sub> (Control) over the other treatments. The reduction in titratable acidity may be attributed to conversion of organic acids and photosynthates into sugar during fruit ripening by applying biofertilizers [19]. These findings are in close conformity with the results of [20] who observed that the reduction in titratable acidity may also be due to utilization of acids as a substrate for respiration during the ripening and neutralization of organic acids due to potassium in tissues. Similar findings were also reported [21,22] in strawberry.

**Table 1. Effect of organic manures in combination with biofertilizers on quality parameters of strawberry**

Treatments	TSS ( <sup>o</sup> Brix)		Acidity (%)		Ascorbic acid (mg/100 ml)		Anthocyanin (mg/100 ml)	
	2018	2019	2018	2019	2018	2019	2018	2019
T <sub>1</sub> : Sand: FYM (3:1)	6.58	6.70	0.82	0.82	42.48	43.04	34.48	35.65
T <sub>2</sub> : Sand: FYM (2:1)	6.64	6.79	0.82	0.80	42.69	43.38	35.82	36.90
T <sub>3</sub> : Sand: Vermicompost (3:1)	6.95	7.10	0.79	0.78	47.11	47.66	39.33	40.54
T <sub>4</sub> : Sand: Vermicompost (2:1)	7.05	7.11	0.78	0.77	47.83	48.32	40.62	41.62
T <sub>5</sub> : Sand: FYM (3:1) + <i>Azotobacter</i>	7.08	7.18	0.77	0.76	45.98	47.11	39.75	40.66
T <sub>6</sub> : Sand: FYM (2:1) + <i>Azotobacter</i>	7.13	7.22	0.76	0.74	46.11	47.32	41.15	42.02
T <sub>7</sub> : Sand: Vermicompost (3:1) + <i>Azotobacter</i>	7.45	7.56	0.73	0.71	50.22	51.39	43.52	44.15
T <sub>8</sub> : Sand: Vermicompost (2:1) + <i>Azotobacter</i>	7.59	7.73	0.70	0.68	51.01	52.15	45.27	45.92
T <sub>9</sub> : Control: (Sand only)	6.03	6.08	0.83	0.82	40.17	40.70	31.24	32.07
CD at 5%	0.62	0.60	0.05	0.07	2.43	2.28	1.99	1.79

**Table 2. Effect of organic manures in combination with biofertilizers on leaf N, P and K content of strawberry**

Treatments	N (%)		P (%)		K (%)	
	2018	2019	2018	2019	2018	2019
T <sub>1</sub> : Sand: FYM (3:1)	2.54	2.61	0.34	0.41	1.23	1.39
T <sub>2</sub> : Sand: FYM (2:1)	2.65	2.72	0.39	0.46	1.38	1.46
T <sub>3</sub> : Sand: Vermicompost (3:1)	2.75	2.78	0.41	0.50	1.41	1.51
T <sub>4</sub> : Sand: Vermicompost (2:1)	2.81	2.85	0.48	0.57	1.53	1.59
T <sub>5</sub> : Sand: FYM (3:1) + <i>Azotobacter</i>	2.70	2.74	0.40	0.51	1.40	1.52
T <sub>6</sub> : Sand: FYM (2:1) + <i>Azotobacter</i>	2.77	2.86	0.49	0.57	1.51	1.63
T <sub>7</sub> : Sand: Vermicompost (3:1) + <i>Azotobacter</i>	2.96	3.01	0.50	0.60	1.56	1.62
T <sub>8</sub> : Sand: Vermicompost (2:1) + <i>Azotobacter</i>	3.06	3.08	0.60	0.69	1.69	1.71
T <sub>9</sub> : Control: (Sand only)	1.98	2.04	0.24	0.31	1.05	1.13
CD at 5%	0.09	0.07	0.06	0.05	0.12	0.07

### 3.3 Ascorbic Acid (mg/ 100 ml)

The maximum ascorbic acid content was observed under sand: vermicompost (2:1) + *Azotobacter* (T<sub>8</sub>), whereas the minimum was recorded from the control (T<sub>9</sub>) plants during both years of investigation as shown in Table 1. The increase in ascorbic acid content might be due to the increased efficiency of microbial inoculants to fix atmospheric nitrogen, increase in availability of nitrogen and excretion of growth promoting hormones which accelerates the physiological process like carbohydrates synthesis, etc. An increase in ascorbic acid content with *Azotobacter* and vermicompost application might be due to the quick metabolic transformation of starch and pectin into soluble compounds and rapid translocation of sugars from leaves to the developing fruits [14]. Similar results were found by authors [23,24] in strawberry.

### 3.4 Anthocyanin (mg/100 mg)

Anthocyanin significantly increased by the application of combination of organic manures and bio-fertilizer over the control. The maximum anthocyanin contents were observed under sand: vermicompost (2:1) + *Azotobacter* (T<sub>8</sub>), whereas the minimum was recorded from the control (T<sub>9</sub>) plants during both years of investigation as shown in Table 1. These results may be due to the synergistic effect of macronutrients supplied through organic and bio-fertilizer which improved the vegetative characteristics of the plants thereby affecting the synthesis of chlorophyll that enhanced the process of photosynthesis and the assimilation of carbon dioxide which led to increased fruit quality [25].

### 3.5 N, P and K Status of Leaf

The leaf nutrient content was influenced significantly by the organic manure in combination with biofertilizer (Table 2). The maximum leaf N, P and K content were recorded under (T<sub>8</sub>) sand: vermicompost (2:1) + *Azotobacter* treatment followed by T<sub>7</sub>, T<sub>6</sub> and T<sub>5</sub> and minimum in the control (T<sub>9</sub>). Plants fertilized with vermicompost have shown greater ability to assimilate essential macronutrients as also observed by other researchers [26]. The higher uptake of nitrogen and phosphorous may be due to improved symbiotic N<sub>2</sub> fixation and also due to improved phosphatase activity, thereby improving phosphorous mobilization and uptake from the root zone. The increase in uptake of nutrients may be due to extra amount of nutrients supplied by the organic fertilizers and provision of conducive physical environment which helps in better root growth and absorption of nutrients from the soil [27]. The results of present study are in line with the findings of [4] who observed that maximum contents of nitrogen, phosphorus and potassium were observed in guava with recommended dose of fertilizers along with vermicompost in guava cv. Sardar; and [28] who recorded higher leaf macro nutrient content in strawberry plants fertilized with *Azotobacter* in combination with organic fertilizers in comparison to control.

## 4. CONCLUSION

From the results, it is concluded that quality strawberry plants are raised with organic manures fortified by biofertilizers. This was shown by the application of Sand: vermicompost

(2:1) along with the combination of *Azotobacter* which showed significantly increased quality parameters as well as maximum leaf NPK content compared to other treatments. Then what is the most plausible recommendation arising from your study that can inform farmers, decision makers, organizations biased on promoting strawberries and other fruits as well as firms dealing in production of vermicompost and bio fertilizers.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Hedley MJ, White RE, Nye PH. Plant-induced changes in the rizosphere of rape (*Brassica napus* var. Emerald) seedlings: iii. changes in L value, soil phosphate fractions and phosphatase activity. *New phytology* 1982; 91(1): 45-5.
- Seneviratne G, Henakaarchchi MPNK, Weerasekara MLMAW, Nandasena KA. Soil organic carbon and nitrogen pools as influenced by polyphenols in different particle size fractions under tropical conditions. *Journal of National Science Foundation of Sri Lanka*. 2009; 37(1) 205-208.
- Singh A, Singh JN. Effect of biofertilizers and bioregulators on growth, yield and nutrient status of strawberry cv. Sweet Charlie. *Indian Journal of Horticulture*. 2009; 66(2): 220-224.
- Cheng GW, Breen PJ. Activity of phenylalanine ammonia-lyase (PAL) and concentrations of anthocyanins and phenolics in developing strawberry fruit. *Journal of American Society of Horticulture Sciences*. 1991;116(5): 865-869.
- Indian Horticulture Data base. National horticultural board, Ministry of Agriculture and Family Welfare, New Delhi, India. 2018.
- Sindhu SS, Nisha V, Seema D, Deepika C. Biofertilizer application for growth stimulation of horticultural crops. *Haryana Journal of Horticultural Sciences*. 2010; 39 (1&2); 48-70.
- Karma B, Kumar S, Gupta AK, Syamal MM. Effect of organic, inorganic and bio-fertilizer on growth, flowering, yield and quality of strawberry (*Fragaria x ananassa* Duch.) cv. Chandler. *International Journal of Current Microbiology and Applied Sciences*. 2017; 6(5): 2932-2939.
- Sexana M, Gandhi PC. Indian Horticulture Data base. National horticultural board, Ministry of Agriculture and Family Welfare, New Delhi, India. 2018.
- A.O.A.C. Official Methods of Analysis, 15<sup>th</sup> Edn. Association of Official Analytical Chemist, Washington, D.C. 1990.
- Jackson ML. Soil Chemical Analysis Prentice Hall of India Private Limited, New Delhi. 1967. p. 498.
- Jackson D. Temperate and sub-tropical fruit production. Academic Press. NY. 1985. pp. 114-115.
- Sheoran OP. Statistical Package for Agricultural Research Workers. CCS Haryana Agricultural University, Hisar. (2004) <http://hau.ernet.in/>.
- Yadav SK, Khokhar UU, Sharma SD, Kumar P. Response of strawberry to organic versus inorganic fertilizers. *Journal of Plant Nutrition*. 2016; 39(2): 194-203.
- Tripathi VK, Sanjeev K, Kaushal K, Sarvesh K, Vishal D. Influence of *Azotobacter*, *Azospirillum* and PSB on vegetative growth, flowering, yield and quality of strawberry cv. Chandler. *Progressive Horticulture*. 2016; 48(1): 48-52.
- Tonutare T, Moor U, Szajdak L. Strawberry anthocyanin determination by pH differential spectroscopic method- how to get true results. *Acta Scientific Polonium, Horticulture*. 2014; 13(3): 35-47.
- Hassan AH. Effect of nitrogen fertilizer in the form of organic, inorganic and bio fertilizer applications on growth, yield and quality of Strawberry. *Middle East Journal of Applied Sciences*. 2015; 5(2): 604-617.
- Mahavir S, Devi SK, Sandeep S, Mahendra B. Effect of integrated nutrient management on growth, yield and quality traits of strawberry (*Fragaria x ananassa* Duch.) cv. Chandler. *Journal of Pharmcognacy and Photochemistry*. 2018;1: 712-715.
- Motsara MR. Fertilizer development and consultancy organization. In: Biofertilizer technology marketing and uses-a source book cum, glossary, New Delhi. 1995; 184.
- Ghosh B, Irenaeus TKS, Kundu S, Datta P. Effect of organic manuring on growth, yield and quality of sweet orange. *Acta Horticulture*. 2011;1024: 121-125.

20. Ahmed R, Karim MR, Quddus MA, Ahmed S, Siddiky MA. Response of strawberry to NPKS on yield in terrace soil. Bangladesh Journal of Agriculture Research. 2018; 43(1): 89-97.
21. Tomic JM, Milivojevic JM, Pesakovic MI. The response to bacterial inoculation is cultivar-related in strawberries. Turkish Journal of Agriculture and Forestry. 2015; 39(2): 332-341.
22. Kumar N, Ram RB, Mishra PK (2015) Response of vermicompost and *Azotobacter* on growth and yield of Sweet Charlie strawberry. Int J Agri Sci Res 5(4): 13-20
23. Hazarika BN, Ansari S. Biofertilizers in fruit crops - A review. Agriculture Review. 2007; 28(1): 69-747.
24. Singh R, Sharma RR, Kumar SI, Gupta RK, Patil RT. Vermicompost substitution influence growth, physiological disorders, fruit yield and quality of strawberry. Bio resource Technology. 2008; 99: 8507-8511
25. Umar I, Wali VK, Kher R, Jamwal M. Effect of FYM, urea and *Azotobacter* on growth, yield and quality of strawberry cv. Chandler. Notulae Botanicae Horti Agrobot Cluj-Napoca. 2009; 37(1): 139-143.
26. Singh SR, Zargar MY, Najar GR, Ishaq MI, Hakeem SA (2012) Effect of integrated nutrient supply on yield, fertility and quality of strawberry under rain fed temperate conditions. Journal of Indian Society of Soil Science 60(1): 79-82.
27. Yadav SK, Khokhar UU, Sharma SD, Kumar P. Response of strawberry to organic versus inorganic fertilizers. Journal of Plant Nutrition. 2016; 39(2): 194-203.
28. Amir K, Fouzia I. Chemical nutrient analysis of different composts and their effect on the growth of a vegetative crop *Pisum sativum*. Asian Journal of Plant Sciences and Research. 2011; 1(1): 116-130.

© 2021 Reddy et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/75266>