

Migraine: Prevalence, Predisposing Factors, and Impact on Quality of Life among Saudi Board Family Medicine Residents in Riyadh, KSA

Rajab Ali ^a, Fahad Alshareef ^a, Mohammed Al-Qahtani ^a,
Osamah Almuqrin ^a, Hamad Aloraini ^a, Assaf Alshibani ^{a*}
and Abdullah Hommadi ^a

^a King Saud Medical City, Riyadh, Saudi Arabia.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/INDJ/2023/v20i1386

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102166>

Original Research Article

Received: 22/04/2023

Accepted: 24/06/2023

Published: 27/06/2023

ABSTRACT

Background: Migraine is a prevalent and incapacitating neurological illness that affects a considerable section of the global population. In a community context, the purpose of this study was to examine the prevalence of migraine and its associated comorbidities, triggers, and impact on quality of life.

Methodology: A cross-sectional study was undertaken among Saudi Board Family residents. A standardized questionnaire was used to collect information on migraine prevalence, related comorbidities, triggers, and impact on quality of life. Additionally, demographic information was collected. Using descriptive statistics, chi-square tests, and logistic regression analysis, the data were evaluated.

*Corresponding author: E-mail: asaf.oti@hotmail.com;

Results: The prevalence of migraine was 23.8% among the individuals. Multiple comorbidities, including bronchial asthma, depression, and irritable bowel syndrome, were found to be strongly linked with migraine. Participants cited stress, sleep deprivation or disturbance, and dehydration as the most prevalent migraine triggers. Participants with migraine reported a considerably higher prevalence of physical activity limitation, missed work, and abstinence from social events as a result of their migraine headache compared to participants without migraine.

Conclusion: This study provides important insights into the prevalence, comorbidities, triggers, and impact of migraine on quality of life among Saudi Board Family residents. The findings underscore the substantial impact of migraine on individuals and society and emphasize the need for appropriate therapy techniques that account for the unique nature of migraine and its associated comorbidities and triggers.

Keywords: Migraine; quality of life; family medicine.

1. INTRODUCTION

Migraine headaches characterizes by throbbing headaches that mostly occur on one side of the head and are accompanied by following symptoms nausea, vomiting, photophobia, and phonophobia [1]. It is typically a chronic condition that affects persons under the age of 35 years and worsens over time [2].

The attacks of this type of disease, characterized by severe pain in the head, can last for hours or even for days. The pain can be so intense that it might even affect the daily activities of the affected person [3]. The patient experience of chronic migraine is more narrative dysfunction of the pain pause network. The preventive treatment of migraine and the biomarkers could predict the patient's response through treatment [4]. Along with a better understanding of migraine will lay the way of development for improvement on treatments that are specially designed for those migraine affected patients. This type of pain experience includes affective, cognitive, and sensory brain networks. The practical perspective of pain includes prefrontal, anterior cingulate, insular and cortex of somatosensory, which is demonstrated for chronically migraine patients.

This type of pain experience is similar to the former findings for the patient in episodic migraine [5].

The exact pathophysiology of migraine is still not fully understood, but it is believed to involve complex interactions between genetic, environmental, and neurochemical factors. Current theories suggest that migraine is initiated by activation of the trigeminovascular system, which involves the release of pro-inflammatory peptides such as calcitonin gene-related peptide

(CGRP) and substance P in response to various triggers. This leads to vasodilation and neurogenic inflammation, which in turn activate pain fibers in the meninges and other structures surrounding the brain [6].

There are several different treatment options available for migraine, depending on the severity and frequency of the attacks. Acute treatments include over-the-counter pain relievers, such as ibuprofen or acetaminophen, as well as prescription medications such as triptans and ergotamine's. These drugs work by targeting specific receptors involved in the pathophysiology of migraine, such as serotonin receptors and CGRP receptors. However, their efficacy is limited by side effects and contraindications, and they may not be effective for all patients [7]. Preventive treatments are also available for patients who experience frequent or severe migraine attacks.

These include beta blockers, calcium channel blockers, antidepressants, and anti-seizure medications. They work by decreasing the frequency and severity of attacks by modulating various mechanisms involved in migraine pathophysiology. However, their efficacy is also limited by side effects and potential drug interactions [8].

Family Medicine residents are a critical group of healthcare professionals who play a significant role in patient care. If they are suffering from migraine, it can impact their ability to provide quality patient care and affect patient outcomes. Therefore, it is essential to identify the risk factors associated with migraine in this population and develop interventions to mitigate them. Migraine can have a significant impact on an individual's quality of life, affecting their work, social life, and personal relationships. By

identifying the prevalence of migraine and associated risk factors among family medicine residents, we can help improve their quality of life and well-being.

This study will provide valuable insights into the prevalence of migraine among Family Medicine residents, which can inform future research in this area. It can also serve as a baseline for future studies on the effectiveness of preventive and management strategies for migraine in this population.

2. METHODOLOGY

This was a cross-sectional study, and the gathering of first-hand data is needed. Therefore, detailed data gathering was needed to assess the retrieved data and get the needed outcome. The data collection and analysis process started on April 31, 2023, and completed prior to June 2023. Data gathered would be analyzed using appropriate statistical packages.

2.1 Study Area/Setting

This study was conducted at the primary health care of all health sectors which they have residency program accredited by SCFHS in Riyadh City, Saudi Arabia.

2.2 Study Subjects

The study included all Saudi board family medicine residents training in Riyadh, KSA.

2.3 Study Design

The study was a cross-sectional study since we contacted the subjects of the study one time only.

2.4 Eligibility Criteria

Inclusion criteria: The study targets all levels of Saudi board family medicine residents training in Riyadh city.

Exclusion criteria: The study excluded all physicians who are not involved in a residency program accredited by SCFHS.

2.5 Sample Size

This is a convenience sampling. It is estimated that almost more than 400 Saudi board family medicine residents training in Riyadh, KSA. Accordingly, the representative sample size in

the current study was determined using the epi Info (<http://www.raosoft.com/sampleize.html>) with a margin of error of 5%, a confidence level of 95%, and the response distribution of the prevalence counted for 50% for the lack of local studies. To which we added 10% as a non-response rate.

Therefore, the final sample size was 199 participants.

2.6 Data Collection Methods

Multiple-choice questionnaires would be implemented through the conduction of a survey for gathering the data, and the analysis would involve quantitative assessment of the data (Spector, 2019). The questionnaires were electronically distributed by a team through WhatsApp, Telegram, etc. (using Google Forms) among the family residents across all residency programs in Riyadh. The questionnaire was anonymous to ensure confidentiality and increase the response rate. The questionnaire covered areas of demographics, trigger factors, history of migraine or any other chronic diseases, the ID Migraine test was used to screen for migraine, and the questionnaire also included their general lifestyle and the impact on work and social activities.

The numeric pain rating scale (NPRS) was used to assess the severity of the headache. This scale was validated in previous studies [9,10].

2.7 Statistical Consideration

The data analysis was conducted using the Statistical Package for Social Science (SPSS) Application for Windows Operating System (version 25). Descriptive statistics were applied to both continuous and categorical variables to describe them. The significant differences between categorical variables were determined by employing the chi-square test and Fisher's exact test. On the other hand, the independent samples t-test was used to test the differences between means. Any P-value below 0.05 was regarded as statistically significant.

3. RESULTS

According to Table 1, there were 199 participants in the study, of which 59.3% were female and 40.7% were male. The participants' mean age was 27.91 years, with a standard deviation of 2.19 years. In terms of training level, 27.1% were

in R1, 28.6% were in R2, and 44.2% were in R3. The bulk of participants (75.9 %) were unmarried, while 23.1% were married and 0.5% were either divorced or widowed. Regarding a migraine family history, 25.6% of subjects reported having a migraine family history. Regarding smoking status, 32.7% of participants reported having smoked at least one cigarette in their lifetime. Regarding comorbidities, the majority of individuals (76.8 %) did not indicate having any. In contrast, 11.1% reported having irritable bowel syndrome, 7.6% depression, and 5.1% asthma or generalized anxiety disorder.

Participants reported hypothyroidism and gastrointestinal disorders at rates of 4.5% and 2.0%, respectively.

Only 0.5% of individuals reported having type 2 diabetes.

Fig. 1 shows the prevalence of migraine among the participants in the study. Out of 199 participants, 54 of them reported having migraine, which represents a prevalence rate of 27.1%. The majority of participants (72.9%) reported not having migraine.

Table 1. Demographic factors of the participants (N=199)

		Count	Column N %
Gender	Female	118	59.3%
	Male	81	40.7%
Age	Mean (SD)	27.91 (2.19)	
Residency level	R1	54	27.1%
	R2	57	28.6%
	R3	88	44.2%
Marital status	Single	151	75.9%
	Married	46	23.1%
	Divorced	1	0.5%
	Widow	1	0.5%
Do you have family history of migraine?	No	148	74.4%
	Yes	51	25.6%
Have you ever smoked a cigarette?	No	134	67.3%
	Yes	65	32.7%
Comorbidities	None	152	76.8%
	Bronchial asthma	10	5.1%
	Depression	15	7.6%
	Irritable bowel syndrome	22	11.1%
	Gastrointestinal disorder	4	2.0%
	Generalized Anxiety disorder	10	5.1%
	Hypothyroidism	9	4.5%
Diabetes mellitus	1	0.5%	




Fig. 1. The prevalence of Migraine among the participants

The relationship between the occurrence of migraine and the participants' demographic characteristics is depicted in Table 2. Males had a greater migraine prevalence (37.0%) than females (20.3%), and the difference was statistically significant ($p=0.009$). Male participants were 2.38 times more likely than female participants to have migraine (95 percent CI: 1.35-4.19), showing that male participants were 2.38 times more likely to have migraine than female participants. There was no connection between migraine prevalence and age ($p=0.682$) or level of education ($p=0.890$). In terms of marital status, the prevalence of migraine was greater among married participants (34.8%) than among single participants (25.2%), but the difference was not statistically significant ($p=0.493$). Participants who had a family history of migraine had a substantially higher migraine prevalence (49.0 %) than those without a family history of migraine (19.6 %; $p = 0.000$). The odds ratio (OR) for having migraine with a family history of migraine was 4.89 (95 percent confidence interval [CI]: 2.71-8.81), indicating that participants with a family history of migraine were nearly 5 times more likely to experience migraine than those without a family history of migraine. There was no significant correlation between migraine prevalence and smoking ($p=0.253$). Those who reported having

comorbidities had a considerably higher prevalence of migraine (51,1 percent) than those who did not report having comorbidities (19.7 %) ($p = 0.000$). The odds ratio (OR) for having migraine with comorbidities was 4.90 (95 % confidence interval [CI]: 2.52-9.54), showing that people with comorbidities were nearly 5 times more likely to have migraine than those without comorbidities.

The relationship between various comorbidities and the prevalence of migraine among the individuals is depicted in Table 3. Participants with bronchial asthma, depression, and irritable bowel syndrome had significantly higher migraine prevalence than those without these comorbidities, as indicated by p-values of 0.002, 0.018, and 0.000, respectively. The odds ratios for migraine with these comorbidities were 5.55 (95 percent confidence interval: 1.66-18.56), 2.31 (95 percent confidence interval: 1.00-5.35), and 3.67 (95 percent confidence interval: 1.73- 7.27), respectively. These findings indicate that bronchial asthma, depression, and irritable bowel syndrome are important migraine risk factors. In contrast, no significant connection was found between migraine prevalence and gastrointestinal illness, generalized anxiety disorder, hypothyroidism, or diabetes.

Table 2. The relation between prevalence of migraine and demographic factors of the participants

		Migraine				P-value
		No		Yes		
		Count	Row N %	Count	RowN %	
Gender	Female	94	79.7%	24	20.3%	0.009*
	Male	51	63.0%	30	37.0%	
Age	Mean (SD)	27.95 (2.26)		27.81 (1.98)		0682
Residency level	R1	38	70.4%	16	29.6%	0.890
	R2	42	73.7%	15	26.3%	
	R3	65	73.9%	23	26.1%	
Marital status	Single	113	74.8%	38	25.2%	0.493
	Married	30	65.2%	16	34.8%	
	Divorced	1	100.0%	0	0.0%	
	Widow	1	100.0%	0	0.0%	
Do you have family history of migraine?	No	119	80.4%	29	19.6%	0.000*
	Yes	26	51.0%	25	49.0%	
Have you ever smoked a cigarette?	No	101	75.4%	33	24.6%	0.253
	Yes	44	67.7%	21	32.3%	
Comorbidities	Yes	23	48.9%	24	51.1%	0.000*
	No	122	80.3%	30	19.7%	

Table 3. The relation between different comorbidities and prevalence of migraine

		Migraine				P-value
		No		Yes		
		Count	Row N %	Count	RowN %	
Bronchial asthma	No	142	75.1%	47	24.9%	0.002*
	Yes	3	30.0%	7	70.0%	
Depression	No	138	75.0%	46	25.0%	0.018*
	Yes	7	46.7%	8	53.3%	
Irritable bowel syndrome	No	136	76.8%	41	23.2%	0.000*
	Yes	9	40.9%	13	59.1%	
Gastrointestinal disorder	No	143	73.3%	52	26.7%	0.299
	Yes	2	50.0%	2	50.0%	
Generalized Anxiety disorder	No	139	73.5%	50	26.5%	0.348
	Yes	6	60.0%	4	40.0%	
Hypothyroidism	No	139	73.2%	51	26.8%	0.669
	Yes	6	66.7%	3	33.3%	
Diabetes mellitus	No	144	72.7%	54	27.3%	0.541
	Yes	1	100.0%	0	0.0%	

Table 4. Symptoms reported by the participants related to headache and its relation to migraine

		Total		Migraine				P-value
		N	Percent	No		Yes		
				N	Percent	N	Percent	
Symptoms	None	107	53.8%	102	95.3%	5	4.7 %	0.000*
	Depressedmode	41	20.6%	18	43.9%	23	56.1%	0.000*
	Difficulty concentrating	66	33.2%	27	40.9%	39	59.1%	0.000*
	Photophobia	38	19.1%	7	18.4%	31	81.6%	0.000*
	Phonophobia	30	15.1%	7	23.3%	23	76.7%	0.000*
	Irritability	41	20.6%	16	39.0%	25	61.0%	0.000*
	Extremefatigue	37	18.6%	15	40.5%	22	59.5%	0.000*
	Nausea	32	16.1%	5	15.6%	27	84.4%	0.000*
	Aura	1	0.5%	1	100%	0	0.0%	0.541

Table 4 displays the headache symptoms reported by participants and their relationship to migraine. Participants who reported symptoms of depression, trouble concentrating, photophobia, phonophobia, irritability, extreme weariness, and nausea had a significantly higher prevalence of migraine, as measured by p-values of 0.000, compared to those who did not report these symptoms. The odds ratios for having migraine with these symptoms ranged between 2.89 and 11.37, showing that persons with these symptoms were considerably more likely to have migraine than those without them. With a p-value of 0.541, there was no significant connection between the prevalence of migraine and the presence of aura.

Fig. 2 depicts how individuals cope with migraine headaches under intense pain. Taking analgesics (42,2 %) and napping were the most

often cited pain relief methods (36.2 %). Resting (29.1 %) and dimming the lights (20.6%) were also frequently cited techniques. Less frequently stated tactics included consuming coffee (10,6 %), experimenting with hot or cold compressors (2.5 %), and engaging in regular physical activity (3.5 %). Notably, a sizeable minority of individuals (45.7%) reported not employing any specific migraine headache pain management method during intense pain.

Fig. 3 depicts the common migraine headache triggers identified by participants. The most often cited cause was stress (26.8 %), followed by sleep deprivation or disturbance (22.7 %) and dehydration (18.2 %). Participants also indicated fasting (14.6%), severe job load (18.2%), and prolonged reading as triggers (10.1 percent). Notably, a considerable majority of patients (53.5%) reported having no identifiable migraine

headache cause. Other identified triggers by participants included continuous exposure to loud sound (5.1%), an overly lighted environment (5.6%), extremely loud noises (8.1%), menstruation (4.5%), smoking (3.5%), caffeine (4.5%), and physical activities (2.5 %).

The relationship between migraines and participants' quality of life is depicted in Table 5. With a p-value of 0.000, participants with migraine reported a considerably higher prevalence of physical activity limitation than those without migraine. The odds ratios for migraine and physical activity limitation were 16.93 (95 % confidence interval: 5.83-49.18) for "often" and 51.29 (95 % confidence interval: 14.99-175.51) for "always" compared to "never" Participants with migraine were substantially more likely to miss work and withdraw from social activities due to their migraine headache, with p-values of 0.000 and 0.001, respectively, compared to those without migraine. The odds ratios for

migraine and missing work or avoiding social activities were 5.77 (95 % confidence interval: 1.92-17.32) and 5.91 (95 % confidence interval: 2.45-14.21), respectively.

4. DISCUSSION AND CONCLUSION

In the current study, the prevalence of migraine among Saudi Board Family residents was 23.8 % which is similar to that reported by different studies. In a comprehensive review undertaken by Albalawi M et al. of 36 Saudi Arabia-based research, the authors observed a 22.56 % prevalence (95 % confidence interval of 17.27-28.33 percent). In addition, several studies conducted in Saudi Arabia reported a prevalence of migraine among general population to be ranged between 2.5 %- 37.5 % [9,11–17] while among the university students, the prevalence of migraine increased to between 7.1 %- 71.6 % [18–21]. This result highlights the significant burden of migraine on individuals and society.




Fig. 2. How do you deal with the migraine headache at times of immense pain?




Fig. 3. What is a general trigger for your migraine headache?

Table 5. The relation between migraine and quality of life

		Total		Migraine				P-value
		Count	Column N %	No		Yes		
				Count	RowN %	Count	RowN %	
Does your headache prevent you from doing any type of physical activities?	Never	99	49.7%	90	90.9%	9	9.1%	0.000*
	Rarely	58	29.1%	41	70.7%	17	29.3%	
	Often	35	17.6%	12	34.3%	23	65.7%	
	Always	7	3.5%	2	28.6%	5	71.4%	
Have you ever missed work due tomigraine headache?	No	180	90.5%	142	78.9%	38	21.1%	0.000*
	Yes	19	9.5%	3	15.8%	16	84.2%	
Have you ever refrained from participating in social activity due to migraine headache?	No	161	80.9%	136	84.5%	25	15.5%	0.00*
	Yes	38	19.1%	9	23.7%	29	76.3%	

In addition, it was reported in the current study, that there is a significant difference in the prevalence of migraine among genders where the prevalence of migraine was significantly higher among male residents than females with higher level of risk of migraine in men by 2.38 times than females. This finding contradicts prior study that demonstrated a higher prevalence of migraines in women [22–24]. The study found no significant correlation between migraine prevalence and age or residency level. These results are in agreement with the results reported by different previous studies which showed no correlation between the prevalence of migraine and the age or education of the population [25,26]. However, this results may be associated with some limitations of the study including the small sample size, therefore, further research using higher sample size is important to investigate the possible association between migraine prevalence and those demographic factors. In addition, the current study found that the prevalence of migraine was insignificantly different between participants according to the marital status which is consistent with the results of previous study which showed that there is no association between marital status and the prevalence of migraine [27]. Moreover, the current study showed that having family history of migraine increase the risk for development migraine by 4.89 times compared with those without family history of migraine. This result is consistent with previous research indicating that migraine has a major hereditary component [26,28].

Moreover, the current study showed that there are many comorbidities that was found to be

associated with the incidence of migraine. The study showed that prevalence of migraine with higher among participants reported having bronchial asthma, depression, and irritable bowel syndrome (IBS). On the other hand, no association was reported between incidence of migraine and incidence of gastrointestinal disorders, generalized anxiety disorders, diabetes mellitus, or hypothyroidism. As reported by different previous study, there is a strong association between incidence of migraine and presence of bronchial asthma, depression, and IBS [13,29–31]. These findings provide support for this conclusion. Our results showed that having those comorbidities increase the risk for developing migraine by 2.31-5.55 times.

One reason for the correlation between incidence of migraine and bronchial asthma is the sharing of the two disorders with some underlying causes as inflammation and autonomic dysfunction [32,33]. Different previous studies showed that both disorders share similar pathophysiological pathways including the higher level of pro-inflammatory cytokine levels, the activation of mast cell, and the dysfunction of the autonomic nervous system [34]. Considering the relation between depression and incidence of migraine, some shared genetic and environmental risk factors between the two disorders as stressful life events, neurotransmitter imbalances, and fluctuations of different hormones may be the reason of this relation [35,36]. Multiple studies have shown that depression is a migraine risk factor and can worsen migraine symptoms [26,37]. Moreover, there is different reasons for the correlation between migraine and IBS including the common underlying processes as

the disturbance of gut-brain axis, the altered gut flora, and visceral hypersensitivity [38,39]. Several previous studies have demonstrated that IBS and migraine have similar pathophysiological pathways, including the presence of higher levels of pro-inflammatory cytokines, altered gut microbiota, and central nervous system sensitivity [40].

According to the study, stress, sleep deprivation or disturbance, and dehydration were the most often reported migraine triggers. In addition to hunger, heavy workload, and prolonged reading, participants cited these as additional triggers. These results are consistent with prior research identifying stress and sleep disturbance as key migraine triggers [41–43]. The fact that 53.5% of patients reported having no identified migraine headache trigger demonstrates the complexity of migraine and the need for tailored treatment choices [44].

Moreover, the current study also investigate the impact of migraine of the quality of life of the participants where patients with migraine were found to report a higher prevalence of physical activity restriction, missing of work day, and social event avoidance because of the migraine. This is consistent with several previous studies which showed that migraine has a significant negative impact on the patients' daily activities, social functioning, job productivity [45–48]. The odds ratios for having migraine and limiting physical activity, missing work, or refraining from social activities ranged from 5.77 to 51.29, indicating that individuals with migraine were substantially more likely to have these adverse impacts than those without migraine.

In conclusion, the findings of this study provide valuable insights into the prevalence, comorbidities, triggers, and quality of life impact of migraine among Saudi Board Family residents. In addition, our results shed the light on the significant negative impact of migraine on patients and society and the importance for presence of management strategies that consider the particular characteristics of migraine and its triggers. This study's findings can inform the development of individualized migraine management approaches that take into consideration the distinct demands and limitations faced by individuals with this condition.

CONSENT AND ETHICAL APPROVAL

The research was conformed to Helsinki Declaration. The protocol of the study was

approved by the institutional Review Board (IRB) of King Saud Medical City (KSMC), with a reference number of H1R1-09-Apr23-01. A written informed consent was taken from each resident accepted participant. Administrative approvals were also taken.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Shaik MM, Hassan NB, Tan HL, Gan SH. Quality of Life and Migraine Disability among Female Migraine Patients in a Tertiary Hospital in Malaysia. *Biomed Res Int.* 2015;2015:1-9. DOI:10.1155/2015/523717
2. AlHarbi F, AlAteeq M. Quality of life of migraine patients followed in neurology clinics in Riyadh, Saudi Arabia. *J Fam Community Med.* 2020;27(1):37. DOI:10.4103/jfcm.JFCM_185_19
3. Koverech A, Cicione C, Lionetto L, et al. Migraine and cluster headache show impaired neurosteroids patterns. *J Headache Pain.* 2019;20(1):61. DOI:10.1186/s10194-019-1005-0
4. Fu C, Zhang Y, Ye Y, et al. Predicting response to tVNS in patients with migraine using functional MRI: A voxels-based machine learning analysis. *Front Neurosci.* 2022;16. DOI:10.3389/fnins.2022.937453
5. Gollion, C., Lerebours, F., Nemmi, F. et al. Insular functional connectivity in migraine with aura. *J Headache Pain.* 2022;23:106. DOI:10.1186/s10194-022-01473-1
6. Iyengar S, Johnson KW, Ossipov MH, Aurora SK. CGRP and the Trigeminal System in Migraine. *Headache J Head Face Pain.* 2019;59(5):659-681. DOI:10.1111/head.13529
7. Vargas BB. Acute Treatment of Migraine. *Contin Lifelong Learn Neurol.* 2018;24(4):1032-1051. DOI:10.1212/CON.0000000000000639
8. Silberstein SD, Holland S, Freitag F, Dodick DW, Argoff C, Ashman E. Evidence-based guideline update: Pharmacologic treatment for episodic migraine prevention in adults: Table 1. *Neurology.* 2012;78(17):1337-1345. DOI:10.1212/WNL.0b013e3182535d20
9. Bamalan BA, Khojah AB, Alkhateeb LM, et al. Prevalence of migraine among the

- general population, and its effect on the quality of life in Jeddah, Saudi Arabia. *Saudi Med J.* 2021;42(10):1103-1108. DOI:10.15537/smj.2021.42.10.20210575
10. Ibrahim NK, Aloitabi AK, Alhazmi A, Alshehri R, Saimaldhr R, Murad M. Prevalence, predictors and triggers of migraine headache among medical students and interns in King Abdulaziz University, Jeddah, Saudi Arabia. *Pakistan J Med Sci.* 2017;33(2). DOI:10.12669/pjms.332.12139
 11. Abukanna AMA, Alenezi NAA, Alenezi AAT, Alenezi MAT. Prevalence and impact of primary headache on quality of life in the general population of Arar, Northern Saudi Arabia. *Med Sci I Anal Artic.* 2021;25(112):2431-2439.
 12. Al Jumah M, Al Khathaami AM, Kojan S, Hussain M, Thomas H, Steiner TJ. The prevalence of primary headache disorders in Saudi Arabia: a cross-sectional population-based study. *J Headache Pain.* 2020;21(1):11. DOI:10.1186/s10194-020-1081-1
 13. AlQarni M, Fayi K, Al-sharif M, Siddiqui A, Alhazzani A. Prevalence of migraine and non-migraine headache and its relation with other diseases in the adults of Aseer Region, Saudi Arabia. *J Fam Med Prim Care.* 2020;9(3):1567. DOI:10.4103/jfmpc.jfmpc_962_19
 14. Alotaibi S, Deshwali S, Alotaibi B, et al. Prevalence of migraine headache among people in dawadmi city, kingdom of Saudi Arabia: cross sectional study. *Eur. J Pharm Med Res.* 2018;5:270-273.
 15. Almalki ZA, Alzhrani MAG, Altowairqi AT, et al. Prevalence of Migraine Headache in Taif City, Saudi Arabia. *J Clin Med Res.* 2018;10(2):125-133. DOI:10.14740/jocmr3277w
 16. Muayqil T, Al-Jafen BN, Al-Saaran Z, et al. Migraine and Headache Prevalence and Associated Comorbidities in a Large Saudi Sample. *Eur Neurol.* 2018;79(3-4):126-134. DOI:10.1159/000487317
 17. Jabbar MA, Ogunniyi A. Sociodemographic Factors and Primary Headache Syndromes in a Saudi Community. *Neuroepidemiology.* 1997;16(1):48-52. DOI:10.1159/000109670
 18. Gouhar G, Tamimm R, Almahri S, et al. Migraine among Princess Nourah Bint Abdulrahman University students in Riyadh, Saudi Arabia. *Int J Med Dev Ctries.* Published online 2018:103-108. DOI:10.24911/IJMDC.51-1522164334
 19. Garah M, Neyaz H, Shaqrun F, et al. Prevalence of migraine among female students at taibah university, kingdom of Saudi Arabia. *Int J Adv Res.* 2016;4(7):1526-1534. DOI:10.21474/IJAR01/995
 20. Al Jumah M, Awada A, Al Azzam S. Headache syndromes amongst schoolchildren in Riyadh, Saudi Arabia. *Headache J Head Face Pain.* 2002; 42(4):281-286. DOI:10.1046/j.1526-4610.2002.02081.x
 21. Kanjo M, H.Alsaati R, M.Jassomah O, H.Alhindi S, F.Jamjoom L, A.Albogami M. The prevalence of migraine headache among students of Fakeeh College in Jeddah, Saudi Arabia. *Med Sci I Anal Artic.* 2021;25(108):320-327.
 22. Burton WN, Conti DJ, Chen C-Y, Schultz AB, Edington DW. The Economic Burden of Lost Productivity Due to Migraine Headache: A Specific Worksite Analysis. *J Occup Environ Med.* 2002;44(6):523-529. DOI:10.1097/00043764-200206000-00013
 23. Peterlin BL, Gupta S, Ward TN, MacGregor A. Sex Matters: Evaluating Sex and Gender in Migraine and Headache Research. *Headache J Head Face Pain.* 2011;51(6):839-842. DOI:10.1111/j.1526-4610.2011.01900.x
 24. Al-Hassany L, Haas J, Piccininni M, Kurth T, Maassen Van Den Brink A, Rohmann JL. Giving Researchers a Headache – Sex and Gender Differences in Migraine. *Front Neurol.* 2020;11. DOI:10.3389/fneur.2020.549038
 25. Stewart WF, Roy J, Lipton RB. Migraine prevalence, socioeconomic status, and social causation. *Neurology.* 2013;81(11):948-955. DOI:10.1212/WNL.0b013e3182a43b32
 26. Amiri P, Kazeminasab S, Nejadghaderi SA, et al. Migraine: A Review on Its History, Global Epidemiology, Risk Factors, and Comorbidities. *Front Neurol.* 2022;12. DOI:10.3389/fneur.2021.800605
 27. Rafi A, Islam S, Hasan MT, Hossain G. Prevalence and impact of migraine among university students in Bangladesh: findings from a cross-sectional survey. *BMC Neurol.* 2022;22(1):68. DOI:10.1186/s12883-022-02594-5
 28. Lateef TM, Cui L, Nakamura E, Dozier J, Merikangas K. Accuracy of Family History

- Reports of Migraine in a Community-Based Family Study of Migraine. *Headache J HeadFace Pain.* 2015;55(3):407-412. DOI:10.1111/head.12501
29. Buse DC, Reed ML, Fanning KM, et al. Comorbid and co-occurring conditions in migraine and associated risk of increasing headache pain intensity and headache frequency: results of the migraine in America symptoms and treatment (MAST) study. *J Headache Pain.* 2020;21(1):23. DOI:10.1186/s10194-020-1084-y
30. Kikui S, Chen Y, Ikeda K, Hasebe M, Asao K, Takeshima T. Comorbidities in patients with migraine in Japan: a cross-sectional study using data from National Health and Wellness Survey. *BMJ Open.* 2022;12(11):e065787. DOI:10.1136/bmjopen-2022-065787
31. Chang F-Y, Lu C-L. Irritable Bowel Syndrome and Migraine: Bystanders or Partners? *J Neurogastroenterol Motil.* 2013;19(3):301-311. DOI:10.5056/jnm.2013.19.3.301
32. Wang L, Deng Z-R, Zu M-D, Zhang J, Wang Y. The comorbid relationship between migraine and asthma: a systematic review and meta-analysis of population-based studies. *Front Med.* 2021;7. DOI:10.3389/fmed.2020.609528
33. Kang L-L, Chen P-E, Tung T-H, Chien C-W. Association between asthma and migraine: A systematic review and meta-analysis of observational studies. *Front Allergy.* 2021;2. DOI:10.3389/falgy.2021.741135
34. Salahi M, Parsa S, Nourmohammadi D, et al. Immunologic aspects of migraine: A review of literature. *Front Neurol.* 2022;13. DOI:10.3389/fneur.2022.944791
35. Zhang Q, Shao A, Jiang Z, Tsai H, Liu W. The exploration of mechanisms of comorbidity between migraine and depression. *J Cell Mol Med.* 2019;23(7):4505-4513. DOI:10.1111/jcmm.14390
36. Stam AH, de Vries B, Janssens ACJW, et al. Shared genetic factors in migraine and depression: Evidence from a genetic isolate. *Neurology.* 2010;74(4):288-294. DOI:10.1212/WNL.0b013e3181cbcd19
37. Ashina S, Serrano D, Lipton RB, et al. Depression and risk of transformation of episodic to chronic migraine. *J Headache Pain.* 2012;13(8):615-624. DOI:10.1007/s10194-012-0479-9
38. Raskov H, Burcharth J, Pommergaard H-C, Rosenberg J. Irritable bowel syndrome, the microbiota and the gut-brain axis. *Gut Microbes.* 2016;7(5):365-383. DOI:10.1080/19490976.2016.1218585
39. Cámara-Lemarroy CR, Rodríguez-Gutierrez R, Monreal-Robles R, Marfil-Rivera A. Gastrointestinal disorders associated with migraine: A comprehensive review. *World J Gastroenterol.* 2016;22(36):8149. DOI:10.3748/wjg.v22.i36.8149
40. Arzani M, Jahromi SR, Ghorbani Z, et al. Gut-brain Axis and migraine headache: a comprehensive review. *J Headache Pain.* 2020;21(1):15. DOI:10.1186/s10194-020-1078-9
41. Fernández-de-las-Peñas C, Fernández-Muñoz JJ, Palacios-Ceña M, Parás-Bravo P, Cigarán-Méndez M, Navarro-Pardo E. Sleep disturbances in tension-type headache and migraine. *Ther Adv Neurol Disord.* 2018;11:175628561774544. DOI:10.1177/1756285617745444
42. Houle TT, Butschek RA, Turner DP, Smitherman TA, Rains JC, Penzien DB. Stress and sleep duration predict headache severity in chronic headache sufferers. *Pain.* 2012;153(12):2432-2440. DOI:10.1016/j.pain.2012.08.014
43. Tiseo C, Vacca A, Felbush A, et al. Migraine and sleep disorders: a systematic review. *J Headache Pain.* 2020;21(1):126. DOI:10.1186/s10194-020-01192-5
44. Mayans L, Walling A. Acute Migraine Headache: Treatment Strategies. *Am Fam Physician.* 2018 Feb 15;97(4):243-251.
45. Burton WN, Landy SH, Downs KE, Runken MC. The Impact of Migraine and the Effect of Migraine Treatment on Workplace Productivity in the United States and Suggestions for Future Research. *Mayo Clin Proc.* 2009;84(5):436-445. DOI:10.1016/S0025-6196(11)60562-4
46. Begasse de Dhaem O, Sakai F. Migraine in the workplace. *eNeurologicalSci.* 2022;27:100408. DOI:10.1016/j.ensci.2022.100408
47. Wong LP, Alias H, Bhoo-Pathy N, et al. Impact of migraine on workplace productivity and monetary loss: a study of employees in banking sector in Malaysia. *J*

- Headache Pain.2020;21(1):68.
DOI:10.1186/s10194-020-01144-z
48. Mannix S, Skalicky A, Buse DC, et al. Measuring the impact of migraine for evaluating outcomes of preventive treatments for migraine headaches. Health Qual Life Outcomes.2016;14(1):143.
DOI:10.1186/s12955-016-0542-3

APPENDIX: QUESTIONNAIRE

1. Gender

- a) Male
- b) Female

2. Age:

3. Residency level

- a) R1
- b) R2
- c) R3

4. Marital status

- a) Single
- b) Married
- c) Widow
- d) Divorced

5. Do you have family history of migraine?

- a) Yes
- b) No

6. Have you ever smoked a cigarette?

- a) Yes
- b) No

7. Have you been diagnosed with any other chronic diseases?

- a) Diabetes mellitus
- b) Hypertension
- c) Irritable bowel syndrome
- d) Gastrointestinal disorder
- e) Bronchial asthma
- f) Anxiety
- g) Depression
- h) Endocrine disease

For questions 8 – 10. During the last three months did you have the following with your headache

8. Experienced a headache that limited your activities for a day or more?

- a) Yes
- b) No

9. Felt nauseated or sick to your stomach when you have a headache?

- a) Yes
- b) No

10. Light bothered you when you have a headache?

- a) Yes
- b) No

11. Describe how intense your pain on a scale of '0' to '10' with '0' representing no pain and '10' representing extreme pain?

- a) 0
- b) 1
- c) 2
- d) 3
- e) 4
- f) 5
- g) 6
- h) 7
- i) 8
- j) 9
- k) 10

12. What symptoms do you experience during your migraine headache?

- a) Difficulty concentrating
- b) Photophobia
- c) Phonophobia
- d) Nausea
- e) Irritability
- f) Depressed mood
- g) Extreme fatigue
- h) None

13. How do you deal with the problem at times of immense pain?

- a) Take analgesics
- b) Try sleeping
- c) Try hot or cold compressors
- d) Drink coffee
- e) Turning off the light
- f) Sleep
- g) Rest
- h) Exercising regularly

14. What is a general trigger for migraine?

- a) Stress
- b) Intense work-load
- c) Continuous exposure to loud sound
- d) Caffeine
- e) Physical activities
- f) Fasting
- g) Dehydration
- h) Overly lit room
- i) Extreme loud noises
- j) Smoking
- k) Sleep deprivation or disturbance
- l) Menstruation
- m) Reading for long period of time

15. Does your headache prevent you from doing any type of physical activities?

- a) Always
- b) Often
- c) Rarely
- d) Never

16. Have you ever missed work due to migraine headache?

- a) Yes
- b) No

17. Have you ever refrained from participating in social activity due to migraine headache?

- a) Yes
- b) No

© 2023 Ali et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/102166>