



# Influence on Crop Health and Yields as Affected by Soil Organic Matter and Crop Residue Management: An Overview

Jahnabi Hazarika <sup>a\*</sup>, Shilpi Gupta <sup>a</sup>, Nanhe Lal Saroj <sup>b++</sup>,  
Tikendrajit Saikia <sup>a</sup> and Manuranjan Roy <sup>c#</sup>

<sup>a</sup> Department of Soil Science, College of Agriculture, Assam Agricultural University, Jorhat-13, Assam, India.

<sup>b</sup> Department of Agricultural Science, SRM University, Delhi-NCR, Sonapat, Haryana-131029, India.

<sup>c</sup> Krishi Vigyan Kendra, Arunachal, Silchar, Cachar, India.

## Authors' contributions

This work was carried out in collaboration among all authors. Author JH finalized the topic and wrote the overview and the first draft of the manuscript. Authors SG and NLS managed the tabulated value of the study and find review related to this topic. However, authors TS and MR managed the literature searches. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJECC/2023/V13i103131

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/106898>

Review Article

Received: 24/07/2023

Accepted: 27/09/2023

Published: 03/10/2023

## ABSTRACT

For growth of plants and crop productivity, soil is a crucial component. However, just a very small portion of the soil is genuinely fertile for farming these days, and if we managed it incorrectly, it could be depleted. It has been demonstrated that organic nutrients increase crop production, quality, and yield, which in turn improves the quality as well as richness of the soil's characteristics and creates a more favorable environment for the soil's beneficial microorganisms. According to

<sup>++</sup> Assistant Professor;

<sup>#</sup> Subject Matter Specialist;

<sup>\*</sup>Corresponding author: E-mail: jahnabihazarika555@gmail.com;

reports, organic carbon in the soil and soil matter are the most significant indicators of soil quality and soil health. They are also advantageous for the sustainability of agriculture. Crop residue can be used effectively to improve soil quality and increase or maintain SOM's physical and chemical qualities.

**Keywords:** Soil organic matter; organic carbon; crop yield; crop residue.

## 1. INTRODUCTION

The component of soil known as soil organic matter (SOM) is made up of plant and animal debris in different stages of decomposition, soil microbe cells and tissues, and compounds that soil microbes produce [12]. The element carbon (C), which is a component of all organic compounds by definition, makes up approximately fifty percent of the mass of SOM. Four to six times as much carbon (C) as there is in the entire world's plants can be found in the organic matter (OM) in soil profiles. Therefore, the global C balance, which essentially regulates global climate change, depends on soil organic matter [54]. Large amounts of plant nutrients are present in soil organic matter, which also serves to act as a slow-release nutrient storage facility (particularly for nitrogen), creates soil aggregation, promotes nutrient exchange, holds moisture, decreases compaction, minimizes surface crusting, and improves water to penetrate into the soil. In addition, OM gives the majority of soil-dwelling organism's nutrients and energy. Some organic chemicals in soil have direct impacts on plants that stimulate development in addition to the actions outlined above [14,52,54].

Land conservation techniques used extensively in agriculture include crop residue management (CRM). It offers different levels of soil nutrients to boost crop output. Crop residues have an impact on the physical, chemical, and biological qualities of the soil as well as the infiltration rate and runoff water. By shielding the soil surface from the sun's heat, residue affects soil temperature. Crop residue accumulation on the soil surface slows evaporation rates. The organic matter content of residue-covered soils is higher than that of bare soils. According to observations, soil maintains greater amounts of moisture when it is maintained on the surface of the soil rather than being incorporated into the soil through residues [49]. Food production in agriculture depends on soil. Maintaining soil quality is essential for continued food production, waste digestion, and storage residue returning nutrients to the soil, carbon sequestration, and gas exchange. Today,

just a very small portion of the soil may be used to grow food crops, and if we don't take care of it, it could be damaged, polluted, or depleted [11].

## 2. SOURCES OF SOIL ORGANIC MATTER

### 2.1 Following are some of the Sources of Soil Organic Matter

**Plants:** Plant tissue is the initial source of soil organic matter. Fallen leaves are the main source of organic matter given to the soil in a forest. Under natural circumstances, trees, shrubs, grasses, and various other native plants seasonally produce significant amounts of organic wastes from the crowns and roots. After the crop is harvested, the part of the crop plant (such as the tops, twigs, roots, etc.) left in the soil also functions as an important source of organic matter. As a result, the main sources of organic material are found in plant tissue.

**Animals:** Animals are typically regarded as secondary suppliers of organic matter. The organic matter in the soil is also contributed by soil organisms like insects, millipedes, nematodes, and others. After they pass away, soil microorganisms significantly contribute to the organic matter of the soil.

**Manures:** Manures are a valuable source of organic matter for the soil and their addition to the soil enhances the organic matter to some level. Examples of manures include farmyard manure, compost, green manure, fish, meal, and oil cake.


### 2.2 Effect of Soil Organic Matter Content on Soil Health

Healthy crops, healthy livestock, and a healthy human population supported by nutrient-balanced diets and environmentally sound environments all depend on healthy soils. In order to achieve such a crucial interconnection, favorable SOM content is essential. Ancient civilizations have known for millennia about the significance of SOM content to agricultural output [35], and soil scientists have known about it for a minimum of two centuries [19,20]. In the modern

day, [1] vividly described the importance of the content of SOM for agricultural production, which established the way for a growing interest in the research of global terrestrial soil organic carbon, or SOC, sequestration and processes affecting its stabilization [50]. Crop productivity is impacted by SOM content because it contributes to improving and maintaining the condition of the soil [32] and quality [46]. SOM content and various biological, chemical, and physical features and functions are tightly related. Using organic manure and manure combined with chemical fertilizers on a regular basis improved soil quality, stabilized crop productivity, and aided adaptation to climate change, according to a long-term research study in eastern Europe (the Czech Republic, Slovakia, and Poland) [36]. Through its beneficial effects on soil characteristics and processes, SOM concentration is a crucial measure of soil health [17]. Two important characteristics that are impacted by SOM concentration in terms of agronomic productivity are plant accessible water capacity (PAWC) and plant available vital nutrients, particularly nitrogen (N).

A field experiment was conducted by [47] on mango (*Mangifera indica* L.) to study the effects of farmyard manure (FYM) on soil organic carbon (SOC) content and on the build-up of fertility over three growing seasons (2006–2007, 2007–2008, and 2008–2009) at Orissa, India. According to them soil moisture contents declined continuously in all treatments during the growing seasons, probably due to the extraction of water from the soil profile by the mango trees (Fig. 1). FYM-treated plots had 47.7 – 59.7% higher gravimetric moisture contents compared to the RDF-treated and control plots. This could be attributed to the incorporation of organic manure, which conserved soil moisture, favoring higher levels of uptake of water from deeper layers.

Saha et al. [47] Also conclude that (Table 1) the soil organic carbon contents increased significantly in the FYM-treated plots. The highest soil organic carbon density (554 g m<sup>-2</sup>) and stock (5.55 Mg ha<sup>-1</sup>) were recorded in the treatment with FYM at 10 kg plant<sup>-1</sup>



**Fig. 1. Soil moisture content [% (w/w)] under ‘Mallika’ mango trees treated with FYM or an inorganic fertilizer, or untreated (control) trees**

Data are mean values (n = 36) pooled from three growing seasons (2006–2009)

**Table 1. Effect of organic manure or inorganic fertilizer on the soil organic carbon content under ‘Mallika’ mango trees**

| Treatment                       | SOC density (g m <sup>-2</sup> )<br>Status (2009) | SOC stock (Mg ha <sup>-1</sup> )<br>Status (2009) |
|---------------------------------|---|---|
| Control                         | 234 ± 19*   | 2.33 ± 0.09                                       |
| 5.0 kg FYM plant <sup>-1</sup>  | 376 ± 32  | 3.75 ± 0.33                                       |
| 7.5 kg FYM plant <sup>-1</sup>  | 420 ± 25  | 4.22 ± 0.32                                       |
| 10.0 kg FYM Plant <sup>-1</sup> | 554 ± 44  | 5.55 ± 0.65                                       |
| RDF                             | 275 ± 17  | 2.75 ± 0.11                                       |
| LSD (P ≤ 0.05)                  | 12.48   | 0.98  |

### 3. PLANT AVAILABLE WATER CAPACITY

Aggregation [30], porosity, and pore size distribution are all impacted by SOM content. According to the majority of soil physicists over the 50 years between 1940 and 1990, SOC enhances water retention at field capacity and at the permanent wilting point with no additional gains in plant available water capacity [21,5 and 44]. According to [25], reducing SOM content in UK soils from 7% to 3% can result in a 10% drop in soil water retention. Similar findings were reported by [29] on the loss in water retention caused by a drop in SOM content. The fall in SOM content-induced decrease in plant available water is partially related to a decrease in aggregate stability. According to [4] rather than a drop in plant available water content, the loss of soil productivity caused by erosion in the northern Great Plains may be attributable to a decline in nutrients and biological activity. According to [37], a rise in SOM content has little effect on Plant Available Water Capacity.

However, several researchers [9,48,44 and 13] have noted a significant impact of SOM content on Plant Available Water Content. For a variety of soil textural groups, [27] discovered highly significant positive relationships between SOM content and PAWC. He came to the conclusion that throughout all textural groups, as SOM concentration increased from 0.5% to 3%, soil PAWC more than doubled. Another scientist [55] suggested spending money on the restoration of SOM content to improve Plant Available Water Capacity and lessen volatility and downside risks in American rain fed maize (*Zea mays* L.). Also [2,3] showed that in California, United States, an increase in SOM water retention results in up to 8.8 cm (3.46 in) more transpiration, or 35 more days without water stress. Even while the rise in Plant Available Water Capacity caused by an increase in SOM content is very minor, it may be crucial for crop growth between rainy times, especially for dry land farming [28].

### 4. SOIL ORGANIC MATTER CONTENT AND CROP YIELD

The mutual improvement of crop production and SOM content further obscures the direct cause-and-effect relationship in field conditions, when crop productivity is influenced by a variety of variables, including biotic and abiotic stressors. According to [40], an increase in SOM content could result in yield increases of 10%–11% for maize and 23%–37% for wheat. According to

[23]'s findings in Denmark, SOC level significantly affected wheat grain yield and aboveground biomass at just 0 to 100 kg N ha<sup>-1</sup> (0 to 89.2 lb N ac<sup>-1</sup>), and this influence dwindled as N input rates increased. Oldfield and colleagues came to the conclusion that improving SOM stocks can help close the yield gap in degraded soils and other situations where there is a significant yield gap. These yield improvements reduce fertilizer inputs by 5% to 7% while bridging the 30% yield gap for maize and 55% yield gap for wheat. Additionally, there was a favorable correlation between Plant Available Water Capacity and SOC content. Plant Available Water Capacity increased modestly for the range of 0.7% to 2.0% SOC and increased with rise in SOC content up to 0.7% [23]. Contrarily, a number of studies [39,53 and 26] suggested no effects of SOM content on crop yield and backed the idea that it is challenging to distinguish the effects of SOM on crop yield from those of nutrients [38] and other factors.

Agricultural management practices have a significant impact on organic matter, including soil carbon, soil quality, and soil health [31,18]. The terrestrial reservoir for the cycling of carbon, nitrogen, and phosphorus is soil organic carbon. Increased soil organic matter has an impact on soil characteristics, including water and nutrient availability. The quantity of agricultural output is ultimately increased by these increments [6,41]. Another scientist [33] recorded that when soil's organic carbon and organic matter levels drop too low, it has a significant impact on the soil's productivity. As a result, soil characteristics deteriorate and nutrient cycle processes are hampered.

Also [43] was conducted an field experiment during rabi 2013-14 and 2014-15 at Agricultural Research Station, Rajasthan, India to find out suitable organic nutrient management practices for higher productivity and economics of wheat. Results (from Table 2) in terms of growth and yield attributes, grain yield and economics of wheat under different treatments were significantly different than control. Pooled data of two years' experimentation showed that the maximum plant height (85.83 cm), number of tillers per meter row length (91.00), number of seed per ear head (55.18), ear head length (9.60 cm), test weight (37.17 g) and grain yield (39.14 q ha<sup>-1</sup>) were recorded with application of FYM @ 10 t ha<sup>-1</sup>+Vermicompost @ 1.25 t ha<sup>-1</sup> followed by treatment FYM @ 5 t ha<sup>-1</sup>+Neem green leaves @ 2 t ha<sup>-1</sup>+Vermicompost @ 1.25 t ha<sup>-1</sup>.

**Table 2. Effect of organic nutrient sources on growth and yield attributes and grain yield of wheat (Pooled data)**

| Treatments     | Plant Height (cm) | No. of tillers m <sup>-1</sup> row | No. of grains ear <sup>-1</sup> head | Ear head Length (cm) | Test Weight (g) | Grain yield(q ha <sup>-1</sup> ) |
|----------------|-------------------|------------------------------------|--------------------------------------|----------------------|-----------------|----------------------------------|
| T <sub>1</sub> | 70.00             | 70.50                              | 44.58                                | 8.31                 | 31.72           | 19.00                            |
| T <sub>2</sub> | 81.17             | 84.83                              | 50.83                                | 9.40                 | 34.90           | 33.09                            |
| T <sub>3</sub> | 72.17             | 75.67                              | 48.07                                | 9.07                 | 32.73           | 27.87                            |
| T <sub>4</sub> | 77.00             | 81.83                              | 50.05                                | 9.49                 | 33.22           | 30.18                            |
| T <sub>5</sub> | 85.83             | 91.00                              | 55.18                                | 9.60                 | 37.17           | 39.14                            |
| T <sub>6</sub> | 73.33             | 79.00                              | 48.75                                | 9.15                 | 33.23           | 29.25                            |
| T <sub>7</sub> | 83.50             | 87.33                              | 51.65                                | 9.42                 | 35.03           | 37.02                            |
| SEm±           | 2.14              | 2.55                               | 1.81                                 | 0.26                 | 1.07            | 1.31                             |
| CD (p=0.05)    | 6.50              | 7.72                               | 5.49                                 | 0.78                 | 3.24            | 3.98                             |

T<sub>1</sub> : control; T<sub>2</sub> : FYM @ 15 t ha<sup>-1</sup>; T<sub>3</sub>: Neem green leaves @ 6 t ha<sup>-1</sup>; T<sub>4</sub> : FYM @ 10 t ha<sup>-1</sup>+Neem green leaves @ 2 t ha<sup>-1</sup>; T<sub>5</sub> :FYM @ 10 t ha<sup>-1</sup>+Vermicompost @ 1.25 t ha<sup>-1</sup>; T<sub>6</sub> : Neem green leaves @ 4 t ha<sup>-1</sup>+Vermicompost @ 1.25 t ha<sup>-1</sup>; T<sub>7</sub>: FYM @ 5t ha<sup>-1</sup>+Neem green leaves @ 2 t ha<sup>-1</sup>+Vermicompost @ 1.25 t ha<sup>-1</sup>

## 5. CROP RESIDUE MANAGEMENT AND CROP HEALTH


Depending on the cultivation method, crop residue may be fully or partially incorporated into the field. The nutritional and soil organic carbon contents are enhanced by the introduction of crop straw. It is useful for reusing leftover nutritional material. The immobilization of nutrients, particularly nitrogen, is facilitated by ploughing, and the better C: N ratio must be adjusted by adding additional nitrogen fertilizer during the time of residues in filling [57]. According to several research, the immobilization of soil nitrogen produced by crop residue put in the field during the first three years of straw integration 30 days before rice planting decreased the production of rice. Later on, however, it had little impact on the yield. According to (58)'s findings, adding straw to the soil significantly increased organic carbon level and storage levels compared to not adding straw, and this increase was likely caused by adding more crop residues to the soil, as previously suggested [34]. According to [16], retention and incorporation of residue led to a significant increase in the total amount of water stable aggregates (15.65%) in surface soil that is 0–15 cm deep and 7.53% in subsurface soil that is 15–30 cm deep. This indicated that using crop residue can result in 2.1-fold greater water stable aggregates than other treatments that do not incorporate or retain residue.

Increasing the amount of crop residue on the soil surface slows the rate of evaporation [24,45].

Therefore, applying crop residue to the soil is the greatest way to add organic amendment and cover the soil's surface. Crop residue effectively manages to concurrently improve organic carbon in the soil, soil nutrients, water availability, productivity demand, and livestock fodder in order to achieve sustainable development. According to [8 and 7], the accessible amount of plant water content was significantly less in conventional tillage than zero tillage under the rice wheat cropping system. According to [10], mulching agricultural residue or partially incorporating it into the soil through conservation tillage improves infiltration by lowering surface sealing and lowering runoff velocity.

## 6. CROP RESIDUE MANAGEMENT AND CROP YIELD

According to [51] incorporating residue into the field resulted in 40 kg nitrogen saving per hectare. According to [58], adding straw to the soil (particularly at a rate of 13500 kilo grams per hectare) was an effective way to improve soil fertility and yield production in China's semi-arid region.80 kg N/ha of yield was reported by [42] with the addition of wheat residue. According to [56], the average grain output improved by 2.65% when crop residue was incorporated as opposed to when it wasn't. However, rice crops had a better straw integration effect on biomass yield. The yields of rice's grain and straw were considerably impacted. In comparison to control sets, rice grain yield was higher when wheat straw alone was applied at 5 or 10 t/ha. According to [22], the residue treated field had



**Fig. 2. Grain yield, straw yield and biological yield as influenced by different residue management techniques**

greater summer output and dry weight of maize plant roots by 18.5% and 15.1%, respectively. They advised returning the proper amount of residue from crops to the field in order to replenish the nutrients necessary for maize growth.

Another scientist [15] conducted an experiment on to study the effect of crop diversification and residue management techniques on yield attributes, yield and soil nutrient on rice. Among the different residue management techniques (Fig. 2) maximum grain, straw and biological yield was obtained in treatment T<sub>5</sub> i.e. 30% residue recycling +vermicompost.

## 7. CONCLUSION

This review assesses whether managing agricultural residues and increasing the amount of organic matter of depleted or degraded soils can improve crop output. The increase in plant-available water capacity of coarse-textured soils and the availability of N at low concentrations of N may be the causes of the positive yield response. Additionally, crop residue provides environmentally friendly and sustainable solutions for satisfying the nutritional needs of soil, agricultural yield, and environmental quality. For different soils, climates, and eco regions, further research is required to identify necessary limits or range of soil organic matter concentration. Additionally, different crop residue management techniques should be chosen to increase crop yields with a sustainable soil environment.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Allison FE. Soil organic matter and its role in crop production. Elsevier; 1973.
- Ankenbauer KJ, Loheide SP. The effects of soil organic matter on soil water retention and plant water use in a meadow of the Sierra Nevada, CA. *Hydrological Processes*. 2017;31(4): 891-901.
- Banwart SA, Noellemeyer E, Milne E. (Eds.). *Soil carbon: Science, management and policy for multiple benefits*. 2014;71. CABI.
- Bauer A, Black AL. Organic carbon effects on available water capacity of three soil textural groups. *Soil Science Society of America Journal*, 1992;56(1):248-254.
- Baver LD. *Soil Physics*. New York: John Wiley & Sons, Ltd; 1940.
- Berzsenyi Z, Gyórfy B, Lap D. Effect of crop rotation and fertilization on maize and wheat yields and yield stability in a long-term experiment. *European Journal of Agronomy*. 2000;13(2-3):225-244.
- Bhattacharyya R, Kundu S, Pandey SC, Singh KP, Gupta HS. Tillage and irrigation effects on crop yields and soil properties under the rice-wheat system in the Indian Himalayas. *Agricultural water management*. 2008;95(9):993-1002.
- Bhattacharyya R, Singh RD, Chandra S, Kundu S, Gupta HS. Effect of tillage and irrigation on yield and soil properties under rice (*Oryza sativa*)-wheat (*Triticum aestivum*) system on a sandy clay loam soil of Uttaranchal. *Indian journal of agricultural science*. 2006;76(7):405-409.
- Bouyoucos GJ. Effect of organic matter on the water-holding capacity and the wilting

- point of mineral soils. *Soil Science*. 1939; 47(5):377-384.
10. Box JE, Bruce RR. The effect of surface cover on infiltration and soil erosion. In *Soil Erosion, Conservation, and Rehabilitation*.1996;107-124. (CRC Press)
  11. Brady NC, Weil RR. Elements of the nature and properties of soils. (No. 631.4 B733E.). Upper Saddle River, NJ, USA: Prentice hall; 2000.
  12. Brady NC, Weil RR. Soil organic matter. The nature and properties of soils. Prentice Hall, Upper Saddle River, New Jersey, USA, 1999;446-490.
  13. Bryant L. Organic matter can improve your soil's water holding capacity. New York: Natural Resources Defense Council; 2015. Available:<https://www.nrdc.org/experts/lara-bryant/organic-matter-can-improve-your-soils-water-holding-capacity>.
  14. Carter MR, Stewart BA. Structure and organic matter storage in agricultural soils.1995;8. (CRC press).
  15. Chaudhary S, Chaturvedi S, Dhyani VC. Effect of crop diversification and residue management techniques on yield attributes, yield and soil nutrient. *International Journal of Chemical Studies*. 2020;8:1107-1111.
  16. Choudhury SG, Srivastava S, Singh R, Chaudhari SK, Sharma DK, Singh SK, Sarkar D. Tillage and residue management effects on soil aggregation, organic carbon dynamics and yield attribute in rice–wheat cropping system under reclaimed sodic soil. *Soil and Tillage Research*. 2014; 136:76-83.
  17. Doran JW, Zeiss MR. Soil health and sustainability: Managing the biotic component of soil quality. *Applied soil ecology*. 2000;15(1):3-11.
  18. Farquharson RJ, Schwenke GD, Mullen JD. Should we manage soil organic carbon in Vertosols in the northern grains region of Australia? *Australian Journal of Experimental Agriculture*. 2003;43(3):261-270.
  19. Feller C, Blanchart E, Bernoux M, Lal R, Manlay R. Soil fertility concepts over the past two centuries: the importance attributed to soil organic matter in developed and developing countries. *Archives of Agronomy and Soil Science*. 2012;58(sup1):S3-S21.
  20. Feller C, Bernoux M. Historical advances in the study of global terrestrial soil organic carbon sequestration. *Waste Management*. 2008;28(4):734-740.
  21. Feustel IC, Byers HG. The comparative moisture-absorbing and moisture-retaining capacities of peat and soil mixtures. (No. 532). US Department of Agriculture; 1936.
  22. Gao F, Zhao B, Dong, S, Liu P, Zhang J. Response of maize root growth to residue management strategies. *Agronomy Journal*. 2018;110(1):95-103.
  23. Ghaley BB, Wösten H, Olesen JE, Schelde K, Baby S, Karki YK, Porter JR. Simulation of soil organic carbon effects on long-term winter wheat (*Triticum aestivum*) production under varying fertilizer inputs. *Frontiers in plant science*. 2018;9:1158.
  24. Gill BS, Jalota SK. Evaporation from soil in relation to residue rate, mixing depth, soil texture and evaporativity. *Soil Technology*. 1996;8(4):293-301.
  25. Gregory AS, Watts CW, Griffiths BS, Hallett PD, Kuan HL, Whitmore AP. The effect of long-term soil management on the physical and biological resilience of a range of arable and grassland soils in England. *Geoderma*. 2009;153(1-2):172-185.
  26. Hijbeek R, Van Ittersum MK, ten Berge HF, Gort G, Spiegel H, Whitmore AP. Do organic inputs matter– A meta-analysis of additional yield effects for arable crops in Europe? *Plant and Soil*. 2017;411:293-303.
  27. Hudson BD. Soil organic matter and available water capacity. *Journal of soil and water conservation*.1994;49(2):189-194.
  28. Johnston AE. Soil organic matter, effects on soils and crops. *Soil use and management*.1986;2(3):97-105.
  29. Johnston A, Poulton P, Coleman K. Soil organic matter: Its importance in sustainable agriculture and carbon dioxide fluxes. *Advances in Agronomy*. 2008; 101:1–57.
  30. Kemper WD, Koch EJ. Aggregate stability of soils from Western United States and Canada: Measurement procedure, correlations with soil constituents (No. 1355). Agricultural Research Service, US Department of Agriculture; 1996.
  31. Lal R. World soils and greenhouse effect: an overview. *Soils and global change*. 1995;1-7.
  32. Lal R. Soil health and carbon management. *Food and Energy Security*. 2016;5(4):212-222.

33. Loveland P, Webb J. Is there a critical level of organic matter in the agricultural soils of temperate regions: a review? *Soil and Tillage research*. 2003;70(1):1-18.
34. Malhi SS, Nyborg M, Goddard T, Puurveen D. Long-term tillage, straw management and N fertilization effects on quantity and quality of organic C and N in a Black Chernozem soil. *Nutrient Cycling in Agroecosystems*. 2011;90:227-241.
35. Manlay RJ, Feller C, Swift MJ. Historical evolution of soil organic matter concepts and their relationships with the fertility and sustainability of cropping systems. *Agriculture, Ecosystems & Environment*. 2007;119(3-4):217-233.
36. Menšík L, Hlisenikovsky L, Kunzová E. The state of the soil organic matter and nutrients in the long-term field experiments with application of organic and mineral fertilizers in different soil-climate conditions in the view of expecting climate change. In *Organic fertilizers-history, production and applications*. IntechOpen; 2019.
37. Minasny B, McBratney AB. Limited effect of organic matter on soil available water capacity. *European Journal of Soil Science*. 2018;69(1):39-47.
38. Murphy BW. Impact of soil organic matter on soil properties— A review with emphasis on Australian soils. *Soil Research*. 2015;53(6):605-635.
39. Oelofse M, Markussen B, Knudsen L, Schelde K, Olesen JE, Jensen LS, Bruun S. Do soil organic carbon levels affect potential yields and nitrogen use efficiency? An analysis of winter wheat and spring barley field trials. *European Journal of Agronomy*. 2016;66:62-73.
40. Oldfield EE, Bradford MA, Wood SA. Global meta-analysis of the relationship between soil organic matter and crop yields. *Soil*. 2019;5(1):15-32.
41. Onemli F. The effects of soil organic matter on seedling emergence in sunflower (*Helianthus annuus* L.). *Plant Soil Environ*. 2004; 50(11):494-499.
42. Paikaray RL, Mahapatra BS, Sharma GL. Integrated nitrogen management in rice (*Oryza sativa*)-wheat (*Triticum aestivum*) cropping system. *Indian Journal of Agronomy*. 2001;46(4):592-600.
43. Parewa HP, Ram M, Jain LK, Choudhary A, Ratnoo SD. Impact of organic nutrient management practices on yield attributes, yield and economics of wheat (*Triticum aestivum* L.). *International Journal of Bio-resource and Stress Management*. 2019;10(3):257-260.
44. Petersen GW, Cunningham RL, Matelski RP. Moisture characteristics of Pennsylvania soils: II. Soil factors affecting moisture retention within a textural class— Silt loam. *Soil Science Society of America Journal*. 1968;32(6):866-870.
45. Prihar SS, Jalota SK, Steiner JL. Residue management for reducing evaporation in relation to soil type and evaporativity. *Soil use and management*. 1996;12(3): 150-157.
46. Reeves DW. The role of soil organic matter in maintaining soil quality in continuous cropping systems. *Soil and Tillage Research*. 1997;43(1-2):131-167.
47. Saha R, Nath V, Kumar D. Effects of farmyard manure on soil organic carbon stock, the pattern of fertility build-up, and plant growth in 'Mallika' mango (*Mangifera indica* L.). *The Journal of Horticultural Science and Biotechnology*. 2010; 85(6):539-543.
48. Salter PJ, Williams JB. The effect of farmyard manure on the moisture characteristic of a sandy loam soil. *Journal of Soil Science*. 1963;14(1):73-81.
49. Singh Y, Sidhu HS. Management of cereal crop residues for sustainable rice-wheat production system in the Indo-Gangetic Plains of India. *Proceedings of Indian National Science Academy*. 2014;80(1):95-114
50. Six J, Frey SD, Thiet RK, Batten KM. Bacterial and fungal contributions to carbon sequestration in agroecosystems. *Soil Science Society of America Journal*. 2006;70(2):555-569.
51. Thakur RC, Bindra, AD, Sood RD, Bhargava M. Effect of fertilizer application and green-manuring on physico-chemical properties of soil and grain yield in rice (*Oryza sativa*)-wheat (*Triticum aestivum*) crop sequence. *Indian Journal of Agronomy*. 1995;40(1):4-13.
52. Vander Wal A, De Boer W. Dinner in the dark: Illuminating drivers of soil organic matter decomposition. *Soil Biology and Biochemistry*. 2017;105:45- 48.
53. Wei W, Yan Y, Cao J, Christie P, Zhang F, Fan M. Effects of combined application of organic amendments and fertilizers on crop yield and soil organic matter: An integrated analysis of long-term experiments. *Agriculture, Ecosystems & Environment*. 2016;225:86-92.



54. Weil RR, Brady NC. Soil organic matter. Nature and properties of soils (15th ed). Pearson Education Limited, England, 2017;545-601.
55. Williams A, Hunter MC, Kammerer M, Kane DA, Jordan NR, Mortensen DA, Davis AS. Soil water holding capacity mitigates downside risk and volatility in US rainfed maize: time to invest in soil organic matter? PloS one, 2016;11(8), e0160974.
56. Xu G, Tan G, Wang Z, Liu L, Yang J. (2009). Effects of wheat-residue application and site-specific nitrogen management on grain yield and quality and nitrogen use efficiency in direct-seeding rice. Scientia Agricultura Sinica. 2009; 42(8):2736-2746.
57. Yadvinder-Singh BS, Timsina J. Crop residue management for nutrient cycling and improving soil productivity in rice-based cropping systems the tropics. Advances in Agronomy. 2005;85:269-407.
58. Zhang P, Chen X, Wei T, Yang Z, Jia Z, Yang B, Ren X. Effects of straw incorporation on the soil nutrient contents, enzyme activities, and crop yield in a semiarid region of China. Soil and Tillage Research. 2016;160:65-72.

---

© 2023 Hazarika et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/106898>