



Performance Evaluation of Brinjal (*Solanum melongena* L.) under Mid Hill Conditions of Uttarakhand, India

Harish A. ^{a*}, S. C. Pant ^a and Pankaj Bahuguna ^a

^a Department of Vegetable Science, College of Horticulture, Veer Chandra Singh Garhwali Uttarakhand University of Horticulture & Forestry, Bharsar - 246123 Pauri Garhwal, Uttarakhand, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i193764

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104649>

Original Research Article

Received: 23/06/2023
Accepted: 28/08/2023
Published: 14/09/2023

ABSTRACT

Brinjal (*Solanum melongena* L.), belongs to the family Solanaceae, which is an important vegetable crop of sub-tropics and tropics. The present investigation was carried out at the Vegetable Research and Demonstration Block, College of Horticulture, VCSG UHF, Bharsar, during *Kharif*, 2017 to evaluate diverse germplasm of brinjal under mid hill conditions. The experiment was laid out in RCBD with three replications. In this study, twenty-seven genotypes *viz.*, DRNKV-02-104, IVBL-116-131, NBJ-19, JB-9, Aruna, JB-2, CH-1045, Swarna Shree, JB, DRNKV-104-43, RCMBL-04-04, IBH-2, Kavya, B. Dewariya, DBR-8, DMU-1, Azad Brinjal-2, Co-11, JB-67, KKM-1, BCB-464, PLR-1, JB-18, Bhagyamati, BB-58, Pusa Shyamala including one check cultivar Pant Samrat were evaluated for different horticultural traits. During the investigation, 18 growth, flowering, fruiting, yield and quality parameters were recorded. The result of variance revealed that the mean sum of square due to treatment were significant at 5% level for all the traits. The genotypes, Aruna, DRNKV-02-104, NBJ-19, JB-2, RCMBL-04-04, IVBL-116-131 and Kavya recorded higher fruit yield and also performed better for other horticultural traits than check cultivar. These genotypes need to

*Corresponding author: E-mail: harishhyr23@gmail.com;

undergo additional testing before being made available as a replacement for currently available brinjal varieties, or they can be included in additional breeding programmes to develop improved varieties or hybrids of brinjal.

Keywords: *Brinjal; evaluation; mean performance; genotypes; hill; Uttarakhand.*

1. INTRODUCTION

Eggplant or Aubergine (*Solanum melongena* L.) belongs to the family Solanaceae (Nightshades) with diploid chromosome number $2n=2x=24$. It was originated in India. Brinjal is a good source of Vitamin A, B and C and also used in curing liver related troubles. For diabetic patient consumption of white brinjal is recommended. Bitter taste of brinjal is due to Solasodine which is a glycoalkaloid [1]. Brinjal is a warm season crop hence it is vulnerable to cold. The low night temperature in North India adversely affects the crop during December to February. It is grown year-round in practically all of India, with the exception of higher altitudes. In Himalayan hilly regions it is grown only during summer (May to September) due to prevailing low temperature in other months.

In India it is having a production of 12607 ('000 MT) from an area of 765 ('000 ha) with a productivity of 17.53 MT/ha. In Uttarakhand, production of brinjal is 32.12 ('000 MT) from an area of 2.75 ('000 Ha) with productivity of 11.68 MT/ha (First advance estimate by Department of agriculture and farmers welfare, Government of India) [2]. This clearly indicates that the productivity of brinjal is very low in Uttarakhand state when compared to National values. This may be due to wide variations in the weather and varying night and day temperatures in hilly terrains, heavy rainfall during flowering and fruiting stage, hail storms, lack of high yielding varieties suited for specific agro climatic conditions, lack of varieties meeting the regional references and specific end use, unscientific method of cultivation by farmers, lack of knowledge in selecting suitable varieties, unavailability of quality seeds, pest and diseases, lack of extension services in remote areas of Uttarakhand are the major production constraints that results in low productivity. Considering the importance of the crop and keeping the above issues in mind this experiment was conducted to select elite genotypes which were better in terms of yield and other yield attributing characters in the mid hill conditions of Pauri Garhwal district, Uttarakhand, and the results obtained were further utilized in the crop improvement of brinjal

or the better performing genotypes could be directly considered for the cultivation.

2. MATERIALS AND METHODS

2.1 Experimental Site

The present investigation was conducted at the Vegetable Research and Demonstration Block, Department of Vegetable Science, College of Horticulture, VCSG Uttarakhand University of Horticulture and Forestry, Bharsar, Pauri Garhwal, Uttarakhand during *kharif* season in 2017. The experimental site is situated between latitude $29^{\circ} 30.05^{\circ}$ North and longitude 78.99° East at an altitude of 1900 m above mean sea level. It belongs to Uttarakhand's mid-hill zone (Bisht and Sharma, 2014). The experimental site's environment is characterised by a mild summer, more precipitation, and a colder or more intensely cold protracted winter. Along with dew, hailstorms, fog, frost, snowfall, and other sporadic occurrences of precipitation, rain is the primary output of precipitation. While the North-East monsoon occasionally brings winter precipitation from November to February, the South-East monsoon starts around the end of June. Throughout this area, snowfall occurs frequently throughout the winter. The valley experiences a hot climate during the summer months for only a few hours each day; nevertheless, nights are chilly and the valleys highest recorded temperature during May and June is between 30°C and 35°C . The coldest months are December and January, with lows between 1°C and -4°C . Relative humidity often reaches its peak during the rainy season (July to August), frequently being measured close to saturation point (92 to 97%), and then progressively declines until December [3]. The soil of research field is comprised of medium clay sandy loam with pH of 6 to 6.5.

2.2 Experimental Design and Plant Material Used

The experiment was laid out in Randomized Complete Block Design (RCBD) with three replications. The trial has total of 26 genotypes *viz.*, DRNKV-02-104, IVBL-116-131, NBJ-19, JB-9, Aruna, JB-2, CH-1045, Swarna Shree, JB,

DRNKV-104-43, RCMBL-04-04, IBH-2, Kavya, B. Dewariya, DBR-8, DMU-1, Azad Brinjal-2, Co-11, JB-67, KKM-1, BCB-464, PLR-1, JB-18, Bhagyamati, BB-58, Pusa Shyamala and Pant Samrat (Check cultivar), in total 27 genotypes are used in the experiment. These genotypes were obtained from the germplasm of IIVR, Varanasi which was collected from the various parts of the country by the institute.

2.3 Management and Cultural Practices Followed

The nursery was prepared inside the polyhouse. The raised bed of size 3 m x 1 m and 15 cm height was prepared. A well decomposed farmyard manure (FYM) about 15 kg and 500g of 10:20:20 NPK complex fertilizer were mixed thoroughly at the time of bed preparation. The seeds were sown on 20th April 2017. The field was prepared to fine tilth by giving 3 ploughings. The plots of size 1.8 m x 1.35 m were prepared at 20 cm apart as per the layout plan. Transplanting was done on 1st June 2017; forty days old seedlings were transplanted with a spacing of 60 cm x 45 cm in the evening hours. Total experimental area thus had a total of 81 plots with 9 numbers of plants in each plot among which 5 plants were randomly taken for data collection, covering a total experimental area of 284.55 m² and a net experimental area of 196.83 m².

During the time of field preparation, well-decomposed farmyard manure (FYM) was added at a rate of 20 tonnes per hectare. Following to the recommendations, 100 N: 60 P₂O₅: 50 K₂O kg NPK per hectare of urea, single super phosphate, and murate of potash were used as inorganic fertilizers. Before transplanting, the soil received a full dose of phosphorus and potassium and a half-dose of nitrogen. 45 days after transplantation, the remaining half of the N dosage was administered. During dry periods irrigation was given twice a week or whenever necessary. Weed growth was more vigorous due to heavy rainfall, hence manual weeding was done at an interval of 15 to 20 days or whenever required. Picking of fruits was done, when the fruits attain suitable marketable size, still immature, tender having attractive and glossy appearance. The fruits were picked at least after a week of insecticidal spraying because of the residual effect.

2.4 Data Recording and Analysis

The five plants were selected randomly using random number table and tagged for

identification. The observations were recorded only from these five tagged plants in every plot for all the characters except days to 50% flowering and yield per plot. Then the value was worked out to compute average value for the character. The quality analysis (Ascorbic Acid content) was estimated as per procedure given by Ranganna [4]. Using MS-Excel and OPSTAT, the statistical analysis was performed for each observed character in the study.

3. RESULTS AND DISCUSSION

3.1 Analysis of Variance

The analysis of variance (Table 1) of the experiment indicated that the mean sum of squares due to genotypes was highly significant for all characters studied *viz.*, plant height, number of primary branches per plant, stem girth, leaf area, days to fifty per cent flowering, number of clusters per plant, number of flowers per cluster, number of fruits per cluster, fruit setting percentage, number of fruits per plant, average fruit weight, length of the fruit, diameter of the fruit, yield per plant, yield per plot, shelf life, total soluble solids and ascorbic acid which revealed the existence of good deal of variability in the germplasm. Similar results were reported by the earlier workers like Kumar et al., [5], Nayak and Nagre [6], Madhavi et al., [7] and Yadav et al., [8].

3.2 Mean Performance of Genotypes for Different Traits

3.2.1 Growth parameters

Taller plant is considered to be desirable because it leads to a greater number of branches and ultimately result in increased yield. Maximum plant height was recorded in RCMBL-04-04 (103.60 cm). Among all the genotypes under study, five genotypes gave higher plant height than check cultivar Pant Samrat (72.51 cm). Similar variations for plant height were also reported earlier by Kumar and Aruguman [5], Nayak and Nagre [6], Dhaka and Soni [9], Mili et al., [10], Solaimana et al., [11] and Madhavi et al., [7] for the trait under study. Number of branches per plant is considered to be desirable because it leads to a greater number of flowers and hence a greater number of fruits and ultimately results in increased productivity. Maximum numbers of branch per plant was recorded in Aruna (9.82) and it was found statistically at par with JB-18 (9.38). In overall,

two genotypes were judged to be superior to the check cultivar. Pant Samrat (9.25). Similar results have also been reported earlier by Karak et al., [12], Lokesh [13], Rahman et al., [14] and Madhavi et al., [7] for number of branches per plants in brinjal. Stem girth is also an important parameter in brinjal, as it helps in preventing the breakage of stems due to more weight of fruits and also prevents the plants from lodging. Maximum stem girth was observed in RCMBL-04-04 (2.40 cm) and it was found statistically at par with the genotype CO-11 (2.37 cm). Five genotypes were judged to be superior to the check cultivar Pant Samrat (1.88 cm). Naik (2006) and Madhavi et al., [7] have also reported similar variation for this trait in brinjal. The large number of leaves might increase the photosynthetic activity and ultimately leads to higher plant yield. Maximum leaf area was recorded in RCMBL-04-04 (127.21 cm²) followed by Bhagyamati (99.91 cm²) and PLR-1 (97.21 cm²). Eight genotypes exhibited more leaf area than the reference cultivar Pant Samrat (92.50 cm²) out of all the genotypes under investigation. Kushwah and Bhandhyopadhya [15] also found similar findings for brinjal leaf area (cm²). The results were represented in Table 2.

3.2.3 Flowering parameters

One of the key determinants of how quickly the fruits reach the harvesting stage is earliness. Minimum days to 50% flowering were recorded in the genotype DRNKV-02-104 (44.00 days) followed by Bhagyamati (56.67 days), JB-67 (57.33 days) and DBR-8 (57.33 days). Four genotypes out of all the ones being studied were discovered earlier than the check cultivar Pant Samrat (60.33 days) for days to 50% flowering. The results were represented in Table 2. Similar outcomes had previously been reported by Muniappan et al., [16], Kumar and Anumugam [5], Lokesh [13], Mili et al., [10] and Solaimana et al., [11] in brinjal. Number of clusters per plant is a major yield contributing character. Significantly higher number of clusters per plant were observed in the genotype DBR-8 (13.30) followed by CH-1045 (12.67) and Kavya (10.70). Seven genotypes out of the total under study were found to have more clusters per plant than the control cultivar Pant Samrat (7.83). Additionally, Madhavi et al. [7] and Tripathy et al. [17] reported earlier on substantial variations in the number of clusters per plant. Number of flowers per cluster is a major trait contributing

towards yield. More number of flowers in a cluster leads to more fruits which ultimately results in high production. Significantly higher number of flowers per cluster was observed in the genotype NBJ-19 (6.70) followed by Kavya (5.53) and IVBL-116-131 (4.87). Six genotypes out of the total under study were found to have more blooms per cluster than the control cultivar Pant Samrat (3.58). Also previously documented were significant changes in the number of clusters per plant by Madhavi et al. [7], Yadav et al. [8], and Ravali et al. [18].

3.2.4 Fruit parameters

Number of fruits per cluster is a major trait contributing yield. More number of flowers in a cluster leads to more fruits which ultimately results in high production. Significantly higher number of fruits per cluster was observed in the genotype NBJ-19 (3.87) followed by IVBL-116-131 (2.87) and Kavya (2.57). Amongst all the genotypes under study, three genotypes were found superior than check cultivar Pant Samrat (2.42) for number of fruits per cluster. Wide variations for number of clusters per plant were also reported earlier by Ravali et al., [18] and Tripathy et al., [17]. Fruit setting percentage determines the number of fruits per cluster. Brinjal has heterostyly, high fruit setting percentage results in more of flowers to set into fruits hence resulting in high yield. High fruit setting percentage was observed in the genotype BB-58 (86.67%) and it was statistically at par with Swarna Shree (83.33%). Eight genotypes out of the total research population were determined to be superior to the check cultivar Pant Samrat (70.03%) for fruit setting percentage. Wide variations for fruit setting percentage were also reported earlier by Madhavi et al., [7]. One of the main factors affecting output is the number of fruits per plant. This trait directly influences the yield of the plant. Significantly higher number of fruits per plant was observed in the genotype NBJ-19 (24.17) followed by IVBL-116-131 (20.27) and Kavya (18.43). Four genotypes were found to produce more fruits per plant than the check cultivar pant samrat (12.70) out of all those under evaluation. Dhaka and Soni [9], Mili et al. [10], Rahman et al. [14], Madhavi et al. [7], and Yadav et al. [8] previously documented significant differences in the number of fruits per plant in brinjal. The results were represented in Table 3.

Table 1. Analysis of variance (mean sum of squares) for fruit yield and its component characters in brinjal

Source of Variation	d. f.	Plant Height (cm)	Number of branches per plant	Stem girth (cm)	Leaf area (cm ²)	Days to 50 % flowering	Number of clusters per plant	Number of flowers per cluster	Number of fruits per cluster	Fruit Setting Percentage (%)	Number of fruits per plant	Average fruit weight (g)	Length of Fruit (cm)	Diameter of fruit (cm)	Yield per Plant (kg)	Yield per plot (kg)	Shelf life (Days)	Total soluble solids (°Brix)	Ascorbic acid (mg/100 g)
Replication	2	41.02	0.18	0.0015	15.44	31.37	0.21	0.01	0.01	13.37	0.06	68.96	0.15	0.65	0.0005	0.48	0.92	0.09	0.45
Treatment	26	570.50**	4.83**	0.52**	290.04**	293.60**	23.63**	5.74**	1.30**	645.08**	67.76**	6086.98**	22.33**	6.91**	0.79**	58.08**	51.99**	0.66**	13.14**
Error	52	108.18	0.08	0.0028	57.39	49.15	0.08	0.01	0.0033	5.57	0.19	16.23	2.37	0.62	0.0017	0.15	0.33	0.11	0.08

** Significant at 1% level of significance

Table 2. Mean Performance of brinjal genotypes for different growth and flower characters

Sl No	Genotypes	Plant Height (cm)	No of primary branches per plant	Stem girth (cm)	Leaf area (cm ²)	Days to 50 % Flowering	No of clusters per plant	No of Flowers per cluster
		Mean ± S.E.(m)	Mean ± S.E.(m)	Mean ± S.E.(m)	Mean ± S.E.(m)	Mean ± S.E.(m)	Mean ± S.E.(m)	Mean ± S.E.(m)
1	DRNKV-02-104	82.33 ± 3.20	5.13* ± 0.11	1.42* ± 0.03	92.17 ± 4.55	44.00* ± 0.58	3.75* ± 0.08	2.23* ± 0.01
2	IVBL-116-131	79.60 ± 4.51	7.61* ± 0.17	1.33* ± 0.03	75.49* ± 3.92	66.33 ± 1.45	8.90* ± 0.02	4.87* ± 0.13
3	NBJ-19	55.39* ± 7.90	7.15* ± 0.14	1.16* ± 0.04	81.89 ± 2.84	70.00 ± 2.65	7.64 ± 0.20	6.70* ± 0.14
4	JB-9	52.90* ± 2.27	6.00* ± 0.17	1.05* ± 0.02	90.41 ± 4.68	86.33* ± 3.33	4.35* ± 0.10	2.70* ± 0.06
5	Aruna	90.31* ± 9.02	9.82* ± 0.22	2.09* ± 0.01	84.05 ± 1.32	78.33* ± 6.44	6.10* ± 0.13	1.43* ± 0.03
6	JB-2	65.83 ± 5.73	8.29* ± 0.02	1.24* ± 0.04	79.13* ± 3.20	61.33 ± 4.63	5.90* ± 0.12	1.85* ± 0.05
7	CH-1045	59.43 ± 1.54	7.50* ± 0.17	1.12* ± 0.03	83.29 ± 5.96	88.33* ± 2.33	12.67* ± 0.28	2.22* ± 0.01
8	Swarna Shree	57.70 ± 3.81	7.67* ± 0.17	1.09* ± 0.02	89.93 ± 5.25	71.33 ± 5.49	3.70* ± 0.01	1.50* ± 0.04
9	JB	73.93 ± 4.10	7.73* ± 0.16	1.21* ± 0.03	93.19 ± 2.23	73.00* ± 2.08	7.37* ± 0.19	1.67* ± 0.04
10	DRNKV-104-43	62.35 ± 6.02	5.82* ± 0.16	1.07* ± 0.02	92.59 ± 3.43	75.67* ± 2.60	4.43* ± 0.10	1.83* ± 0.04
11	RCMBL-04-04	103.60* ± 3.98	8.19* ± 0.18	2.40* ± 0.01	127.21* ± 5.24	64.00 ± 0.58	7.90 ± 0.17	2.40* ± 0.05
12	IBH-2	61.93 ± 2.20	7.62* ± 0.02	1.09* ± 0.03	87.48 ± 3.44	61.00 ± 2.65	9.87* ± 0.20	2.47* ± 0.07
13	Kavya	63.47 ± 3.49	7.25* ± 0.19	1.25* ± 0.03	85.71 ± 4.72	70.33 ± 2.96	10.70* ± 0.29	5.53* ± 0.12
14	B. Dewariya	60.85 ± 2.54	5.60* ± 0.11	2.07* ± 0.06	85.48 ± 5.22	77.67* ± 4.33	6.90* ± 0.18	2.00* ± 0.05
15	DBR-8	63.92 ± 5.00	9.13 ± 0.24	1.40* ± 0.03	93.67 ± 9.88	57.33 ± 4.67	13.30* ± 0.37	2.67* ± 0.06
16	DMU-1	46.25* ± 8.22	6.89* ± 0.19	2.06* ± 0.05	95.21 ± 2.51	85.00* ± 3.46	3.43* ± 0.08	2.00* ± 0.05
17	Azad Brinjal-2	57.10 ± 3.32	6.15* ± 0.16	1.69* ± 0.04	83.04 ± 6.31	62.00 ± 10.69	2.13* ± 0.06	1.60* ± 0.04
18	Co-11	70.42 ± 8.60	8.01* ± 0.17	2.37* ± 0.01	86.65 ± 5.35	67.33 ± 2.73	6.43* ± 0.14	1.82* ± 0.04
19	JB-67	54.10* ± 6.12	6.87* ± 0.14	1.59* ± 0.04	77.16* ± 2.63	57.33 ± 2.96	8.22 ± 0.21	3.93* ± 0.09
20	KKM-1	65.99 ± 7.97	7.44* ± 0.16	1.19* ± 0.03	95.08 ± 3.97	68.67 ± 2.60	6.93* ± 0.15	3.75 ± 0.01
21	BCB-464	43.41* ± 8.65	4.70* ± 0.10	1.19* ± 0.02	82.35 ± 4.61	67.67 ± 5.21	3.38* ± 0.01	1.47* ± 0.04
22	PLR-1	56.57 ± 8.80	6.72* ± 0.14	1.21* ± 0.03	97.21 ± 2.84	74.33* ± 4.91	6.35* ± 0.17	4.77* ± 0.10
23	JB-18	65.00 ± 9.46	9.38 ± 0.26	1.17* ± 0.03	84.00 ± 1.46	73.33* ± 5.78	5.97* ± 0.13	2.73* ± 0.06
24	Bhagyamati	55.12* ± 3.66	7.71* ± 0.17	1.46* ± 0.01	99.91 ± 5.31	56.67 ± 2.33	3.43* ± 0.08	1.63* ± 0.03
25	BB-58	42.28* ± 8.11	7.17* ± 0.02	1.05* ± 0.03	82.33 ± 1.13	71.00 ± 4.04	5.35* ± 0.11	1.67* ± 0.05
26	Pusa Shyamala	52.83* ± 4.37	6.47* ± 0.17	1.37* ± 0.03	86.45 ± 6.56	69.33 ± 3.28	4.85* ± 0.13	2.00* ± 0.05
27	Pant Samrat*	72.51 ± 4.17	9.25 ± 0.20	1.88 ± 0.04	92.50 ± 4.89	60.33 ± 4.06	7.83 ± 0.17	3.58 ± 0.01
	Mean	63.52	7.31	1.45	89.02	68.81	6.58	2.70
	Range	42.28 ~ 103.60	4.70 ~ 9.82	1.05 ~ 2.40	75.49 ~ 127.21	44.00 ~ 88.33	2.13 ~ 13.30	1.43 ~ 6.70
	SE(d)	8.49	0.22	0.04	6.19	5.72	0.23	0.09
	C.D. (0.05)	17.09	0.45	0.09	12.45	11.52	0.46	0.18
	C.V.	16.37	3.76	3.67	8.51	10.19	4.24	4.11

Table 3. Mean performance of brinjal genotypes for different fruit characters

SI No	Genotypes	Number of Fruits per cluster	Fruit Setting Percentage (%)	Number of Fruits per plant	Average fruit weight (g)	Length of Fruit (cm)	Diameter of Fruit (cm)
		Mean \pm S.E.(m)	Mean \pm S.E.(m)	Mean \pm S.E.(m)	Mean \pm S.E.(m)	Mean \pm S.E.(m)	Mean \pm S.E.(m)
1	DRNKV-02-104	1.33* \pm 0.04	58.92* \pm 1.29	10.67* \pm 0.23	177.41* \pm 3.59	17.76 \pm 0.39	3.68 \pm 0.49
2	IVBL-116-131	2.87* \pm 0.06	57.70* \pm 1.26	20.27* \pm 0.41	52.00* \pm 1.44	20.43* \pm 1.04	2.04 \pm 0.24
3	NBJ-19	3.87* \pm 0.08	59.29* \pm 1.20	24.17* \pm 0.67	84.14* \pm 1.85	13.53* \pm 0.73	2.96 \pm 0.19
4	JB-9	1.65* \pm 0.04	62.09* \pm 1.72	10.90* \pm 0.24	89.12* \pm 0.26	11.33* \pm 0.99	4.67* \pm 0.36
5	Aruna	1.00* \pm 0.03	75.64* \pm 1.66	4.97* \pm 0.02	226.67* \pm 5.89	12.45* \pm 1.80	9.62* \pm 1.63
6	JB-2	1.23* \pm 0.03	67.09 \pm 0.19	9.53* \pm 0.25	221.33* \pm 4.86	14.43* \pm 1.36	5.19* \pm 0.29
7	CH-1045	1.30* \pm 0.04	58.33* \pm 1.28	10.40* \pm 0.23	64.90* \pm 1.31	12.30* \pm 1.15	4.11* \pm 0.36
8	Swarna Shree	1.17* \pm 0.03	83.33* \pm 1.83	7.20* \pm 0.14	65.82* \pm 1.82	10.20* \pm 0.32	3.67 \pm 0.26
9	JB	1.07* \pm 0.02	80.00* \pm 1.62	7.83* \pm 0.22	107.73* \pm 2.36	11.96* \pm 0.70	4.97* \pm 0.62
10	DRNKV-104-43	1.40* \pm 0.03	76.11* \pm 2.11	9.33* \pm 0.21	53.41* \pm 0.15	15.13 \pm 1.08	2.30 \pm 0.18
11	RCMBL-04-04	1.07* \pm 0.03	45.96* \pm 1.01	6.07* \pm 0.02	146.53* \pm 3.81	14.25* \pm 0.42	6.97* \pm 0.83
12	IBH-2	1.40* \pm 0.03	56.11* \pm 0.16	8.23* \pm 0.21	80.55* \pm 1.77	11.93* \pm 0.87	4.95* \pm 0.11
13	Kavya	2.57* \pm 0.01	46.44* \pm 1.21	18.43* \pm 0.40	78.72 \pm 1.73	9.87* \pm 0.56	4.29* \pm 0.13
14	B. Dewariya	1.30* \pm 0.03	66.48 \pm 1.35	6.30* \pm 0.17	93.28* \pm 2.04	10.64* \pm 0.94	5.07* \pm 0.35
15	DBR-8	1.40* \pm 0.01	56.11* \pm 1.46	14.83* \pm 0.33	109.31* \pm 2.40	10.45* \pm 0.74	4.65* \pm 0.24
16	DMU-1	1.33* \pm 0.03	66.43 \pm 1.84	6.53* \pm 0.14	64.68* \pm 0.19	9.71* \pm 0.88	3.72 \pm 0.17
17	Azad Brinjal-2	1.03* \pm 0.00	68.64 \pm 1.78	7.10* \pm 0.16	80.87* \pm 1.78	9.52* \pm 1.01	4.29* \pm 0.40
18	Co-11	1.22* \pm 0.04	68.26 \pm 1.50	11.57* \pm 0.03	95.22* \pm 2.47	13.27* \pm 0.49	3.43 \pm 0.22
19	JB-67	1.50* \pm 0.04	37.64* \pm 0.76	6.60* \pm 0.19	87.31* \pm 1.92	11.20* \pm 0.60	4.21* \pm 0.16
20	KKM-1	1.60* \pm 0.04	43.54* \pm 0.95	6.67* \pm 0.15	92.13* \pm 1.86	11.93* \pm 0.74	3.66 \pm 0.35
21	BCB-464	1.07* \pm 0.02	78.57* \pm 1.73	5.53* \pm 0.11	125.36* \pm 3.47	14.07* \pm 0.65	4.45* \pm 0.39
22	PLR-1	1.27* \pm 0.03	26.35* \pm 0.53	5.47* \pm 0.15	78.94 \pm 1.73	9.92* \pm 0.55	3.91* \pm 0.38
23	JB-18	1.22* \pm 0.03	45.65* \pm 1.26	10.47* \pm 0.23	85.20* \pm 0.25	12.49* \pm 0.82	3.70 \pm 0.37
24	Bhagyamati	1.10* \pm 0.03	70.83 \pm 1.55	5.80* \pm 0.02	83.53* \pm 2.17	9.41* \pm 0.40	4.98* \pm 0.23
25	BB-58	1.40* \pm 0.03	86.67* \pm 0.25	11.77* \pm 0.31	106.38* \pm 2.34	12.85* \pm 0.39	3.65 \pm 0.06
26	Pusa Shyamala	1.40* \pm 0.01	75.00* \pm 1.95	10.00* \pm 0.22	63.00* \pm 1.38	14.44* \pm 1.18	2.29 \pm 0.09
27	Pant Samrat*	2.42 \pm 0.06	70.03 \pm 1.54	12.70 \pm 0.28	73.11 \pm 1.48	17.55* \pm 0.98	2.46 \pm 0.11
	Mean	1.53	62.49	9.98	99.51	12.70	4.22
	Range	1.00~3.87	26.35~86.67	4.97 ~ 24.17	52.00 ~ 226.67	9.41 ~ 20.43	2.04 ~9.62
	SE(d)	0.05	1.93	0.36	3.29	1.26	0.64
	C.D. (0.05)	0.09	3.88	0.73	6.62	2.53	1.29
	C.V.	3.76	3.78	4.42	4.05	12.12	18.68

* Significant at 5 % level of significance compared with check cultivar (Pant Samrat) 3.2.5 Yield parameters

Table 4. Mean performance of brinjal genotypes for different yield and quality characters

SI No	Genotypes	Yield per Plant (kg)	Yield per plot (kg)	Shelf life (Days)	Total soluble solids (°Brix)	Ascorbic acid (mg/100 g)
		Mean ± S.E.(m)	Mean ± S.E.(m)	Mean ± S.E.(m)	Mean ± S.E.(m)	Mean ± S.E.(m)
1	DRNKV-02-104	2.02* ± 0.01	18.82* ± 0.49	13.00* ± 0.28	4.20 ± 0.12	10.19* ± 0.21
2	IVBL-116-131	1.07* ± 0.03	10.97* ± 0.24	14.00* ± 0.31	4.40 ± 0.31	8.91* ± 0.25
3	NBJ-19	2.09* ± 0.05	18.72* ± 0.41	7.67* ± 0.16	5.60* ± 0.31	6.37* ± 0.14
4	JB-9	0.96 ± 0.02	9.68 ± 0.20	10.00* ± 0.28	4.50 ± 0.25	11.46* ± 0.04
5	Aruna	1.20* ± 0.02	12.07* ± 0.34	20.33* ± 0.45	5.97* ± 0.03	6.37* ± 0.17
6	JB-2	2.12* ± 0.06	19.79* ± 0.43	18.67 ± 0.06	5.43* ± 0.32	6.37* ± 0.14
7	CH-1045	0.68* ± 0.00	7.44* ± 0.19	13.67* ± 0.30	4.03 ± 0.03	5.09* ± 0.10
8	Swarna Shree	0.48* ± 0.01	5.46* ± 0.12	16.00* ± 0.35	4.67 ± 0.33	6.37* ± 0.18
9	JB	0.84* ± 0.02	8.66* ± 0.19	17.00* ± 0.34	5.17* ± 0.03	5.09* ± 0.11
10	DRNKV-104-43	0.48* ± 0.01	5.44* ± 0.11	12.67* ± 0.35	5.43* ± 0.07	5.09* ± 0.02
11	RCMBL-04-04	0.90 ± 0.02	9.22 ± 0.26	21.00* ± 0.46	4.87* ± 0.09	6.37* ± 0.17
12	IBH-2	0.68* ± 0.02	7.16* ± 0.16	15.00* ± 0.04	5.10* ± 0.15	8.91* ± 0.20
13	Kavya	1.44* ± 0.03	14.02* ± 0.04	18.33* ± 0.48	4.90* ± 0.06	7.64* ± 0.17
14	B. Dewariya	0.60* ± 0.02	6.50* ± 0.14	17.67* ± 0.36	4.90* ± 0.00	10.19* ± 0.22
15	DBR-8	1.63* ± 0.04	15.81* ± 0.05	12.33* ± 0.32	4.40 ± 0.35	6.37* ± 0.14
16	DMU-1	0.43* ± 0.01	5.23* ± 0.11	6.33* ± 0.18	4.40 ± 0.25	8.91* ± 0.03
17	Azad Brinjal-2	0.55* ± 0.01	6.11* ± 0.02	15.33* ± 0.40	4.80* ± 0.00	8.91* ± 0.20
18	Co-11	1.20* ± 0.02	11.86* ± 0.33	18.00* ± 0.39	4.77* ± 0.03	11.46* ± 0.30
19	JB-67	0.60* ± 0.02	6.64* ± 0.15	9.33* ± 0.19	4.88* ± 0.06	10.19* ± 0.22
20	KKM-1	0.61* ± 0.00	6.62* ± 0.17	18.67 ± 0.41	4.87* ± 0.07	7.64* ± 0.16
21	BCB-464	0.68* ± 0.02	7.40* ± 0.16	16.33* ± 0.36	4.80* ± 0.00	8.91* ± 0.25
22	PLR-1	0.45* ± 0.01	5.08* ± 0.11	19.00 ± 0.38	4.93* ± 0.07	6.37* ± 0.14
23	JB-18	0.89 ± 0.02	9.07* ± 0.18	21.67* ± 0.60	4.60 ± 0.31	6.37* ± 0.02
24	Bhagyamati	0.48* ± 0.01	6.10* ± 0.17	17.00* ± 0.38	4.40 ± 0.31	8.91* ± 0.23
25	BB-58	1.32* ± 0.04	12.98* ± 0.29	12.00* ± 0.04	4.27 ± 0.18	8.91* ± 0.20
26	Pusa Shyamala	0.65* ± 0.02	6.93* ± 0.02	9.67* ± 0.25	4.27 ± 0.18	8.91* ± 0.20
27	Pant Samrat*	0.94 ± 0.00	9.71 ± 0.25	19.33 ± 0.42	4.20 ± 0.12	12.73 ± 0.26
	Mean	0.96	9.76	15.19	4.77	8.11
	Range	0.43~2.12	5.08~19.79	6.33 ~ 21.67	4.03 ~ 5.97	5.09 ~ 12.73
	SE(d)	0.03	0.31	0.47	0.27	0.24
	C.D. (0.05)	0.07	0.63	0.95	0.54	0.47
	C.V.	4.22	3.93	3.79	6.95	3.54

* Significant at 5 % level of significance compared with check cultivar (Pant Samrat)

Fruit weight directly affects yield, and consumers are drawn to this quality. Maximum average fruit weight was recorded in Aruna (226.67 g) and it was found statistically at par with JB-2 (221.33 g). The results were represented in Table 3. Amongst all the genotypes under study, twenty genotypes were found superior over check cultivar (73.11 g) for average fruit weight. Significant variations in average fruit weight were also noted previously by Senapati et al. (2014), Rahman et al., [14], Solaimana et al., [11], Madhavi et al., [7], Vidhya and Kumar [19] and Tripathy et al., [17] in brinjal. Fruit length is a crucial factor that directly affects fruit weight and, in turn, the overall yield. Maximum fruit length was recorded IVBL-116-131 (20.43 cm) followed by DRNKV-02-104 (17.76 cm) and Pant Samrat (17.55 cm). Two genotypes viz, DRNKV-02-104 (17.76 cm) and IVBL-116-131 (20.43 cm) had higher fruit length than check cultivar Pant Samrat (17.55 cm). The results of present studies are in line with Ansari et al., [20], Madhvi et al., [7], Yadav et al., [8], Ravali et al., [18] and Tripathy et al., [17] for the trait under study. Fruit diameter directly impacted fruit weight, which in turn affected overall fruit yield. Maximum fruit breadth was observed in Aruna (9.62 cm) followed by RCMBL-04-04 (6.97 cm) and JB-2 (5.19 cm). All the genotypes were found superior than check cultivar Pant Samrat (2.46 cm) except three genotypes viz, IVBL-116-131 (2.04 cm), DRNKV-104-43 (2.30 cm) and Pusa Shyamala (2.29 cm). Madhavi et al., [7], Ravali et al., [18] and Pujer et al., [21] had also reported significant variations for fruit breadth in brinjal.

The main focus of cultivating a crop is to have the maximum yield per unit area for better returns. Moreover, high fruit yield is the ultimate goal of any breeding programme, hence it requires the highest consideration. The genotype JB-2 had the highest yield (2.12 kg), which was statistically equal to NBJ-19 (2.09 kg). The outcomes were shown in Table 4. Ten genotypes out of the total under research were determined to be superior to the control cultivar Pant Samrat (0.94 kg). Tremendous variations regarding yield parameter in brinjal have also been reported earlier by Dhaka and Soni [9], Rahman et al., [14], Solaimana et al., [11] and Pujer et al., [21] in brinjal. A rough notion of the yield that can be obtained over a broad area is provided by yield per plot. Maximising yield per unit area is the major goal of crop cultivation in order to maximise returns. High fruit output is also the ultimate goal of any breeding programme, thus it demands the most care. Maximum yield was

recorded in the genotype JB-2 (19.79 kg) followed by DRNKV-02-104 (18.82 kg) and NBJ-19 (18.72 kg). Nine genotypes out of the total under research were determined to be superior to the control cultivar Pant Samrat (9.71 kg). In the past, Shekar et al. [22], Karak et al. [12], Lokesh [13], and Madhavi et al. [7] observed enormous fluctuations in the yield parameter in brinjal.

3.2.5 Quality parameters

Fruit quality is significantly influenced by shelf life. Fruits with a longer shelf life can be shipped to far-off markets, whereas fruits with a shorter shelf life are more susceptible to damage from long distance shipping and illness. Maximum shelf life was observed in the genotype JB-18 (21.67 days) and it was found statistically at par with RCMBL-04-04 (21.00 days). Three genotypes viz, Aruna (20.33 days), RCMBL-04-04 (21.00 days) and JB-18 (21.67 days) were found superior for shelf life than check cultivar Pant Samrat (19.33 days). Earlier workers like Sharma et al., [23,24] and Mili et al., [10] have also observed similar results for shelf life in brinjal. The data for total soluble solid revealed notable variations across all genotypes. Maximum TSS was observed in Aruna (5.97 °Brix) and it was found statistically at par with JB-2 (5.43 °Brix) and NBJ-19 (5.60 °Brix). Whereas, minimum TSS was observed in CH-1045 (4.03 °Brix). All the genotypes were superior than the check cultivar Pant Samrat (4.20 °Brix) except CH-1045 (4.03 °Brix) and DRNKV-02-104 was same as check cultivar for this trait. Similar results had also reported by Koundinya et al., [25]. Ascorbic acid content is important quality parameters of brinjal. Maximum ascorbic acid was observed in Pant Samrat (12.73 mg/100 g) followed by JB-9 and CO-11 (11.46 mg/100 g each). In the meanwhile, minimum ascorbic acid was recorded in CH-1045, JB and DRNKV-104-43 (5.09 mg/100g each). A higher ascorbic acid level than the population mean was present in fourteen genotypes, including the check cultivar. Ravali et al., [18] and Pujer et al., [20] had also reported wide variability ascorbic acid content in different genotype of brinjal [26,27]. The results were represented in Table 4.

4. CONCLUSION

According to the results of the investigation, seven genotypes viz, Aruna, DRNKV-02-104, NBJ-19, JB-2, RCMBL-04-04, IVBL-116-131, and Kavya (Plate 1) recorded higher fruit yield and




Plate 1. Best performing genotypes for yield and other attributing characters

also outperformed check cultivar Pant Samrat for other yield-attributing traits. These genotypes require additional testing before being made available as an alternative to currently available brinjal varieties or they can be used in future breeding programmes to create superior varieties or hybrids for improving yield and quality in brinjal grown in Uttarakhand hilly regions.

CONFERENCE DISCLAIMER

Some part of this manuscript was previously presented in the conference: 6th International Conference on Strategies and Challenges in Agricultural and Life Science for Food Security and Sustainable Environment (SCALFE-2023) on April 28-30, 2023 in Himachal Pradesh

University, Summer Hill, Shimla, HP, India. Web Link of the proceeding: <https://www.shobhituniversity.ac.in/pdf/Souvenir-Abstract%20Book-Shimla-HPU-SCALFE-2023.pdf>

ACKNOWLEDGEMENT

The authors cordially thank, The Director, Indian Institute of Vegetable Research (IIVR), Varanasi, for furnishing the seed materials of different genotypes used in the study through the Material Transfer Agreement.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Chaudhary B. Vegetables, National Book Trust. New Delhi, 4th Edition, New Delhi. 2015;50-58.
2. Anonymmous. First advance estimate. Department of Agriculture and Farmers Welfare. Government of India; 2022-23. Available:<https://agricoop.nic.in/en/StatHortEst>
3. Bisht AS, Sharma KD. Plant utilization by the communities of bharsar and adjoining area of pauri garhwal district, Uttarakhand, India. Biodiversitas. 2014;15(1):94-100.
4. Ranganna S. Handbook of analysis and quality control for fruit and vegetable products. 2nd ed. Tata McGraw Hill, New Delhi. 2015;105-106.
5. Kumar RS, Anumugam T. Genetic variability and genetic advance for fruit yield quality and pest disease incidence in eggplants (*Solanum melongena* L.). Vegetables Science. 2013;40(1):111-113.
6. Nayak BR, Nagre PK. Genetic variability and correlation studies in brinjal (*Solanum melongena* L.). International Journal of Applied Biology and Pharmaceutical Technology. 2013;4(1).
7. Madhavi N, Mishra AC, Prasad O, Bahuguna N. Studies on variability, heritability and genetic advance in brinjal (*Solanum melongena* L.). Plant Archives. 2015;15(1):277-281.
8. Yadav N, Dhankar SK, Chandanshive AV, Kumar V. Studies on variability, heritability and genetic advance in brinjal (*Solanum melongena* L.). The Bioscan. 2016;11(4): 3001-3005.
9. Dhaka SK, Soni AK. Genetic variability in brinjal (*Solanum melongena* L.). Asian Journal of Horticulture. 2014;7(2):537-540.
10. Mili C, Bora GC, Das B, Paul SK. Studies on variability, heritability and genetic advance in *Solanum melongena* L. (Brinjal) genotypes. Direct Research Journal of Agriculture and Food Science. 2014;2(11): 192-194.
11. Solaimana AHM, Nishizawa T, Khatuna M, Shahabuddin A. Physio Morphological characterization genetic variability and correlation studies in brinjal genotypes in Bangladesh. Computational and Mathematical Biology. 2015;4(1):1-36.
12. Karak C, Ray U, Akhtar S, Naik A, Hazra P. Genetic variation and character association in fruit yield components and quality character in brinjal (*Solanum melongena* L.). Journal of Crop and Weed. 2012;8(1):86-89.
13. Lokesh B, Reddy PS, Reddy RVSK, Sivaraj N. Variability, heritability and genetic advance studies in Brinjal (*Solanum melongena* L.). Electronic Journal of Plant Breeding. 2013;4(1):1097-1100.
14. Rahman MO, Rabbani MG, Yesmin R, Garvey, EJ. Genetic diversity of brinjal (*Solanum melongena* L.) through multivariate analysis. International Journal of Natural and Social Sciences. 2014;1:85-93.
15. Kushwah S, Bandhyopadhy BB, Sharma RN. Genetic divergence in brinjal. Haryana Journal of Horticultural Sciences. 2005;34(3-4):316-317.
16. Muniappan S, Saravanan K, Ramya B. Studies on genetic divergence and variability for certain economic characters in eggplant (*Solanum melongena* L.). Electronic Journal of Plant Breeding. 2010;1(4):462-465.
17. Tripathy B, Sharma D, Jangde BP, Bairwa PL. Genetic variability and heritability studies in brinjal (*Solanum melongena* L.). The Bioscan. 2017;10:109-116.
18. Ravali B, Reddy KR, Saidaiah P, Shivraj N. Variability, Heritability and Genetic Advance in Brinjal (*Solanum melongena* L.). International Journal of Current Microbiology and Applied Sciences. 2017; 6(6):42-47.
19. Vidhya C, Kumar N. Genetic variability studies in Brinjal (*Solanum melongena* L.) for fruit yield and quality. Electronic

- Journal of Plant Breeding. 2015;6(3):668-671.
20. Ansari SF, Mehta N, Ansari S, Gavel GP. Variability studies in brinjal (*Solanum melongena* L.) in Chhattisgarh plains. Electronic Journal of Plant Breeding. 2011;2(2):275-281.
 21. Pujer P, Jagadeesha RC, Cholin S. Genetic variability, Heritability and Genetic Advance for Yield, Yield Related Components of Brinjal [*Solanum melongena* (L.)] Genotypes. International Journal of Pure and Applied Bioscience. 2017;5(5):872-878.
 22. Shekar KC, Ashok P, Sasikala K. Studies on heritability and multivariate analyses in brinjal (*Solanum melongena* L.). Vegetable Crops Research Bulletin. 2012;76:79-88.
 23. Sharma SK, Talukdar P, Barbora MC. Genetic divergence in brinjal. Annals of Biology. 2000;16(1):67-70.
 24. Sharma TVRS, Swaroop K. Genetic variability and character association in brinjal (*Solanum melongena* L.). Indian Journal of Horticulture. 2000;57(1):59-65.
 25. Koundinya AVV, Das A, Layek S, Chowdhury R, Pandit MK. Genetic variability, characters association and path analysis for yield and fruit quality components in Brinjal. Journal of Applied and Natural Science. 2017;9(3):1343-1349.
 26. De Candolle AP. Origin des plantes, Paris. 1883;150.
 27. Vavilov NI. Proceedings of 5th International Congress of Genetics, New York. 1928;342-369.

© 2023 Harish et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104649>