



# **Effect of Nutrients Omission on Growth, Yield and Economics of Rabi Sorghum (*Sorghum bicolor* L.) in Vertisols under Rainfed Conditions**

**C. Sudha Rani <sup>a++</sup>, K. Sandya Rani <sup>b#</sup>, K. Shekar <sup>a#</sup>  
and C. Sudhakar <sup>a++</sup>**

<sup>a</sup> Agricultural Research Station, Tandur, Jayashankar Telangana State Agricultural University, Rajendranagar, Telangana, India.

<sup>b</sup> Agricultural Research Station, Kampasagar, Jayashankar Telangana State Agricultural University, Rajendranagar, Telangana, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/IJECC/2023/v13i123773

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110944>

**Original Research Article**

**Received: 17/10/2023**

**Accepted: 21/12/2023**

**Published: 26/12/2023**

## **ABSTRACT**

A field experiment was conducted at Agricultural Research Station, Tandur, Professor Jayashankar Telangana State Agricultural University during 2020-21 and 2021-22 to study the impact of nutrient omission on productivity and profitability of Rabi sorghum. The experiment was laid out in randomized block design with 3 replications. Pooled data results indicated that among the nutrient omissions treatments, significantly higher plant height (211.7 cm), days to 50% flowering (75 days), panicle length (17.9 cm), number of grains panicle<sup>-1</sup> (946), grain yield (1234 kg ha<sup>-1</sup>), fodder yield

<sup>++</sup> Principal Scientist (Agronomy);

<sup>#</sup> Scientist (Agronomy);

\*Corresponding author: E-mail: ksrgrico@gmail.com;

(5788 kg ha<sup>-1</sup>), harvest index (18.17), net returns of Rs. 44,790 ha<sup>-1</sup> and BC ratio of 2.68 were recorded in no omission treatments and it was on par with K omission followed by N omission and lowest with NPK omission (control) treatments. When compared to the no omission (NPK) treatment, the *rabi* sorghum grain yields were decreased by 2% with the K omission, 10% with N omission, 15% with P omission, 22% with PK omission, 23% with NK omission, 29% with NP omission and 39% with NPK omission.

**Keywords:** Nutrient omission; *rabi* sorghum; vertisols; yield.

## 1. INTRODUCTION

“*Rabi* sorghum is valued mainly as staple food of humans in rainfed ecosystems and fodder for livestock. In India, *Rabi* Sorghum is cultivated in an area of 27.35 lakh ha with a production of 28.25 lakh tons and productivity of 1033 kg ha<sup>-1</sup>. In Telangana, this crop occupies an area of 0.48 lakh ha with a production 1.20 lakh tons and an average yield of 1711 kg ha<sup>-1</sup>” [1]. “The production and productivity of this crop is restricted by insufficiency of present fertilizer recommendations, low fertilizer efficiency, no application of micronutrients and continuous deterioration of soil quality. An adequate supply of plant nutrients decides optimum productivity of any crop or cropping system. All other factors of crop production are in the optimum, the fertility of a soil largely determines the ultimate yield. When the soil does not supply sufficient nutrients for normal plant development and optimum productivity, application of supplemental nutrients are required. Since soil reserves alone are not sufficient to meet the nutrient requirement of the crop, a balanced nutrient supply has to be provided keeping in view of the soil inherent nutrient supplying capacity to achieve higher crop yields (Das et al., 2014). Fertilizer is one of the most important sources to meet the target yield. Indiscriminate use of fertilizers may cause adverse effects on soils and crops both nutrient toxicity and deficiency either by over use or inadequate use” [2]. Thus the current investigation was designed to determine the effect of nutrient omission under rainfed conditions on growth and yield parameters of *Rabi* sorghum.

## 2. MATERIALS AND METHODS

A field experiment was conducted at Agriculture Research Station, Tandur, Professor Jayashankar Telangana State Agricultural University for two years 2020-21 and 2021-22. The soil of the experimental field was clayey in texture, low in organic carbon (0.26 %), low in available nitrogen (225 kg ha<sup>-1</sup>), high in available

phosphorus (22 kg ha<sup>-1</sup>), high in available potassium (532 kg ha<sup>-1</sup>) and slightly alkaline in reaction (pH 6.79). The recommended dose of NPK for *Rabi* sorghum was 60:40:40 (NPK kg ha<sup>-1</sup>). An experiment was laid out in randomized block design by comprising eight treatment plots 1. T<sub>1</sub>: No Omission (NPK), T<sub>2</sub>: N omission (PK), T<sub>3</sub>: P omission (NK), T<sub>4</sub>: K omission (NP), T<sub>5</sub>: NP Omission (K), T<sub>6</sub>: NK Omission (P), T<sub>7</sub>: PK omission (N) and T<sub>8</sub>: NPK omission (control) with 3 replications. *Rabi* Sorghum was sown CSV-29 R was with spacing of 45 cm x 10 cm in second fortnight of October and harvested in second fortnight of February for two years. Net plot yield was converted into hectare basis. The results were analyzed using standard statistical procedure given by Panse and Sukhatme [3].

## 3. RESULTS AND DISCUSSION

### 3.1 Growth Parameters

Significantly maximum plant height (211.7 cm) was recorded with No omission (NPK) treatment and it was on par with K omission (NP) followed by N omission (PK) treatment. There was no significant difference in a number of plants m<sup>-2</sup> (Table.1). The increase in the plant height availability of adequate nutrient supply. The results were similar to Joshi et al. [4] reported that growth were significantly influenced by nutrient omission. In line with this result, Sebnie and Mengesha [5] also stated that nitrogen and phosphorus fertilizers significantly affect the plant height of sorghum in the Wag-Lasta area of Ethiopia.

### 3.2 Yield Attributes

Significantly higher panicle length (17.9 cm) and number of grains (946) were recorded without omission (NPK) and it was on par with K omission (17.7 cm) followed by N omission. There was no significance difference in test weight among the treatments (Table.2). The highest panicle length and number of grains panicle<sup>-1</sup> might be due to better photosynthates and translocation of nutrients [6].

**Table 1. Effect of nutrition omission on growth parameters of Rabi Sorghum**

Treatments	Plant height (cm)			No. of plants m <sup>-2</sup>			Days to 50 % flowering		
	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled
T1: No omission (NPK)	213.0	210.3	211.7	18	19	19	75	75	75
T2: N omission (PK)	193.0	190.0	191.5	16	18	17	74	74	74
T3: P omission (NK)	188.7	187.3	188.0	15	17	17	74	74	74
T4: K omission (NP)	208.7	206.0	207.3	17	20	19	74	74	74
T5: NP omission (K)	178.3	176.0	177.2	15	16	16	73	73	73
T6: NK omission (P)	170.3	168.3	169.3	14	16	16	71	71	71
T7: PK omission (N)	174.7	172.3	173.5	15	17	16	71	71	71
T8: NPK omission (Control)	164.7	162.1	163.4	14	16	16	69	69	69
SEm ±	9.55	9.87	9.69	1.15	1.45	1.32	0.55	0.55	0.55
C.D. (p=0.05)	20.69	21.37	21.00	2.50	NS	NS	1.19	1.19	1.19
CV (%)	6.28	6.57	6.41	9.06	10.14	9.57	0.93	0.93	0.93

**Table 2. Effect of nutrition omission on yield attributes of Rabi Sorghum**

Treatments	Panicle length (cm)			No. of grains panicle <sup>-1</sup>			Test wt (gm)		
	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled
T1: No omission (NPK)	17.9	17.8	17.9	940	951	946	42.7	42.8	23.6
T2: N omission (PK)	16.7	16.7	16.7	908	866	887	41.4	41.8	23.0
T3: P omission (NK)	16.4	16.4	16.4	855	916	886	40.7	41.1	22.6
T4: K omission (NP)	17.7	17.7	17.7	917	926	922	43.7	44.0	24.2
T5: NP omission (K)	16.0	16.0	16.0	811	827	819	39.1	39.4	21.7
T6: NK omission (P)	15.7	15.7	15.7	788	803	796	38.4	38.7	21.3
T7: PK omission (N)	15.7	16.1	15.9	792	806	799	35.9	45.9	24.7
T8: NPK omission (Control)	14.1	13.3	13.7	547	558	553	33.3	33.7	18.5
SEm ±	0.67	0.611	0.59	50.07	49.736	47.311	2.84	4.52	1.65
C.D. (p=0.05)	1.45	1.32	1.28	108.44	107.71	102.45	NS	NS	Ns
CV (%)	5.02	4.61	4.44	7.48	7.33	7.02	12.59	13.51	12.77

**Table 3. Effect of nutrition omission on yield of Rabi Sorghum**

Treatments	Grain Yield (kg ha <sup>-1</sup> )			Fodder Yield (kg ha <sup>-1</sup> )			Harvest Index (%)		
	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled
T1: No omission (NPK)	1232	1237	1234	6206	5370	5788	16.76	19.58	18.17
T2: N omission (PK)	1039	1188	1114	5320	5471	5395	16.30	18.12	17.22
T3: P omission (NK)	954	1154	1053	5249	5587	5418	15.41	16.84	16.13
T4: K omission (NP)	1134	1294	1214	5720	5853	5786	16.58	17.44	17.01
T5: NP omission (K)	906	853	880	5029	4109	4569	15.27	17.55	16.41
T6: NK omission (P)	859	1032	946	4461	4505	4483	16.17	18.79	17.48
T7: PK omission (N)	900	1023	961	4768	4851	4810	15.96	16.86	16.41
T8: NPK omission (Control)	797	805	801	3904	3653	3779	16.96	18.22	17.59
SEm ±	66.72	185.00	99.87	366.88	826.97	418.21	-	-	-
C.D. (p=0.05)	144.50	NS	216.28	794.51	NS	905.67	-	-	-
CV (%)	8.35	21.11	11.93	8.84	20.56	10.24	-	-	-

**Table 4. Effect of nutrition omission on economics of Rabi Sorghum**

Treatments	Gross returns (Rs ha <sup>-1</sup> )			Net returns (Rs ha <sup>-1</sup> )			BCR		
	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled
T1: No omission (NPK)	64849	58194	61522	47349	42231	44790	2.71	2.65	2.68
T2: N omission (PK)	54992	47308	51150	38452	30928	34690	2.32	1.90	2.11
T3: P omission (NK)	51747	45076	48411	36697	29193	32944	2.44	1.86	2.15
T4: K omission (NP)	59716	55441	57579	43416	39878	41647	2.67	2.56	2.61
T5: NP omission (K)	49327	42164	45745	35237	27338	31287	2.50	1.84	2.17
T6: NK omission (P)	45687	34870	40279	30347	20363	25355	1.98	1.40	1.69
T7: PK omission (N)	48184	39360	43772	34334	25350	29842	2.48	1.80	2.14
T8: NPK omission (Control)	41568	32576	37072	28678	18149	23414	2.22	1.26	1.74

### 3.3 Grain and Stover Yield

The highest grain yield (1234 kg ha<sup>-1</sup>) and fodder yield (5788 kg ha<sup>-1</sup>) were recorded with no omission treatment (NPK) and it was on par with K omission (NP), N omission (PK) and P omission (NK). Compared to the no omission (NPK) treatment, the rabi sorghum grain yields were decreased 2% with the K omission, 10% with N omission, 15% with P omission, 22% with PK omission, 23% with NK omission, 29% with NP omission and 39% with NPK omission. The highest fodder yield (5788 kg ha<sup>-1</sup>) was recorded with no omission treatment (NPK) and the lowest in NPK omission (Control) treatment (Table.3). Compared to the no omission (NPK) treatment, the Rabi sorghum fodder yields were decreased 0% with the K omission, 6% with P omission, 7% with N omission, 17% with PK omission, 21% with NP omission, 23% with NK omission and 35% with NPK omission. The highest harvest index was also followed the same trend. In comparison with the other treatments, the higher yield loss was recorded in the unfertilized plots and omitted N plots. These results were expected since that they could be the result of poor nutrient supply in the soil, which did not satisfy the N demand of the sorghum crop [7]. Omission of P and K resulted in 12.4 and 17.5% reduction in grain yield, respectively in wheat [8].

### 3.4 Economics

Maximum net returns of Rs. (44,790 ha<sup>-1</sup>) with BCR of 2.68 was recorded in no omission (NPK) treatment (Table.4). The improvement in economic returns was mainly due to higher grain and fodder yields. Hargilas [9] reported that N and P were proved to be the most limiting nutrient in crop production.

## 4. CONCLUSION

Results indicated that the highest growth, yield attributes, yield and economics of Rabi sorghum were recorded with no omission (NPK) and lowest with NPK omission plots (Control). Omission of K followed by N and P has less reduction in all growth parameters and yield. Omission of PK followed by NK and NP has significantly recorded more reductions in all growth parameters and yield.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Indiastat; 2020-21. Available: <https://www.indiastat.com/table/telangana-state/agriculture/selected-state-season-wise-area-production-product/1423628>.
2. Mohammed Azharuddin BR, Bandiwaddar TT, Durgannavar FM, Shaila HM. Effect of nutrient omission on growth and yield of Rabi Sorghum under rainfed and irrigated ecosystems; 2022.
3. Panse VG, Sukhatme PV. Statistical methods for agricultural workers. ICAR, New Delhi. 1967;199-200.
4. Joshi E, Vyas AK, Dass A, Shiva D. Nutrient omission effects on yield, water productivity and profitability of maize (*Zea mays*). Indian Journal of Agronomy. 2016;61(2):204-209.
5. Sebnie W, Mangesha M. Response of nitrogen and phosphorus fertilizer rate for sorghum (*Sorghum bicolor*) production in Wag-Lasta area of Ethiopia. Archives of Agriculture and Environmental Science. 2018;3(2):180-186.
6. Mohammed Azharuddin R, Bandiwaddar TT, Shaila HM. Effect of nutrients omission on yield, nutrient uptake and economics of rabi sorghum in vertisols under rainfed and irrigated conditions. Journal of Cereal Research. 2014;14(Spl - 2):43-48.
7. Habtemariam Teshome, Eyayu Molla, Tesfaye Feyisa. Identification of yield limiting nutrients for Sorghum (*Sorghum bicolor* (L.) Moench) yield, nutrient uptake and use efficiency on vertisols of Raya Kobo District, Northeastern Ethiopia. International Journal of Agronomy. 2023;1-11.
8. Ekta J, Vyas AK, Dass A, Shiva D, Prajapat K. Nutrient omissions effects on growth, yield, water productivity and profitability of wheat (*Triticum aestivum*) in maize (*Zea mays*)-wheat cropping system. Indian Journal of Agricultural Sciences. 2018;88(6):924-30.

9. Hargilas. Response of Nutrient Omission on Growth, Productivity and Profitability of Maize (*Zea mays*)-Wheat (*Triticum aestvum* L) Cropping System in Southern Rajasthan. International Journal of Current Microbiology and Aplplied Science. 2019; 8(10):1522-1532.

---

© 2023 Rani et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/110944>