



Influence of Various Nutrient Sources on Rice (*Oryza sativa* L.) Productivity and Quality under Organic Farming

Janak Raj ^{a*}, P. K. Dubey ^b, A. R. Kaswala ^b, A. P. Italiya ^c
and Krutika Subodh Patel ^a

^a Department of Agronomy, N.M. College of Agriculture, NAU, Navsari, Gujarat, India.

^b Department of Natural Resource Management, ASPEE College of Horticulture, India.

^c Soil and Water Management Research Unit, NAU, Navsari, Gujarat, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2024/v46i42341

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113466>

Original Research Article

Received: 23/12/2023

Accepted: 26/02/2024

Published: 02/03/2024

ABSTRACT

The field investigation aimed to examine the impact of various organic nutrient sources on rice var. GNR-7 at the organic farm of ASPEE College of Horticulture, Navsari Agricultural University, Navsari, Gujarat during the *kharif* season of 2021. The experiment was structured using a Randomized Block Design with a factorial concept, incorporating two levels: soil application and foliar application. Total of twelve treatment i.e., S₁: 100% RDN through NADEP compost, S₂: 80% RDN through NADEP compost along with *Azospirillum* and PSB @ 2 l/ha each S₃: 60% RDN through NADEP compost along with *Azospirillum* and PSB @ 2 l/ha each and S₄: Ghan-jivamrut @ 500 kg/ha + Jivamrut @ 500 l/ha and that of foliar application were, F₀: Control, F₁: Novel organic liquid nutrient @ 1% and F₂: *moringa* leaf extract @ 3%. The application of foliar spray was done thrice at 15, 30 and 45 days after sowing. were replicated thrice. The result revealed that S₁ treatment significantly recorded the grain yield and S₂ treatment significantly recorded the straw

*Corresponding author: Email: 1997janakraj@gmail.com;

yield. While for foliar application, the treatment F₂ significantly recorded the highest grain and straw yield but remained at par with F₁ treatment. The mean data of quality parameters did not significantly change due to soil and foliar application of different organic nutrient sources. It can be determined that the soil application of 100 per cent RDN through NADEP compost and foliar application of either 3 per cent *Moringa* leaf extract or 1 per cent Novel organic liquid nutrient at 15, 30 and 45 DAS resulted in higher yield and also improved the quality parameters of rice under organic farm.

Keywords: NADEP compost; ghan-jivamrut; jivamrut; biofertilizer; rice; novel organic liquid nutrient; *Moringa* leaf extract; organic farming; quality parameters; yield.

1. INTRODUCTION

The escalating population in our country has heightened the demand for food grain, prompting Indian agriculture to embrace green revolution technology. However, the relentless pursuit of increased food grain production has resulted in the excessive use of chemical fertilizers, leading to soil degradation. The imbalanced and excessive application of agricultural chemicals, including fertilizers and synthetic pesticides, has led to soil degradation, thereby compromising the quality of farm produce. Products tainted with unnecessary chemicals pose a potential threat to consumer health. Cultivating crops through eco-friendly methods emerges as a viable solution in addressing these concerns, particularly as the demand for high-quality food continues to surge [1]. Organic farming is a production system that eschews or significantly limits the use of synthetic fertilizers, pesticides, growth regulators, genetically modified organisms, and livestock feed additives. To the fullest extent feasible, organic farming relies on practices such as crop rotations, utilization of crop residues, animal manures, legumes, green manures, off-farm organic wastes, biofertilizers, mechanical cultivation, mineral-bearing rocks, and various forms of biological control to maintain soil productivity and tilth. These natural inputs, readily accessible, facilitate slow nutrient release, supply both macro and micronutrients, and foster a conducive soil environment for microbial populations [2].

Rice (*Oryza sativa* L.) stands as the primary food source for almost half of the world's population, making it a crucial staple in Asian diets. With over 50 percent of the global population relying heavily on rice consumption, its significance extends beyond mere sustenance, impacting economies, employment, cultural practices, and historical narratives [3]. In India, rice holds the status of a staple food for nearly 65 percent of the population. Asia is commonly referred to as

the 'rice bowl' of the world, as it accounts for over 90% of the global production and consumption of rice, making it the premier food crop worldwide. Following China, India holds the position of the second-largest producer of rice globally. In the year 2021-2022, India's rice cultivation covered an area of 46.38 million hectares, yielding a production of 130.29 million tonnes with an average productivity of 2809 kg/ha. The leading rice-producing states within India include West Bengal, Uttar Pradesh, Punjab, Odisha, Andhra Pradesh, Bihar, Chhattisgarh, Tamil Nadu, and Gujarat [4].

In organic farming, composting, particularly the NADEP method, presents an environmentally friendly solution. Cow-based organic manures such as jivamrut, bijamrut, and panchgavya serve as excellent alternatives to chemical fertilizers, playing a pivotal role in organic farming [5]. Biofertilizers containing nitrogen-fixing and phosphate-solubilizing microorganisms enhance soil health by accelerating specific microbial processes, thereby increasing the availability of nutrients in a form readily assimilated by plants [6]. *Moringa* leaf extract, acting as a bio-stimulant with zeatin, efficiently enhances crop growth, reducing reliance on chemical fertilizers [7]. Enriched banana pseudo-stem sap serves as a remedy for the harmful effects of inorganic farming. This value-added product, known as "NOVEL-Liquid Organic Nutrient," developed by the banana pseudo-stem Processing Unit at Navsari Agricultural University, Navsari, Gujarat, contains essential macro and micronutrients, plant growth regulators, and beneficial microbes, thus promoting crop growth and protecting against pests [8]. These organic practices, addressing both the environmental and economic burdens of chemical fertilizers, are in line with the growing global trend towards sustainable agriculture. As awareness grows, integrating these methods can ensure a healthier and more environmentally

friendly future for pulse cultivation, thereby contributing to both food security and sustainable agriculture [9]. Given the importance and anticipated increase in demand for quality food, this study proves invaluable in addressing these challenges.

2. MATERIALS AND METHODS

The field experiment took place at the Organic Farm of ASPEE College of Horticulture, Navsari Agricultural University, Navsari during the *kharif* season of 2021, focusing on rice var. GNR-7. It was structured using a Randomized Block Design with a Factorial concept (FRBD), with two factors: soil application and foliar application replicated thrice. The fourth levels of soil application included: S₁: 100% RDN through NADEP compost, S₂: 80% RDN through NADEP compost along with *Azospirillum* and PSB @ 2 l/ha each S₃: 60% RDN through NADEP compost along with *Azospirillum* and PSB @ 2 l/ha each and S₄: Ghan-jivamrut @ 500 kg/ha + Jivamrut @ 500 l/ha and that of foliar application were, F₀: Control, F₁: novel organic liquid nutrient @ 1% and F₂: *Moringa leaf extract* @ 3%. Foliar spraying was carried out thrice at 15, 30, and 45 days after sowing. Twenty-one days old seedlings were transplanted at the spacing of 20 cm row to row and 15 cm plant-to-plant distance. Two to three seedlings per hill were transplanted. The experimental field's soil had a clayey texture (57.20%) with high organic carbon content (0.77%) and medium levels of available K₂O (495.52 kg/ha), available N (266.8 kg/ha), and available P₂O₅ (60.52 kg/ha). Favorable weather conditions prevailed throughout the cropping period, facilitating optimal crop growth and development.

2.1 Quality Content

Starch content (%) The estimation of starch by anthrone reagent. The sample is treated with 80% alcohol to remove sugars and then starch is extracted with perchloric acid [10.] **Amylose content (%)** A method for the determination of the amylose content in starch has been developed which is based on the colorimetric measurement of the iodine complexes formed with amylose and amylopectin. The method requires measurement at only one wavelength and avoids the use of harsh dispersants for the starch [11]. **Amylopectin content (%)** Amylopectin content was determined as the difference between the total starch content and amylose content and expressed as the percentage of starch. **Amylose-amylopectin ratio:** Amylose-amylopectin ratio is the main factor for classifying rice into waxy and non-waxy. Amylose-amylopectin ratio of rice samples was estimated as per the method suggested [12,13].

2.2 Cooking Quality

Gel consistency (mm) Gel consistency is a measure of the flow characteristics of milled rice (100 mg) in 2.0 ml 0.2N KOH and is indexed by the length of cold horizontal gel in mm in 13 x 100 mm test tube. Gel consistency test together with the amylose test is a good index of cooked rice texture [14]. **Paste clarity (Transmittance %)** The procedures with slight modifications were used for determination of starch clarity. 1% dry basis aqueous dispersions of starch were boiled at 100 °C for 30 min at constant stirring, and then transferred to plastic tubes and stored at 4 °C for 4 weeks. The transmittance was measured at 620 nm using a spectrophotometer every week, obtaining triplicate readings per sample [15].

Table 1. Treatment details

| Factor I: | Soil application (S) |
|------------|---|
| | S ₁ : 100% RDN through NADEP compost |
| | S ₂ : 80% RDN through NADEP compost along with <i>Azospirillum</i> and PSB @ 2 lit/ha each |
| | S ₃ : 60% RDN through NADEP compost along with <i>Azospirillum</i> and PSB @ 2 lit/ha each |
| | S ₄ : Ghan-jivamrut @ 500 kg/ha + Jivamrut @ 500 lit/ha |
| Factor II: | Foliar application (F) |
| | F ₀ : Control |
| | F ₁ : Novel Organic liquid nutrient @ 1% |
| | F ₂ : <i>Moringa leaf extract</i> @ 3% |

3. RESULTS AND DISCUSSION

3.1 Productivity

3.1.1 Grain yield

Based on the results, it was concluded that the grain yield was significantly influenced by the soil application and foliar application of the organic nutrient sources (Table 2). The S₁ 100% RDN through NADEP compost treatment recorded significantly higher grain yield of 3749 kg/ha and it remained at par with S₂ 80% RDN through NADEP compost along with *Azospirillum* and PSB @ 2 l/ha each treatment which recorded 3538 kg/ha grain yield. The result revealed that the foliar application of *Moringa leaf extract* @ 3 per cent viz., F₂ treatment recorded significantly higher grain yield of 3312 kg/ha but was statistically similar with F₁ treatment where, Novel organic liquid nutrient @ 1 per cent was sprayed and it recorded 3257 kg/ha grain yield. The F₀ treatment, which served as the control with no spray applied, recorded the lowest grain yield at 2959 kg/ha. The influence of the interaction effect of soil and foliar application on grain yield was found to be statistically non-significant, with no observed variation in grain yield. Analysis of grain yield data indicated that the combination of soil application of organic nutrient sources and foliar spray of liquid organic sources significantly influenced seed production,

as evident from the substantial variation detected, which positively impacted yield. The higher seed yield attributed to *Moringa leaf extract* may be attributed to its capacity to enhance the loading and unloading of assimilates across membrane boundaries of vascular tissues, thereby increasing yield. Additionally, cytokinins present in *Moringa leaf extract* are known to promote carbohydrate metabolism and establish new source-sink relationships, further contributing to increased crop yield. These findings align with previous studies by Sahare and Mahanpatra [16] in rice, Ghube et al. [17] in organic rice, Jhilik et al. [18] in wheat, Biswas et al. [19] in maize.

3.1.2 Straw yield

The results revealed that the application of S₂ 80% RDN through NADEP compost along with *Azospirillum* and PSB @ 2 l/ha each significantly recorded the highest straw yield i.e. 5442 kg/ha. While the S₄ treatment where Ghan-jivamrut @ 500 kg/ha + Jivamrut @ 500 l/ha was applied recorded the lowest straw yield of 4329 kg/ha. The foliar spray of 3 per cent *Moringa leaf extract* (F₂ treatment) resulted in significantly higher straw yield (5265 kg/ha) and was statistically similar with foliar spray of 1 per cent Novel organic liquid nutrient (F₁ treatment) which recorded 5117 kg/ha straw yield. While the F₀

Table 2. Effect of different organic nutrient sources on yield of rice var. GNR-7

| Treatments | Grain yield (kg/ha) | Straw yield (kg/ha) | Harvest index (%) |
|---|---------------------|---------------------|-------------------|
| Factor I: Soil application (S) | | | |
| S ₁ - 100% RDN through NADEP compost | 3749 | 5437 | 40.9 |
| S ₂ - 80% RDN through NADEP compost along with <i>Azospirillum</i> and PSB @ 2 l/ha each | 3538 | 5442 | 39.4 |
| S ₃ - 60% RDN through NADEP compost along with <i>Azospirillum</i> and PSB @ 2 l/ha each | 2908 | 4774 | 37.9 |
| S ₄ - Ghan-jivamrut @ 500 kg/ha + Jivamrut @ 500 l/ha | 2511 | 4329 | 36.8 |
| SEm± | 96 | 149 | 0.9 |
| CD at 5% | 281 | 437 | 2.8 |
| Factor II: Foliar application (F) | | | |
| F ₀ - Control | 2959 | 4605 | 39.1 |
| F ₁ - Novel Organic liquid nutrient @ 1% | 3257 | 5117 | 38.7 |
| F ₂ - <i>Moringa leaf extract</i> @ 3% | 3312 | 5265 | 38.5 |
| SEm± | 83 | 129 | 0.8 |
| CD at 5% | 243 | 378 | NS |
| Interaction | | | |
| S×F | NS | NS | NS |
| CV (%) | 9.0 | 8.9 | 7.1 |

treatment as control where no foliar application was given recorded the lowest grain yield *i.e.* 4605 kg/ha (Table 2). The statistically non-significant result was obtained for the straw yield due to the interaction effect between the soil and foliar application of various nutrient organic sources. The findings of the present research work were closely related to previous experiment results observed by Parmar et al. [20] in maize, Abusuwar and Abohassan [21] in cereals forages, Safiullah et al. [22] in sweet corn.

3.1.3 Harvest index

The harvest index was calculated based on the grain yield and biological yield, and subsequently subjected to statistical analysis depicted in Table 2. The results revealed that the application of S₁ 100% RDN through NADEP compost significantly recorded the highest harvest index at 40.9%, followed by S₂ and S₃ treatments at 39.4% and 37.9%, respectively. The foliar application of different treatment did not show significant difference in the resulting harvest index. The F₀ treatment (Control) showed the highest numerical harvest index of 39.1%, compared to the F₁ treatment and F₂ treatment, which recorded harvest indices of 38.7% and 38.5%, respectively. The analysis revealed that the harvest index was not significantly affected by the interaction between soil and foliar application of various organic nutrient sources, with the obtained result being statistically non-significant.

3.2 Quality Content

3.2.1 Starch content

The statistically non-significant result was obtained for the starch content. It was noted that S₁ treatment recorded numerically the highest starch content of 76.5% and foliar application of F₂ treatment *Morigna leaf extract* recorded numerically the highest starch content of 76.4% but, slightly lowest starch content observed in S₃ and F₀ (Fig. 1). The interaction effect of soil and foliar application of various organic nutrient sources showed non-significant result for the starch content.

3.2.2 Amylose content

The soil and foliar application of the organic sources failed to influence of the amylose content and the obtained result was statistically

non-significant. Numerically the highest amylose content was recorded the S₂ 80% RDN through NADEP compost along with *Azospirillum* and PSB @ 2 l/ha each treatment *i.e.* 21.4% and foliar application of F₁ *i.e.* Novel Organic liquid nutrient @ 1% (21.5%), but slightly lowest amylose content observed in S₃ and F₀ (Fig. 1). The interaction effect of soil and foliar application of various organic nutrient sources showed non-significant result for the amylose content.

3.2.3 Amylopectin content

The statistically non-significant result was obtained for the amylopectin content. The result showed that the numerically the highest amylopectin content was recorded the S₁ *i.e.* 100% RDN through NADEP compost of 55.2% and foliar application of F₂ *i.e.* *Morigna leaf extract* @ 3% 55.8% but, slightly lowest amylopectin content observed in S₃ and F₀: *i.e.* control. The interaction effect of soil and foliar application of various organic nutrient sources showed non-significant result for the amylopectin content.

3.2.4 Amylose-amylopectin ratio

The soil and foliar application of the organic sources failed to influence of the amylose-amylopectin ratio and the obtained result was statistically non-significant (Fig. 1). The result showed that numerically the highest amylose-amylopectin ratio was recorded the S₂ 80% RDN through NADEP compost along with *Azospirillum* and PSB @ 2 l/ha each treatment *i.e.* 0.40 and foliar application of Novel Organic liquid nutrient @ 1% (F₁) 0.40, but lowest amylose-amylopectin ratio observed in S₄ and F₀ control. The interaction effect of soil and foliar application of various organic nutrient sources showed non-significant result for the amylose-amylopectin ratio.

3.3 Cooking Quality

3.3.1 Gel consistency

The data pertaining to gel consistency (mm) of rice as influenced by different organic nutrient source presented in Fig. 2. The result showed that soil application of S₂ 80% RDN through NADEP compost along with *Azospirillum* and PSB @ 2 l/ha each was recorded numerically higher gel consistency (75.22 mm) and foliar application of F₁ Novel Organic liquid nutrient




Fig. 1. Effect of different organic nutrient sources on starch, amylose, amylopectin content and amylose-amylopectin ratio




Fig. 2. Effect of different organic nutrient sources on gel consistency and paste clarity (Transmittance %)

@1% (75.08 mm), while numerically lowest gel consistency observed in S₄ and F₂. The interaction effect of soil and foliar application of various organic nutrient sources showed non-significant result for the gel consistency.

3.3.2 Paste clarity (Transmittance%)

The data pertaining to paste clarity (transmittance %) of rice as influenced by different organic nutrient source was presented in Fig. 1. The result showed that soil application

of S₁ i.e. 100% RDN through NADEP compost was recorded numerically higher paste clarity 1st week (36.81%), 2nd and 3rd week (27.72%, 7.68%) was recorded with S₄ i.e. Ghan-jivamrut @ 500 kg/ha + Jivamrut @ 500 l/ha and 4th week (4.61%) recorded with S₃: i.e., 60 % RDN through NADEP compost along with *Azospirillum* and PSB @ 2 l/ha each. Foliar application result showed that numerically higher paste clarity 1st week and 4th week (36.77 %, 4.58 %) was recorded with F₂ i.e. *Moringa leaf extract* @ 3%, 2nd week (27.63%) recorded with F₁ i.e. Novel Organic liquid nutrient @ 1% and 3rd week (7.63%) with F₀ i.e. control. The interaction effect of soil and foliar application of various organic nutrient sources showed non-significant result for the paste clarity (Transmittance%).

4. CONCLUSION

Based on the findings of the above study, it is concluded that adopting soil application of 100% RDN through NADEP compost, along with foliar application of either 3% *Moringa leaf extract* or 1% Novel organic liquid nutrient at intervals of 15, 30, and 45 days after sowing, is recommended for achieving higher yields and enhancing quality content for rice cultivation under organic farming practices.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Yadav AK. Organic agriculture (concept, scenario, principals and practices), national centre of organic farming department of agriculture and cooperation, ministry of agriculture, government of India, Ghaziabad, Uttar Pradesh; 2015.
2. Singh RS, Singh D. Effect of organic nutrient management on productivity and economics of rice (*Oryza sativa* L.). Indian J. Agron. 2007;52(3):291-293.
3. Amudha K, Thiyagarajan K, Sakthivel K. Aerobic rice: A review. Agric. Rev. 2009;30(2):145-149.
4. Anonymous. Directorate of economics and statistics, department of agriculture and farmers welfare, ministry of agriculture and farmers welfare government of India, New Delhi; 2022.
5. Kumawat N, Kumar R, Sahu Y. Yield, economics and soil health of mungbean as influenced by organic manures and phosphorus levels. FARM. 2017;1(1):10-13.
6. Agarwal P, Gupta R, Gill IK. Importance of biofertilizers in agriculture biotechnology. Ann. Biol. Res. 2018;9(3):1-3.
7. Abdalla MM. The potential of *Moringa oleifera* extract as a bio-stimulant in enhancing the growth, bio-chemical and hormonal contents in rocket (*Eruca vesicaria*) plants. Int. J. Plant Physiol. Biochem. 2013;5(3):42-49.
8. Desai CS, Patel JM, Pawar SL, Usadadia VP, Naik VR, Savani NG. Value added products from banana pseudostem. Research Scientist, Soil and Water Management Research Unit, Navsari Agricultural University, Navsari. 2016;55-56.
9. Ramprasad V, Srikanthamurthy HS, Ningappa Kakol, Shivakumar, Nagaraju B, Ningaraju, Shashidhara Doddappa, Vijay AR, Shivanna M, Obanna N, Pandu AC, Rama SSM, Veena P. Sustainable Agricultural Practices. Green Foundation Bangalore, First edition, India; 2009.
10. Clegg KM. The application of the anthrone reagent to the estimation of starch in cereals. J. Sci. Food Agric. 1956;7(1):40-44.
11. Sowbhagya CM, Bhattacharya KR. A simplified colorimetric method for determination of amylose content in rice. Starch-Stärke. 1971;23(2):53-56.
12. Mccready RM, Hassid WZ. The separation and quantitative estimation of amylose and amylopectin in potato starch. J. American Chem. Soc. 1943;65(6):1154-1157.
13. Bharti RR, Khichar ML, Godara M, Beniwal G, Duhan R, Jakhar A. Performance of Clusterbean (*Cyamopsis tetragonoloba* L.) Varieties under Different Temperature and Relative Humidity Profiles. International Journal of Plant & Soil Science. 2024 Feb 3;36(3):45-58.
14. Cagampang GB, Perez CM, Juliano BO. A gel consistency test for eating quality of rice. J. Sci. Food Agric. 1973;24(12):1589-1594.
15. Perera C, Hoover R. Influence of hydroxypropylation on retrogradation properties of native, defatted and heat-moisture treated potato starches. Food Chem. 1999;64(3):361-375.
16. Sahare D, Mahapatra A. Effect of organic manures and liquid organic manures on growth, yield and economics of aerobic

- rice cultivation. Int. J. Agric. Sci. 2015;11(1):183-188.
17. Ghube NN, Ghodpage RM, Badole WP, Balpande SS, Nagdeote VG. Soil sustainability and nutritional quality of organically grown rice under farmer's fields of nagpur district. J. soils crops. 2018;28(1):165-171.
 18. Jhilik NZ, Hoque TS, Moslehuddin AZM, Abedin MA. Effect of foliar application of *Moringa leaf extract* on growth and yield of late sown wheat. Asian J. Medical Biol. Res. 2017;3(3):323-329.
 19. Biswas AK, Hoque TS, Abedin MA. Effects of *Moringa leaf extract* on growth and yield of maize. Progressive Agric. 2016;27(2):136-143.
 20. Parmar VP, Shinde RD, Kokani JM, Patel KG, Kolambe BN. Effect of organic nutrient management on maize production under south Gujarat condition. Int. J. Chem. Stud. 2017;5(6): 2067-2069.
 21. Abusuwar AO, Abohassan RA. Effect of *Moringa olifera leaf extract* on growth and productivity of three cereal forages. J. Agric. Sci. 2017;9(7): 236-243.
 22. Safiullah K, Durani A, Durani H, Ansari MA. Effect of solid and liquid organic manures on growth, yield and economics of sweet corn (*Zea mays L.*) under South Gujarat condition. Int. J. Pure App. Biosci. 2018b;6:(2):567-574.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/113466>