



Bovine Renal Lipofuscinosis in a Tanzanian Crossbred Yearling Male Calf: A Case Report

Seleman Nasib Masola ^{a*}

^a Tanzania Livestock Research Institute (TALIRI), Southern Highlands Zone office, P. O. Box-6191, Mbeya, Tanzania.

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: <https://doi.org/10.9734/air/2024/v25i41085>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116593>

Case Report

Received: 01/03/2024

Accepted: 04/05/2024

Published: 18/05/2024

ABSTRACT

Aim: To report a rare case of bovine renal lipofuscinosis (BRL) that was encountered incidentally at necropsy.

Presentation of the Case: In June 2023 an yearling male calf was found dead in one of the dairy cattle farms in Mbeya city, Tanzania; after being sick for about a week. The owner requested a necropsy service to establish the cause of the death of the animal. Necropsy revealed significant gross pathological changes in a liver that was cooked-like and shrunken, with flabby consistence and large pale necrotic foci. Furthermore, dark blue discoloration of both kidneys with normal morphology was evident. Based on the history and gross pathological findings, the animal was diagnosed to have died from acute hepatitis. Differential diagnosis for kidney discoloration included hemachromatosis, renal hemosiderosis, BRL, and melanotic renal cell carcinoma (MRCC). After ruling out hemachromatosis, renal hemosiderosis and MRCC in the list of possible causes of kidney discoloration based on the lack of gross pathological findings, particularly hepatic fibrosis, pallor, and a mass of tumor tissue in the kidneys that are characteristic of hemachromatosis, renal hemosiderosis and MRCC, respectively; the cause of kidney discoloration was diagnosed as BRL.

*Corresponding author: E-mail: selemani.masola@taliri.go.tz, snmasola58@yahoo.com;

Cite as: Masola, S. N. (2024). Bovine Renal Lipofuscinosis in a Tanzanian Crossbred Yearling Male Calf: A Case Report. *Advances in Research*, 25(4), 99–105. <https://doi.org/10.9734/air/2024/v25i41085>

Discussion: In this necropsy, the history and gross pathology suggested that the animal's death could be attributed to acute liver failure, which is one of the sequelae of acute hepatitis. Kidney discoloration was not considered an attribute factor to the animal's death because, with the exception of discoloration, the kidneys did not have gross pathological changes. Moreover, to date there has been no report of a clinical disease in cattle associated with BRL. However, BRL is an important disorder because it leads to condemnation of the affected kidneys as unfit for human consumption.

Conclusion: The case of BRL reported here was an incidental finding at necropsy, and is not associated with the death of the animal.

Keywords: Dark blue discoloration; black kidneys; hemochromatosis; renal hemosiderosis; melanotic renal cell carcinoma.

1. INTRODUCTION

Bovine renal lipofuscinosis (BRL) is an incidental finding in cattle at slaughter [1,2]. It is associated with brown to black kidney discoloration, which is condemned as unfit for human consumption [1,3]. Grossly, the main characteristic is a brownish to dark brown discoloration of the kidneys in an apparently healthy animal [2,4]. The discoloration affects the entire organ and both kidneys are equally affected to the same extent. The extent of discoloration varies. In some instances, it may be so intense that the kidneys appear dark blue [4]. This discoloration has led to the more common name "black kidney disease" [2,4,5].

Histopathology of "black kidneys" demonstrates no renal lesions and a preserved renal parenchyma. For histological sections stained with hematoxylin and eosin (H&E) stain, a granular refringent pigment varying from light yellow to brown or red can be seen within proximal tubules. For histological sections stained with periodic acid Schiff (PAS) stain, pigment varying from deep purple to a light pinkish-purple can be seen within the proximal tubules [2,3].

BRL has been reported as incidental findings in slaughter cattle for more than a century [3]. Initially, some studies identified the accumulation of melanin pigment as the cause of this condition. Others identified the accumulation of hemosiderin as the cause of the condition, and others identified the accumulation of lipofuscin pigment as the cause of the condition [3]. Since the 1960s, several studies have further classified this pigment as lipofuscin based on its staining and ultrastructural properties [2, 3], a view currently generally accepted.

Generally, lipofuscin pigment comes from the degradation of proteins and lipids that

accumulate within cells, especially cells of the heart [2,6-8], liver [2,6], neurons [2, 6-9], kidneys [6], dermal tissue [6,8,10] and skeletal muscle fibers [7] as part of the normal cellular aging process [2,8,10] or in association with multiple diseases or disorders [2].

According to reports from epidemiological and genealogical studies, the disorder has a genetic etiology with simple autosomal recessive inheritance in the Danish Red and Danish Holstein breeds [1,3], but with incomplete penetrance of the genotype in Danish Holsteins [1]. The disorder has been reported in Danish Red, Danish Holstein, crossbred cattle [1,3], Danish Jersey, Danish Red Holstein, Finnish Ayrshire, Norwegian Red, Jutland cattle, Summertaler, Swiss Brown, Grauvieh, Highland cattle, Danish Gelbvieh, Dexter, Salers, Aberdeen Angus, Galloway, Hereford, Piemontese, Blonde d'Aquitaine, Danish Shorthorn, Danish Charolais, Limousine, and Belgian blue breeds of cattle [1]. This paper reports a case of BRL encountered at necropsy of a crossbred yearling male calf in a dairy cattle farm composed of Friesian crosses, in Mbeya city, Tanzania.

2. PRESENTATION OF THE CASE

On 12 June 2023 an yearling male calf was found dead in the morning in one of the dairy cattle farms in Mbeya city, Tanzania. According to the animal attendant, before death the animal was recumbent with decreased appetite for seven consecutive days and was treated with oxytetracycline and multivitamin. Unfortunately, the prognosis was poor, so the animal died. The owner requested a necropsy service to establish the cause of the death of the animal. Necropsy was carried out using standard procedures described previously [11]. Gross pathological findings included a cooked-like shrunken liver

with flabby consistence and large necrotic foci (Fig. 1). Dark blue discoloration of the kidneys was also evident. The discoloration affected the entire kidneys and both kidneys were equally affected to the same extent (Fig. 2 and 3). However, the kidneys had normal gross morphology, that is, normal size and texture (consistence). Based on the history and gross pathological findings, the animal was diagnosed to have died of acute hepatitis.

Differential diagnosis for kidney discoloration included hemochromatosis [12], renal hemosiderosis, renal lipofuscinosis, and melanotic renal cell carcinoma (MRCC) [13]. Renal hemosiderosis is a disease in which hemosiderin is deposited in the renal cortex as a

form of iron overload. It is a complication of chronic intravascular hemolytic states such as hemolytic anemia, paroxysmal nocturnal hemoglobinuria, and mechanical hemolysis from the prosthetic cardiac valve [14]. Hemochromatosis is a disease characterized by an accumulation of iron in the liver and other organs, including the kidneys [15] that leads to tissue damage, particularly the liver [16]. A carcinoma is a type of cancer that forms solid tumors. Like other types of cancer, carcinomas are abnormal cells that divide without control. Carcinomas form in epithelial cells, which line the outer surface of the skin and the cover and lining of organs and internal passageways, such as the gastrointestinal tract [17].




Fig. 1. An image of a cooked-like shrunken liver showing large pale necrotic foci (arrows)



Fig. 2. An image of a right kidney showing dark blue discoloration of the kidney surface



Fig. 3. An image of a left kidney showing dark blue discoloration of the kidney surface

Based on gross pathology, the disorder was diagnosed as renal lipofuscinosis (black kidneys). The diagnosis was made after ruling out renal hemosiderosis, hemochromatosis, and MRCC in the list of possible causes of kidney discoloration based on the absence of gross pathological findings characteristic of each of these diseases. Renal hemosiderosis was ruled out because the dead animal was not anemic; this was indicated by a lack of pallor. Anemia is a consistent characteristic feature of hemosiderosis [14,18-22]. Hemochromatosis was excluded due to the lack of liver fibrosis, a gross pathological finding involving the liver that has been consistently reported in cases of hemochromatosis [12,15,23-27]. MRCC was ruled out because the kidneys were grossly free of tumor tissue, which is a characteristic feature of renal cell carcinoma [28-32] and other types of cancer. Due to budget limitations, histopathology could not be performed to confirm the diagnosis.

3. DISCUSSION

Generally, BRL is a rare disorder in cattle that has been reported to occur incidentally during meat inspection after slaughter of cattle found to be clinically healthy during antemortem inspection [1-4]. Unlike previous reports, this article reports a case of BRL encountered at necropsy which revealed a shrunken, cooked-like liver with flabby consistence and large pale

necrotic foci, and dark blue discoloration of both kidneys. Renal hemosiderosis, hemochromatosis, MRCC, and BRL were considered for differential diagnosis of the dark blue discoloration of the kidneys. Based on gross pathology, the discoloration was diagnosed as BRL after ruling out renal hemosiderosis, hemochromatosis, and MRCC due to lack of pallor, liver fibrosis, and tumor tissue, respectively; which are gross pathological findings characteristic of each of these diseases.

Although in addition to a cooked-like shrunken liver with flabby consistence and large pale necrotic foci, BRL was also evident at necropsy, the pathological changes found in the liver were found to be more significant. Therefore, based on history and gross pathology, the animal was diagnosed to have died of acute hepatitis. The animal's death could be attributed to acute liver failure, which is one of the sequelae of acute hepatitis [33-37]. BRL was not considered an attribute factor to animal death because, except for discoloration, the kidneys did not have pathological changes (had normal gross morphology). In addition to that, to date, there has been no report of a clinical disease in cattle that is associated with BRL.

In this necropsy, both kidneys were found to have dark blue discoloration and equally discolored to the same extent. These findings are

consistent with similar findings previously reported [4]. The normal gross morphology of the discolored kidneys reported here is in agreement with similar incidental findings by Rude et al. [3] at meat inspection where cattle kidneys with dark brown to black discoloration due to the deposition of lipofuscin pigment were found to have normal gross morphology. Furthermore, the finding is in agreement with a previous incidental finding by Lund and Olsen [38] at surgery of a human patient in which the kidneys with diffuse dark brown discoloration due to the deposition of lipofuscin pigment in the epithelium of the proximal tubules were found to have normal gross morphology. These findings suggest that renal lipofuscinosis does not seem to have had any harmful effect on kidney morphology and functions.

BRL is a rare disorder in cattle that has been reported in Holland, Germany, Sweden, Norway [4] and Denmark [3,4]. Tanzania has been importing heifers, bulls and semen from Europe and the United States for breeding purposes [39-41]. Having a genetic etiology, this case of BRL could be associated with the importation of animals and semen from countries affected by BRL. To my knowledge, this is the first report of BRL in Tanzania and the African continent at large.

4. CONCLUSIONS

Based on the gross pathological findings, it is concluded that a case of BRL reported here was an incidental finding at necropsy. The finding is not associated with the death of the animal.

ACKNOWLEDGEMENTS

Mr. First Taison Mwansama is acknowledged for his assistance during necropsy of the carcass.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Agerholm JS, Christensen K, Nielsen SS, Flagstad P. Bovine renal lipofuscinosis: Prevalence, genetics and impact on milk production and weight at slaughter in Danish cattle. *Acta Vet Scand.* 2009;51:7, DOI:10.1186/1751-0147-51-7
2. Slaughterhouse Support Network (SESC). Black kidneys in cattle (renal lipofuscinosis); 2021. Accessed On: 21 June 2023. Available: <https://sesc.cat/en/black-kidneys-in-cattle-renal-lipofuscinosis/>
3. Rude H, Agerholm JS, Maddox-Hyttel P, Christensen K, Flagstad P. Renal lipofuscinosis in Danish slaughter cattle. *J Comp Pathol.* 2005;132(4):303-312.
4. Svenkerud R, Melanosis Renum Bovis. *Acta Vet Scand.* 1960;1:161-187.
5. Winter H. Black kidneys in cattle – A lipofuscinosis. *J PatholBacteriol.* 1963;86(1):253-258.
6. Jung T, Bader N, Grune T. Lipofuscin: Formation, distribution, and metabolic consequences. *Ann N Y AcadSCI;* 2007. DOI:10.1196/annals.1401.008
7. Ottis P, Koppe K, Onisko B, Dynin I, Arzberger T, Kretschmar H, et al. Human and rat brain lipofuscin proteome. *Proteomics.* 2012;12:2445-2454. DOI:10.1002/pmic.201100668
8. Moreno-García A, Kun A, Calero O, Medina M, Calero M. An overview of the role of lipofuscin in age-related neurodegeneration. *Front Neurosci.* 2018;12:464. DOI:10.3389/fnins.2018.00464
9. Kushwaha SS, Patro N, Patro IK. A sequential study of age-related lipofuscin accumulation in hippocampus and striate cortex of rats. *Ann Neurosci.* 2018;25:223-233. DOI:10.1159/000490908
10. Tonolli PN, Baptista MS, Chiarelli-Neto O. Melanin, lipofuscin and the effects of visible light in the skin. *J PhotochemPhotobiol;* 2021. DOI:10.1016/j.jpap.2021.100044
11. Wagner S. Necropsy Techniques in Cattle. In proceedings of the American association of bovine practitioners. 2007;203-204. Accessed On:16 June 2023. Available: <https://www.google.com/search?q=Necropsy+Techniques+in+Cattle>
12. House JK, Smith BP, Maas J, Lane VM, Anderson BC, Graham TW, et al.

- Hemochromatosis in Salers Cattle. *J Vet Intern Med.* 1994;8(2):105-111.
13. Yarmohamadi A, Rezayat ARA, Memar B, Rahimi HR. Normal black kidney. *Can Urol Assoc J.* 2014;8(3-4):e282-6. DOI:10.5489/cuaj.1207
 14. Lee IH, Kang GW, Kim G-Y, Lee S-J, Kim M-K, Ahn DJ. Renal hemosiderosis secondary to intravascular hemolysis after mitral valve repair: A case report. *Med.* 2020;99:3(e18798). DOI:10.1097/MD.00000000000018798
 15. Lavoie JP, Teuscher E. Massive iron overload and liver fibrosis resembling haemochromatosis in a racing pony. *Equine Vet J.* 1993;25(6):552-554.
 16. Frye FL. Iron storage disease (Hemosiderosis) in an African Rock Hyrax (*Procavia capensis*). *J Zoo Anm Med.* 1982;13(4):152-156.
 17. Knox J. Types of carcinoma; 2023. Accessed On: 8 December 2023. Available: <https://www.verywellhealth.com/carcinoma-5092211>
 18. Enyi it A, Ayy Id z O, Babayi it C, Y Id r m M, Büyükbayram H. Idiopathic pulmonary hemosiderosis. *East J Med.* 2001;6(1):29-32.
 19. Asim M, Igbal Z, Mujeeb IB. Blue kidney in a pale patient – a case of a causal association between renal hemosiderosis in paroxysmal nocturnal hemoglobinuria and chronic kidney disease. *NDT Plus.* 2009;2:365-367. DOI:10.1093/ndtplus/sfp057
 20. Tzouveleakis A, Ntoliou P, Oikonomou A, Koutsopoulos A, Sivridis E, Zacharis G, et al. Idiopathic pulmonary hemosiderosis in adults: A case report and review of literature. *Case Rep Med;* 2012. DOI:10.1155/2012/267857
 21. HealthJade. Hemosiderosis. 2019. Accessed On: 29 November 2023. Available: <https://healthjade.net/hemosiderosis/>
 22. Hamilton JPA. Hemosiderosis. 2022. Accessed On: 3 September 2023. Available: <https://www.msmanuals.com/professional/hematology-and-oncology/iron-overload/hemosiderosis>
 23. Madarame H, Kumagai M, Suzuki J, Watanabe A, Konno S. Pathology of excessive iron storage in the Afghan pika (*Ochotona rufescens rufescens*). *J Comp Pathol.* 1990;103:351-359.
 24. Crawshaw G, Oyarzun S, Valdes E, Rose K. Hemochromatosis (Iron Storage Disease) in fruit bats. In *Proceedings of the First Conference on Zoo and Wildlife Nutrition*, AZA Nutrition Advisory Group, Scarborough, OT; 1995. Accessed On: 29 November 2023. Available: <https://nagonline.net/wpcontent/uploads/2014/01/Hemochromatosis-in-bats.pdf>
 25. Bonar CJ, Trupkiewicz JG, Toddes B, Lewandowski AH. Iron Storage Disease in Tapirs. *J Zoo Wildl Med.* 2006;37(1):49-52. DOI:10.1638/03-032.1
 26. Morrice G. Haemochromatosis in two mobs of sheep; 2010. Accessed On: 23 September 2023. Available: <http://www.flockandherd.net.au/sheep/reader/haemochromatosis.html>
 27. Peters A, Raidal SR, Blake AH, Atkinson MM, Atkinson PR, Eggins GP. Haemochromatosis in a Brazilian tapir (*Tapirus terrestris*) in an Australian zoo. *Aust Vet J.* 2012;90:29-33. DOI:10.1111/j.1751-0813.2011.00867.x
 28. Kamishima T, Fukuda T, Emura I, Tanigawa T, Naito M. Pigmented renal cell carcinoma. *Am J Surg Pathol.* 1995;19(3):350-356.
 29. Lei J-Y, Middleton LP, Guo X-D, Duray PH, McWilliams G, Linehan WM, et al. Pigmented renal cell carcinoma with melanocytic differentiation. *Hum Pathol.* 2001;32(2):233-236. DOI:10.1053/hupa.2001.22009
 30. Shetty J, Chandrika, Laxman P. Renal cell carcinoma with melanin pigment. *Indian J Urol.* 2010;26(2):292-293. DOI:10.4103/0970-1591.65407
 31. Antic T, Taxy JB, Alikhan M, Segal J. Melanotic translocation renal cell carcinoma with a novel ARID1B-TFE3 gene fusion. *Am J Surg Pathol.* 2017;00(00):1-5.
 32. Mohd AB, Ghannam RA, Mohd OB, Elayan R, Albakri K, Huneiti N, et al. Etiologies, gross appearance, histopathological patterns, prognosis, and best treatments for subtypes of renal carcinoma: An educational review. *Cureus.* 2022;14(12):e32338. DOI:10.7759/cureus.32338
 33. Davoudi SM. Study of hepatic problems in livestock. *Eur J Zool Res.* 2013;2(4):124-132.
 34. Manka P, Verheyen J, Gerken G, Canbay A. Liver failure due to acute viral hepatitis (A-E). *Visc. Med.* 2016;32:80-85. DOI:10.1159/000444915

35. Daniel C. Common complications of hepatitis; 2022.
Accessed On:15 December 2023.
Available:<https://www.verywellhealth.com/complications-of-hepatitis-1759905>
36. Gong K, Xu X, Yao J, Ye S, Yu X, Tu H, et al. Acute hepatitis of unknown origin in children: A combination of factors. *Front Pharmacol.* 2022;132:1056385. DOI:10.3389/fphar.2022.1056385
37. Jain M. Acute viral hepatitis A infection in adults: Complications and outcomes. *Trop Gastroenterol.* 2023;44(1):42-44.
38. Lund T, Oslen S. Idiopathic Renal Lipofuscinosis. *Acta Pathol Microbiol Scand A.* 1970;74:414-420.
39. USAID Tanzania. An Evaluation of Dairy Production Assistance – Phase I; 1976.
Accessed On:12 February 2024.
Available:<https://pdf.usaid.gov/pdfdocs/pdaab604f1.pdf>
40. United Republic of Tanzania. Country Report: The state of farm animal genetic resources.
Accessed On:21 February 2024.
Available:<https://www.fao.org/3/a1250e/anexes/CountryReports/Tanzania.pdf>
41. World Integrated Trade Solution. Tanzania animal products: Bovine semen imports in 2018. 2024.
Accessed On:12 February 2024.
Available:<https://wits.worldbank.org/trade/comtrade/en/country/TZA/year/2018/tradeflow/Imports/partner/ALL/product/051110>

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/116593>