



Effects of Organic Manures and Bio-Fertilizers on Soil Properties, Productivity and Nutrients Uptake of Indian Mustard

Narender ^{a*}, Mukesh Kumar ^a, Anil Kumar ^a and Manjeet ^a

^a Deendayal Upadhyay Centre of Excellence for Organic Farming, CCS Haryana Agricultural University, Hisar- 125004, Haryana, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i92352

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100157>

Original Research Article

Received: 27/04/2023

Accepted: 28/06/2023

Published: 17/07/2023

ABSTRACT

A field experiment was conducted during the Rabi season 2020-21 to study the effect of organic manures and bio-fertilizers on soil properties, productivity and nutrient uptake of Indian mustard at Deendayal Upadhyay Centre of Excellence for Organic Farming (DDUCE-OF), CCS Haryana Agricultural University, Hisar. The experiment comprised of eight treatments involving Farm Yard Manure (FYM), Vermicompost (VC) and Poultry Manure (PM) alone and along with bio-fertilizers (BF), an absolute control and a cow-based formulation treatment. The results revealed that the treatment 100% RDN through PM+BF produced significantly higher seed and stover yield of mustard which was at par with treatment 100% RDN through VC+BF and 100% RDN through FYM+BF as compared to other treatments. Maximum seed (29.84 q ha⁻¹) and stover (74.75 q ha⁻¹) yields were received under treatment 100% RDN through PM+BF which was 27.5 % and 16.7 % higher over the control, respectively. Similarly, highest total NPK uptake (116.2 kg ha⁻¹ for total N,

*Corresponding author: E-mail: narenderhisar@gmail.com;

19.9 kg ha⁻¹ for total P and 83.5 kg ha⁻¹ for total K) were recorded in the treatment 100% RDN through PM+BF. The highest soil restorative effect on soil organic carbon (0.67 %), available N (162 kg ha⁻¹), available P (13.7 kg ha⁻¹) and available K (280 kg ha⁻¹) were also observed under treatment 100% RDN through PM+BF. These organic amendments such as FYM, vermicompost, poultry manures alone and along with bio-fertilizers, had significantly improved the seed and stover yields as well as uptake of nutrients by Indian mustard.

Keywords: Farm yard manure; vermicompost; poultry manure; mustard; yield; NPK uptake.

1. INTRODUCTION

“Mustard and Rapeseed/Toria are the third most important edible oilseed crops of the world after soybean and oil palm. Globally, India accounts for 21.7% and 10.7% of the total acreage and production. In India, mustard is grown in an area of 7.40 million hectares and the production is 7.70 million tonnes with the productivity of 1040 kilogram per hectare, respectively” [1]. Both mustard and rapeseed are grown in Haryana in an area of up to 6.0-6.5 lac hectares, out of which, mustard is sown in about 90% of the area of the State. Mustard is grown in both the irrigated and rainfed areas and their yield depends on the weather conditions. “The oil content varies from 37 to 49% and oil cake is used as feed and manure. The nutrient management is one of the most important agronomic factors that affects the Indian mustard”, as reported by Hadiyal et al., [2]. “But application of all the needed fertilizer through chemical fertilizers had deleterious effects on soil fertility and unsustainable yields, while organic manures and bio-fertilizers would be able to maintain soil fertility and sustain crop productivity, by improving the physical and nutritional system of the soil and enhance the activity of the soil micro flora” [3].

“Organic cultivation is an environmentally viable approach to agriculture through its use of farm yard manures, compost, green manuring crops and cow based solid and liquid bio-formulations to provide nutrients and cultural practices to manage weeds, insects and pathogens. On the other hand, organic farming has potential benefits in comparison to conventional farming in promoting soil structure formation, enhancing soil biodiversity, protecting environment, improving soil quality, food quality and safety and also, ensuring premium price. Decomposition of organic manures in the soil helps in preventing various disease-causing pathogens by different types of biological reactions” [4]. Thus, organic production systems have the potential to achieve sustainability of agricultural systems. Keeping in

view the importance, the present investigation was taken up to study the effect of organic manures and bio-fertilizers on soil properties, productivity and nutrient uptake of Indian mustard.

2. MATERIALS AND METHODS

The experiment was laid out in Randomized Block Design (RBD) with three replications during the Rabi 2020 season at DDUCE-OF farm, CCSHAU, Hisar. Eight treatments viz. 100% RDN through Farm Yard Manure (FYM), Vermicompost (VC) and Poultry Manure (PM) alone and along with bio-fertilizer (BF) and cow-based bio-formulation were selected. Accounting the recommended dose of nitrogen (80 kg N ha⁻¹), specific amount of well-decomposed FYM, VC and PM were applied to the corresponding treatment plots before sowing of the crop and thoroughly incorporated into the soil. Cow Urine Based Liquid Organic bio-Formulation (CUBLOF) @400 ltr. /ha was applied after 1st and 2nd irrigation each. Mustard variety RH-725 was used as a test crop and treated with Azotobacter + PSB @ 5 ml/kg seed. Mustard seed was sown with 5 kg/ha in lines at a row spacing of 45 cm as per treatment. The depth of seed was kept 4 to 5 cm. After thinning twice, the plant to plant distance was maintained at 10-15 cm. The crop was sown on 17th October, 2020 and was harvested on 2nd April, 2021, respectively. The samples of mustard crop at harvesting stage were collected and washed properly with tap water, distilled water, acidified deionized water and deionized water sequentially. Henceforth, the samples were shade dried followed by drying in the oven at 65 ± 20°C. The seed and stover samples were then ground and stored in paper bags with proper labelling for analysis of various parameters.

“Soil samples were collected from the surface layer (0-15 cm) of all the plots before treatment application and immediately after crop harvest. The pH and Electrical Conductivity (EC) were determined in (1:2) soil:water suspension with

the help of glass electrode pH meter and conductivity meter bridge, respectively using an Eutech pH meter" [5]. Organic Carbon (OC) was estimated by wet digestion method [6]. Soil texture was determined using qualitative methods by feel method given by Soil Science Division Staff [7]. Available nitrogen was determined using steam distillation method [8] and estimated on Gerhardt's fully automated N analyzer (VAPODEST 500 C). Available P was determined by the method of Olsen et al. [9] and estimated on double beam UV spectrophotometer. Available potassium was determined using neutral normal ammonium acetate method [10] and estimated on flame photometer. DTPA-extractable micronutrients in soil samples were determined by Lindsay and Norvell [11] using Atomic Zeenit 700 P Atomic absorption spectrophotometer (Analytik Jena). The data collected on various parameters were analysed statistically according to the experimental design using OPSTAT software, CCSHAU, Hisar.

2.1 Nutrient Contents of Farm Yard Manure (FYM), Vermicompost (VM) and Poultry Manure (PM)

Nutrient content of Farm Yard Manure (FYM), Vermicompost (VM) and Poultry Manure (PM) were analyzed for Total N, P and K content. A set of 8 treatments comprising 100% of recommended nitrogen dose (RDN) given through three sources of organic manures alone (*viz.*, farmyard manure (FYM), vermicompost (VM) and poultry manure (PM) as well as along with biofertilizers. The NPK contents in FYM, VM and PM were 0.65, 1.87 and 2.40% N, 0.40, 0.80 and 1.60% P and 0.85, 0.95 and 1.70% K, respectively.

2.2 Initial Soil Physico-chemical and Microbial Properties of Experimental Site

The soil was loam in texture having pH 7.80 with EC of 0.80 dS m⁻¹. The soil organic carbon (SOC) was observed at medium 0.69%. The available N, P and K contents of soil were 160, 13.5 and 275 kg ha⁻¹, respectively. The status of available N was low (<250 kg ha⁻¹), P was medium and K was high in content (10-20 kg ha⁻¹ for P & >250 kg ha⁻¹ for K). The initial status of DTPA extractable Zn (1.10 mg kg⁻¹), Fe (10.2 mg kg⁻¹), Mn (13.3 mg kg⁻¹) and Cu (1.46 mg kg⁻¹) was observed sufficient in amount. The initial bacterial count of experimental soil was 3.6x10⁸ CFU Count per g of soil.

3. RESULTS AND DISCUSSION

3.1 Effects of Organic Manures and Bio-fertilizers on Seed and Stover yield of Indian Mustard

The moong bean seed yield during kharif season was poor due to white fly infestation and results were found non-significant. However, seed and stover yield of mustard increased significantly with the application of different organic manures alone (*viz.*, farmyard manure (FYM), vermicompost (VM) and poultry manure) and along with bio-fertilizers over control (Table 1). Maximum mustard yields (29.84 q ha⁻¹ in seed and 74.75 q ha⁻¹ in stover) were obtained under treatment 100% RDN through PM+BF. However, minimum seed and stover yields (23.41 & 64.09 q ha⁻¹) were obtained from the control, where there was no application of organic manure. The interaction effect of different organic amendments on seed and stover yield of mustard were found to be significant. Similar findings were observed by Kumar and Singh [12] and reported that significant enhancement in seed (12.7 %) and stover (10.2 %) yields was recorded, respectively with the application of organic manures along with bio-fertilizers over control.

3.2 Effects of Organic Manures and Bio-Fertilizers on NPK Content and Their Uptake by Indian Mustard

The uptake of N, P and K in mustard seeds was from 47.47 to 74.27, 9.43 to 15.13 and 12.80 to 17.63 kg ha⁻¹, respectively under treatment 100% RDN through PM+BF. The corresponding values were from 25.37 to 41.97, 3.03 to 4.77 and 50.77 to 65.80 kg ha⁻¹ in mustard stover (Tables 2, 3 and 4). Treatment 100% RDN through PM+BF produced significantly higher N, P and K content and their uptake in seeds and stover, however, it was statistically at par to treatment 100% RDN through VC+BF and 100% RDN through FYM+BF. Highest Total N, P and K uptake (116.2 kg ha⁻¹, 19.9 kg ha⁻¹ and 83.5 kg ha⁻¹) were recorded in the treatment 100% RDN through PM+BF (Fig. 1). This might be due to proper establishment of roots, higher absorption of mineral nutrients from soil, transport of more nutrients to seeds, vigorous plant growth and higher seed and straw yields under proper availability of nutrients. The overall increase in uptake of nitrogen was the cumulative effect of rise in their concentration in plant tissues and in yield levels. The

interaction effect of organic amendments i.e. FYM, vermicompost and poultry manure alone and along with bio-fertilizers on NPK content and their uptake by Indian mustard were found to be significant and the results are in

conformity with Singh et al. [13] who revealed that all the growth, nutrient uptake of nitrogen, phosphorus and potassium, increased significantly under the treatment of organic manures.

Table 1. Effects of organic manures and bio-fertilizers on seed and stover yield of Indian mustard

Treatment	Seed yield (q ha ⁻¹)	Stover yield (q ha ⁻¹)	Biological yield (q ha ⁻¹)
Control	23.41	64.09	87.50
100% RDN through FYM	25.00	71.89	96.89
100% RDN through VC	26.59	72.02	98.61
100% RDN through PM	27.78	74.49	102.27
100% RDN through FYM+BF	27.46	73.13	100.59
100% RDN through VC+BF	27.78	73.84	101.62
100% RDN through PM+BF	29.84	74.75	104.59
Cow based bioformulation	24.60	72.15	96.75
CD (P=0.05)	2.72	6.19	7.64

Table 2. Effects of organic manures and bio-fertilizers on nitrogen (N) Content and its uptake by Indian mustard

Treatment	N content (%)		N Uptake (kg ha ⁻¹)	
	Seed	Stover	Seed	Stover
Control	2.03	0.40	47.47	25.37
100% RDN through FYM	2.20	0.46	55.10	32.93
100% RDN through VC	2.28	0.49	60.60	35.23
100% RDN through PM	2.30	0.49	63.77	36.70
100% RDN through FYM+BF	2.36	0.51	64.83	37.73
100% RDN through VC+BF	2.38	0.52	66.07	38.60
100% RDN through PM+BF	2.49	0.56	74.27	41.97
Cow based bioformulation	2.14	0.44	52.60	31.47
CD (P=0.05)	0.18	0.07	7.20	5.24




Fig. 1. Effects of organic manures and bio-fertilizers on total N, P, and K contents and their uptake by Indian mustard

Table 3. Effects of organic manures and bio-fertilizers on phosphorous (P) content and its uptake by Indian mustard

Treatment	P content (%)		P uptake (kg ha ⁻¹)	
	Seed	Stover	Seed	Stover
Control	0.40	0.05	9.43	3.03
100% RDN through FYM	0.44	0.05	11.13	3.60
100% RDN through VC	0.45	0.06	12.00	4.00
100% RDN through PM	0.47	0.06	13.10	4.30
100% RDN through FYM+BF	0.48	0.06	13.13	4.43
100% RDN through VC+BF	0.49	0.06	13.50	4.67
100% RDN through PM+BF	0.51	0.06	15.13	4.77
Cow based bioformulation	0.42	0.05	10.27	3.60
CD (P=0.05)	0.05	0.01	1.95	0.42

Table 4. Effects of organic manures and bio-fertilizers on potassium (K) content and its uptake by Indian mustard

Treatment	K content (%)		K uptake (kg ha ⁻¹)	
	Seed	Stover	Seed	Stover
Control	0.55	0.79	12.80	50.77
100% RDN through FYM	0.57	0.83	14.17	59.50
100% RDN through VC	0.57	0.84	15.27	60.50
100% RDN through PM	0.58	0.84	16.03	62.83
100% RDN through FYM+BF	0.58	0.86	15.97	63.33
100% RDN through VC+BF	0.58	0.87	16.13	64.23
100% RDN through PM+BF	0.59	0.88	17.63	65.80
Cow based bioformulation	0.57	0.83	14.00	59.50
CD (P=0.05)	0.02	0.04	1.65	6.57

Table 5. Effects of organic manures and bio-fertilizers on soil properties after harvest of Indian mustard

Treatment	Soil properties					
	pH (1:2)	EC (ds/m)	SOC (%)	Avail. N (kg/ha)	Avail. P (kg/ha)	Avail. K (kg/ha)
Control	7.80	0.80	0.65	159	13.3	269
100% RDN through FYM	7.79	0.79	0.66	160	13.4	272
100% RDN through VC	7.79	0.79	0.66	161	13.5	272
100% RDN through PM	7.78	0.79	0.67	162	13.4	274
100% RDN through FYM+BF	7.78	0.77	0.67	161	13.7	276
100% RDN through VC+BF	7.78	0.77	0.67	161	13.7	278
100% RDN through PM+BF	7.77	0.78	0.67	162	13.7	280
Cow based bioformulation	7.78	0.79	0.65	159	13.5	270
CD (P=0.05)	0.02	NS	NS	NS	0.15	5.81

3.3 Effects of Organic Manures and Bio-Fertilizers on Soil Properties after Harvest of Indian Mustard

A perusal of the data (Table 5) reveals that organic carbon, available N, P and K contents of post-harvest soil increased and/or maintained with the application of organic manures alone and along with bio-fertilizers. The maximum values of soil organic carbon (0.67 %), available N (162 kg ha⁻¹), available P (13.7 kg ha⁻¹) and

available K (280 kg ha⁻¹) were observed under treatment 100% RDN through FYM, poultry manure and vermicompost with bio-fertilizers and the minimum under control and cow-based bioformulation. The interaction effects of organic amendments i.e. FYM, vermicompost and poultry manure alone and along with bio-fertilizers on post-harvest soil properties i.e. pH, available P and K were found to be significant during first year trial whereas, it was non-significant on EC, SOC and available N. These results are in

agreement with those of Ahlawat et al. [3], Sharma et al. [14] and Kumar and Singh [12] who reported that application of farm yard manure also increased the soil organic carbon and available NPK in post-harvest soil as compared to control. This may be attributed to the beneficial role of organic manures creating favorable conditions for microbial activities in mineralization process, nutrient contents and their uptake, as well as increased availability of available nutrient pool in the soil.

4. CONCLUSIONS AND PERSPECTIVES

From the present findings, it can be concluded that the application of organic manures along with bio-fertilizers would help in the sustainable production of Indian mustard under organic farming in this locality to decrease the dependence on inorganic fertilizers. This research elucidates the fact that regular addition of organic manures alone and along with bio-fertilizers in the soil is the only way to maintain the soil health and quality. In order for farming to help mitigate the potential for devastating environmental effects, there is an apparent need for a shift towards the adoption of modern organic manure methods.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. Area coverage, production and yield of oil seeds rabi and kharif season in India. Economic Survey of India, New Delhi. 2019:21. Available:www.IndiaStat.com
2. Hadiyal JG, Kachhadiya SP, Ichchhuda PK, Kalsaria RN. Response of Indian mustard (*Brassica juncea* L.) to different levels of organic manures and bio-fertilizers. Journal of Pharmacognosy and Phytochemistry. 2017;6(4):873-875.
3. Ahlawat V, Dadarwal RS, Yadav PK, Chaudhary K. Effects of long-term nutrient management practices on physicochemical properties of soils: A review. The Pharma Innovation Journal. 2023;12(1):491-496.
4. Ramesh P, Panwar NR, Singh AB, Ramana S, Yadav SK, Rao AS. Status of organic farming in India. Current Science. 2010;98(9):1190-1194.
5. Jackson ML. Soil chemical analysis, Prentice Hall of India Pvt. Ltd., New Delhi; 1967.
6. Walkley AJ, Black CA. Estimation of soil organic carbon by the chronic acid titration method. Soil Science. 1934;37:29-38.
7. Soil Science Division Staff. Soil Survey sand. Ditzler C, Scheffe K, Monger HC (eds.). USDA Handbook 18. Government Printing Office, Washington, D.C; 2017.
8. Subbiah BV, Asija GL. A rapid procedure for the determination of available nitrogen in soils. Current Science. 1956;25:259-260.
9. Olsen SR, Cole CV, Watanabe FS, Dean LA. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. Circular-United States Department of Agriculture. 1954:939.
10. Hanway JJ, Heidel H. Soil analysis methods as used in Iowa State College, Soil Testing Laboratory. Iowa State College Bull. 1952;57:1-131.
11. Lindsay WL, Norvell WA. Development of a DTPA soil test for zinc, iron, manganese and copper. Soil Science Society of America Journal. 1978;42(3):421-448.
12. Kumar V, Singh S. Effect of fertilizers, biofertilizers and farmyard manure on sustainable production of Indian mustard (*Brassica juncea*). Annals of Plant and Soil Research. 2019;21(1):25-29.
13. Singh D, Gangwar R, Singh A. To study the effect of organic manures on nutrient uptake & economics of mustard crop. International Journal of Chemical Studies. 2020;8(4):257-259.
14. Sharma JK, Jat G, Meena RH, Purohit HS, Choudhary RS. Effect of vermicompost and nutrients application on soil properties, yield, uptake and quality of Indian mustard (*Brassica juncea*). Annals of Plant and Soil Research. 2017;19(1):17-22.

© 2023 Narender et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/100157>