



An Assessment of Anti-Diabetic Potentility of Ethanolic Extract of *Andrographis paniculata* in Alloxan Induced Rat Model

Nur-Neasha Dolon ^a, Juliana Aditi Baroi ^a,
Md. Rahmat Ullah ^a, Sayma Akter Nirzana ^b,
Redwanul Haque ^c, Fairuz Humayra ^d,
Md. Abdullah Hil Baky Rupak ^e,
Shanta Mahbub ^a and Rafat Tashin ^{a*}

^a Department of Pharmacy, University of Asia Pacific, Farmgate, Dhaka, Bangladesh.

^b Department of Pharmacy, East West University, Dhaka 1212, Bangladesh.

^c Department of Pharmacy, Northern University Bangladesh, Dhanmondi, Dhaka, Bangladesh.

^d Department of Pharmacy, Primeasia University, Bangladesh.

^e Department of Pharmaceutical Sciences, North South University, Plot # 15, Block # B, Bashundhara R/A, Dhaka, 1229, Bangladesh.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/103538>

Original Research Article

Received: 18/05/2023

Accepted: 25/07/2023

Published: 04/08/2023

ABSTRACT

Background: Diabetes is one of the most common non-communicable diseases in the world, and it is the fourth largest cause of death in the most developed countries. Plants have long been used as a source of medicine, and preclinical and clinical research has revealed that *Andrographis*

*Corresponding author: E-mail: whitefang229@gmail.com;

paniculata possesses significant anti-diabetic capabilities. The historical and current condition of *Andrographis paniculata* study in terms of medicinal usage, phytochemistry, pharmacological activity, toxicity profile, and therapeutic usage, in order to bridge the gap needing future research potential.

Methods: We divided 50 rats into ten groups and used Alloxan to induce hyperglycaemia in rats, as well as an extract of *Andrographis paniculata* soaked in ethanol. Metformin is a common anti-diabetic medication that has been demonstrated to be effective at lowering blood sugar levels without causing hypoglycaemia. Pre- and postprandial blood glucose levels, as well as cholesterol, Low Density Lipoprotein (LDL), triglycerides, and body weight, as well as liver and kidney functions, were used to measure anti-diabetic activity.

Results and Discussion: When compared to controls, the purified leaves extract substantially ($P < 0.05$) reduced blood glucose, triglyceride, Serum glutamic pyruvic transaminase (SGPT), serum glutamic-oxaloacetic transaminase (SGOT), creatinine, urea, and Low-Density Lipoprotein (LDL) levels. Serum cholesterol, on the other hand, showed no changes. Metformin had comparable impacts on these parameters. Anti-diabetic activity is demonstrated by the bioactive compounds andrographolide and 14-deoxy11-12-didehydroandrographolide.

Conclusion: Active compounds from *Andrographis paniculata* exert a hypoglycemic effect in alloxan-induced rats. It was successful in decreasing blood glucose, triglyceride, and LDL levels. They had no effect on cholesterol levels, however, and more research is required to establish their mechanisms of action. According to the findings of the study, *Andrographis paniculata* leaf extract has high therapeutic potential for the treatment of diabetes.

Keywords: *Andrographis paniculata*; lipids; blood glucose; alloxan; extract.

1. INTRODUCTION

“Diabetes mellitus (DM) is a chronic metabolic condition of glucose, lipids, and proteins that is defined by a lack of insulin synthesis and action and affects a huge number of people globally” [1]. “Because of its micro vascular (retinopathy, neuropathy, and nephropathy) and macro vascular (heart attack, stroke, and peripheral vascular disease) consequences, it has caused severe morbidity and death” [2]. In 2010, it was projected that over 200 million individuals globally suffered from diabetes, and this figure is predicted to rise to 300 million by 2025 [3].

“Currently, treatment for diabetes is primarily based on several approaches aimed at reducing hyperglycaemia, including Sulphonylureas, which increase insulin release from pancreatic islets; Metformin, which reduces hepatic glucose production; peroxisome proliferator-activated receptor-agonists (Thiazolidinediones), which enhance insulin action; α -glucosidase inhibitors, which interfere with gut glucose absorption; and insulin itself. These medicines are ineffective, have low tolerance, and have severe mechanism-based adverse effects” [4].

Traditional medicine (herbal) is utilized to treat diabetes in impoverished nations where the cost of mainstream treatments is prohibitively expensive [5]. Despite great breakthroughs in

the health sciences and medical treatment, many people use alternative therapies alone or in addition to prescription medicine [6]. One of the most significant advantages of medicinal plants is that they are widely available and have relatively few negative effects. Plants have long been an excellent source of medications, with many currently accessible pharmaceuticals produced directly or indirectly from them. According to ethno botanical data, around 800 plants may have anti-diabetic properties [7].

With a few medicinal plants with anti-diabetic activity and elucidating their mechanisms of action, such as *Brassica juncea* [8], *Eugenia jambolana* [9], *Coccinia grandis* [10], *Alangium lamarckii* [11], *Albizia odoratissima* [12], *Artemis sphaerocephala* Krasch [13], *Xonopus compressus* [14], *Berberis vulgaris* [15], *Caesalpinia digyna* [16], *Catharanthus roseus* [17], *Centaurium erythraea* [18], *Chaenomeles sinensis* [19], etc.

“*Andrographis paniculata* (AP) is a significant medicinal plant that is utilized in traditional herbal medicine in Bangladesh, China, Hong Kong, India, Pakistan, the Philippines, Malaysia, Indonesia, and Thailand” [20,21]. “It is also known as King of Bitters in English, Mahatikta in Sanskrit, Kiryato in Gujarati, Chirayetah in Hindi, Kalmegh in Bengali and Urdu, and Fah Talai Jone in Thai” [22].

It's perceived "blood purifying" property results in its use in diseases and has been reported to have a broad range of pharmacological effects, including anticancer [23], anti-diarrheal [24], antihepatitis [25], anti-HIV [26], antihyperglycemic [27], anti-inflammatory [28], antimicrobial, antimalarial [29, 30], antioxidant [31], cardiovascular [32], cytotoxic [33], hepatoprotective [34], immunostimulatory [35], and sexual dysfunctions [36].

"The aerial portion of the plant includes a great variety of chemical elements, primarily lactones, diterpenoids, diterpene glycosides, flavonoids, and flavonoid glycosides, while the leaves contain two bitter principles, andrographolide and kalmeghin" [37]. Andrographolide, a primary Ent-labdane diterpenoid of AP, contributes the most to numerous pharmacological actions. Anti-diabetic action is demonstrated by andrographolide and 14-deoxy-11,12-didehydroandrographolide [38-40].

Andrographolide reduced plasma glucose levels in streptozotocin-induced diabetic rats, the animal model used to study Type 1 diabetes [41]. However, the mechanism(s) by which andrographolide affects glucose homeostasis remain unknown. Exogenous endorphin has been shown to raise circulating insulin in adults with or without diabetes [42]. Indeed, stimulation of opioid receptors by exogenous -endorphin or pharmacological agents such as loperamide and tramadol has been shown to enhance glucose homeostasis in diabetic rats in the absence of insulin [43-44]. Furthermore, stimulation of 1A adrenoceptors in the adrenal medulla may increase -endorphin production from the rat adrenal gland [45-46].

Andrographolide activates the 1A-adrenoceptor, increasing glucose absorption in cultured mouse myoblast C2C12 cells [47]. "In order to evaluate the potential antihyperglycemic potential of *Andrographis paniculata* (AP) and the main active constituent andrographolide (AG), *in vitro* α -glucosidase and α -amylase enzyme inhibition studies, as well as *in vivo* studies to confirm the activity in live animals, were carried out" [48]. He and coworkers showed that after being converted to 14-deoxy-11-12-didehydroandrographolide, α -glucosidase inhibitor *in vivo*, AG may exhibit an anti-diabetic effect by blocking α -glucosidase. Furthermore, AP ethanol extract displayed an anti-diabetic effect via α -glucosidase inhibitory action [49]. 14-deoxy-11-12-didehydroandrographolide inhibits DN

phenotypes more effectively than andrographolide. Both of these drugs lower ROS levels but lack antioxidative action in the cellular environment, indicating that their anti-diabetic nephropathy impact is mediated through intracellular signalling pathways [50].

"It is essential to discover an integrated therapy for diabetes by employing alternative medicine such as herbal medicine or a mix of synthetic medications and herbal medicine. Numerous therapeutic qualities of *A. paniculata* have been documented. Furthermore, *A. paniculata* is included as a major medicinal plant for treating diabetes, high blood pressure, and a variety of other illnesses in the Compendium of Medicinal and Aromatic Plants" [51].

This study will go through the anti-diabetic action of *Andrographis paniculata*, which has been employed as an anti-diabetic treatment in traditional medical systems. As a result, the dry powder of *A. paniculata* was chosen for the current investigation in order to standardize its antidiabetic function in Diabetes mellitus.

2. MATERIALS AND METHODS

2.1 Drugs, Chemicals, and Instruments

The ethanol and alloxan were purchased from Sigma Aldrich in Germany. Healthcare Pharmaceutical Limited provided us with a free sample of the common diabetic medicine metformin. The blood serum analysis kits for total cholesterol, HDL, LDL, triglycerides, SGOT, SGPT, and creatinine were purchased from Plasmatic Laboratory Product Ltd. in the United Kingdom. The glucometer, Alere GI from Alere Inc., USA, was obtained from Shahbag in Dhaka, Bangladesh, and the biochemical parameters were evaluated using the Humalyzer 3000 (a semiautomated clinical chemistry analyzer).

2.2 Plant Collection and Extract Preparation

A. paniculata leaves were collected in three separate areas of Bangladesh: North Bengal, a hill track area, and a low land area. The next stage was authentication and taxonomic identification. The plant specimen was maintained at Bangladesh's National Herbarium in accordance with their laws. The leaves were dried in the shade for seven to ten days before

being finely ground. The powdered leaves were forcefully agitated for 96 hours while steeping in 70% ethanol. The extract was filtered when it had finished soaking, and the filtered liquid was collected. Once the extracted solution was brought there, it was concentrated using a rotating evaporator. The dried extract was then carefully collected and stored for future use.

2.3 Experimental Animal Handling

Adult healthy male Wistar rats (125-200 grams) were recruited from the Pharmacy Department of Jahangirnagar University in Dhaka, Bangladesh, and kept on a 12-hour dark/light cycle at a constant temperature of 25° C at the University of Dhaka's Institute of Nutrition and Food Science. On a regular basis, a normal pellet meal and clean water were provided. The rats were kept there to adapt before the investigation began. All rat trials followed the guidelines provided by the Institutional Animal Ethics Committee (IEAC). Animals were treated and maintained in accordance with the guidelines of the Swiss Academy of Medical Sciences (SAMS) and the Swiss Academy of Sciences (SCNAT).

2.4 Experimental Guidelines

All tests were carried out in accordance with the ethical principles outlined in the 2013 Helsinki Declaration. The "3R" standards, which are a cornerstone of Swiss and global legislation governing the use of animals in research, were scrupulously observed throughout the duration of this study. The term "replacement" is symbolized by the prefix "R," which encompasses both absolute replacements (such as replacing animal models with computer-generated models) and relative replacements (such as replacing live animals with cell or tissue cultures or vertebrates with invertebrates). In order to conduct perfect research, animal model was employed. Rats were selected as test animals because, unlike invertebrates, mammalian vertebrates have a particular pancreas and beta cells for antidiabetic research.

The second "R" stands for "reduction," which refers to any method that result in the use of fewer animals to acquire adequate data to answer the research objectives or in maximizing the information collected from each animal. Ten rats were gathered for this investigation based

on the "power analysis method" sample size estimate, which was utilized to guarantee compliance with this suggestion.

The third "R" stands for "refinement," and it refers to lowering the amount of pain inflicted on the experimental animals by alleviating their anguish. To make the surgery more tolerable and reduce discomfort from pinching, the tail tips of rats were rubbed with isopropyl alcohol before and after each blood glucose level measurement. The rats were fed adequately throughout the trial, and they were painlessly killed at the end in accordance with the 2013 amendment to the Guidelines for the Euthanasia of Animals.

2.5 Experimental Design

Individually weighed rats were separated into groups based on body weight and then tested for anti-hyperglycemic action (Table 1). The rodents were divided into groups based on body weight, with 10 rats in each group. Table 1 depicts the alloxan control group as rats that solely received alloxan therapy. N/A indicates that no therapeutic treatment was administered to the rats in this group.

2.6 Biological Sample Collection

Blood samples were collected by puncturing the tip of the rat's tail in order to measure blood glucose levels. In contrast, blood was immediately collected from the slaughtered animal after a heart puncture and transported to a micro centrifuge tube. The supernatant fluid was produced by centrifuging the collected samples for 5 minutes at 5,000 rpm. This fluid was then transferred to another micro centrifuge tube for biochemical testing. The kidney and liver were promptly removed from the animal body after sacrifice and meticulously washed in ice-cold saline for kidney and liver function testing.

2.7 Estimation of Biochemical Parameters

The blood glucose level was measured using a glucometer. Aside from the Humaluzer 3000, lipid profile, kidney, and liver function tests were performed. In addition, the gluconeogenic and glycolytic enzyme activity of kidney and liver samples was examined.

Table 1. Anti-hyperglycemic activity analysis

Group number	Group Status	Treatment specimen	Dose of treatment specimen (mg/kg)	Group Abbreviation
1	Negative Control	Physiological Saline	10 mL/kg	N
2	Alloxan control	Alloxan	150 mg/kg	A
3	Alloxan + Metformin	Alloxan + Metformin	150 mg/kg + 100mg	A + M100
4	Alloxan + <i>A. paniculata</i>	Alloxan + <i>A. paniculata</i> leaves extract low dose	150 mg/kg + 500 mg/kg	A + AP ₅₀₀
5	Alloxan + <i>A. paniculata</i>	Alloxan + <i>A. paniculata</i> leaves extract medium dose	150 mg/kg + 1000 mg/kg	A + AP ₁₀₀₀
6	Alloxan + <i>A. paniculata</i>	Alloxan + <i>A. paniculata</i> leaves extract high dose	150 mg/kg + 1500 mg/kg	A + AP ₁₅₀₀
7	Metformin	Metformin	100 mg/kg	M
8	<i>A. paniculata</i>	<i>A. paniculata</i> leaves extract low dose	500 mg/kg	AP ₅₀₀
9	<i>A. paniculata</i>	<i>A. paniculata</i> leaves extract medium dose	1000 mg/kg	AP ₁₀₀₀
10	<i>A. paniculata</i>	<i>A. paniculata</i> leaves extract high dose	1500 mg/kg	AP ₁₅₀₀

2.8 Statistical Analysis

For each group, the mean and standard deviation (SD) of each research parameter are displayed. The "one-way ANOVA test" was used to examine the statistical significance of intergroup heterogeneity by evaluating differences across groups in terms of different biological parameters. The application "SPSS 16" was used for the analysis. The result was considered statistically significant when the "P" value was less than 0.05 ($p < 0.05$) and highly significant when it was less than 0.01 ($p < 0.01$).

3. RESULTS AND DISCUSSION

Both the negative control and plant extract-treated groups of our experimental rats gained weight as a result of the call of nature. Alloxan had a devastating impact and significantly reduced body weight; however, because of the antioxidant properties of the plant (*A. paniculata*), the alloxan-induced treatment group did not lose the same amount of weight. However, as seen in Table 2, metformin led to an additional reduction of body weight in Group 3 since this group received both metformin and alloxan, both of which lowered weight significantly. In contrast, in group 4-6 plant extracts are administered with alloxan, and weight does not fall much. As a result, increasing dosages of plant extract have the potential to prevent animal weight loss.

100 rats were separated into ten groups to see how the extract altered blood glucose levels. Blood glucose levels were examined before giving alloxan to rats in groups 2-6. After three days, diabetes was induced. They went 14 days without treatment. The treatment then began on day 14. The drugs were likewise given to the rats in groups 7 through 10, but no alloxan was provided to them. They were treated for 14 to 42 days. Furthermore, rats in Group 1 were fed and watered on a regular basis. Alloxan is one of the most commonly utilized drugs to induce diabetes in experimental animals. As shown in Fig. 1, the blood glucose levels of the group 1 rats were normal. However, due to alloxan, the blood glucose level in the positive control group was greater than in all other groups. When compared to a positive control, metformin appears to lower blood glucose levels. On the contrary, all dosages of plant extract considerably lower blood glucose levels, while medium and high doses outperform low ones. As a result, its anti-diabetic effects will be naturally greater than metformin's.

As shown in Table 3, the treatment group had lower creatinine and urea levels during the renal function test than the alloxan control group. In groups 4-6, all dosages of plant extract significantly lowered creatinine levels when compared to the Alloxan Control group. Group 6 (Alloxan + *A. paniculata* leaves extract, high dosage) beat the Alloxan Control group in terms

of urea levels. There was no significant difference in creatinine deduction between metformin and placebo.

In the liver functional test, alloxan-induced diabetic rats had substantially higher SGOT and SGPT values. The treatment groups' SGOT and

SGPT levels were lower than all other groups, including the Alloxan control group. The hepatoprotective effect of *A. paniculata* leaves extract was seen in the reduction of liver enzymes. The high dose lowered the excessively high level of SGOT, whereas the medium and high doses reduced SGPT levels.




Fig. 1. Blood glucose levels of rats in ten groups over the course of treatment

Values were presented as the mean \pm SD ($n = 10/\text{group}$). * $P < 0.05$ and ** $P < 0.01$ indicate significant difference from the disease group (C = control group, A = alloxan-treated group, M = metformin, A + M = Alloxan + metformin, A + AP = Alloxan + *Andrographis paniculata*, and AP = *Andrographis paniculata*)

Table 2. Body weight of rats before the initiation and after the termination of the experiment

Group number	Group status	Body Weight (gm)	
		Initial	Final
1	Negative Control	121.47 \pm 7.34	148.32 \pm 8.93
2	Alloxan control	124.37 \pm 6.93	103.67 \pm 5.86
3	Alloxan + Metformin	128.45 \pm 4.37	100.35 \pm 5.48
4	Alloxan + <i>A. paniculata</i>	120.84 \pm 5.57	108.87 \pm 7.35
5	Alloxan + <i>A. paniculata</i>	126.38 \pm 5.24	115.19 \pm 6.91
6	Alloxan + <i>A. paniculata</i>	129.35 \pm 7.61	124.25 \pm 4.57
7	Metformin	119.45 \pm 5.47	130.30 \pm 6.14
8	<i>A. paniculata</i>	123.59 \pm 7.32	138.39 \pm 5.54
9	<i>A. paniculata</i>	127.50 \pm 5.50	140.47 \pm 7.30
10	<i>A. paniculata</i>	122.20 \pm 3.99	134.87 \pm 8.38

Table 3. Effect of *A. paniculata* on the kidney, liver, and lipid function of control and experimental rats

Group no.	Group status	Kidney Function test		Liver Function test		Lipid Profile Function Test			
		Creatinine (mg/dl)	Urea	SGOT (u/l)	SGPT (u/l)	Cholesterol (mg/dl)	Triglyceride (mg/dl)	LDL (mg/dl)	HDL (mg/dl)
1	Negative Control	0.6±0.02	24.3 ±1.09	33.52 ±0.39	42.63 ±3.49	97.75 ± 3.14	54.14 ±2.41	37.33 ±1.72	74.42 ±1.73
2	Alloxan control	2.3±0.9253	94.9 ±6.87	81.83 ±4.27	89.75 ±6.41	131.32 ± 7.71	113.46 ±3.63	72.45 ±4.62	53.16 ±2.47
3	Alloxan + Metformin	1.3±0.235	79.3 ±4.31	64.34 ±5.59	71.30 ±4.55	110.43 ± 5.46	84.14 ±7.30	56.40 ±3.32	64.50 ±3.30
4	Alloxan + <i>A. paniculata</i>	1.8±0.23	92.83 ±6.31	78.80 ±4.37	88.34 ±4.60	130.44 ± 6.83	108.49 ±6.30	70.75 ± 3.31	55.93 ±4.14
5	Alloxan + <i>A. paniculata</i>	1.4±0.12	90.33 ±4.82	74.93 ±5.17	86.12 ±5.59	128.46 ± 8.34	104.34 ±4.69	63.22 ±4.50	57.17 ±3.10
6	Alloxan + <i>A. paniculata</i>	0.9±0.09	86.83 ±5.87	71.30 ±4.14	82.82 ±5.46	125.30 ± 7.30	98.41 ±3.89	59.83 ±3.41	60.25 ±4.66
7	Metformin	0.5±0.03	26.30 ±1.32	37.30 ±1.09	39.40 ±2.48	95.50 ± 3.30	50.57 ±2.33	35.52 ±2.73	74.14 ±4.50
8	<i>A. paniculata</i>	0.7±0.359	25.93 ±0.97	30.42 ±0.87	44.23 ±3.38	99.93 ± 4.10	58.32 ±3.93	33.18 ±3.16	71.30 ±3.24
9	<i>A. paniculata</i>	0.7±0.069	28.81 ±1.04	34.14 ±2.01	42.46 ±3.10	92.14 ± 3.38	49.82 ±3.84	38.30 ±4.40	73.91 ±2.48
10	<i>A. paniculata</i>	0.6±0.03	28.74 ±1.81	31.73 ±1.78	41.93 ±2.87	94.38 ± 4.16	56.94 ±4.10	36.36 ±4.01	75.75 ±3.32

All dosages of *A. paniculata* leaves extract demonstrate no reduction in cholesterol equivalency compared to the Alloxan-controlled groups, as shown in Table 3, because alloxan also increased the levels of these indicators. According to Table 3, the negative control group had the highest HDL level, whereas the Alloxan control group had the lowest. Thus, the quantity of HDL was increased in both the metformin and high dose *A. paniculata* plant extract instances. The drug-treated groups 5 and 6 (Alloxan + *A. paniculata* leaves extract medium dosage and Alloxan + *A. paniculata* leaves extract, high dose) outperformed the positive control group in terms of LDL levels. In the case of triglycerides, a higher dose of plant extract has a greater effect and significantly lowers triglycerides.

Human fasting blood glucose levels were dramatically reduced after oral treatment with ethanol extracts of AP. Another bioactive molecule, 14-deoxy-11-12-didehydroandrographolide, was shown to have antihyperglycemic properties. We discovered that groups 5 and 6 performed better in terms of blood glucose-lowering impact, SGOT, SGPT, creatinine, total cholesterol, and LDL, triglyceride, and HDL levels in our study. As a result, researchers will work on identifying new antihyperglycemic components of AP and combining AP with other medicinal plants for a better therapy option for diabetic patients.

4. CONCLUSION

The anti-diabetic effect of *Andrographis paniculata* was observed in the ethanolic extract and the fractional compound (i.e., Andrographolide or its analogue, 14-deoxy-11-12-didehydroandrographolide) in alloxan-induced diabetic rats. As a result, the folk usage of this herb to treat diabetes is warranted. Because of its wide pharmacological actions, the AP may be fairly classified as a modern catholicon. However, the identified pharmacological actions of AP require clinical confirmation. Verification of the efficacy of additional biological activities of AP, such as anti-cancer, anti-inflammatory, and hepatoprotective actions, on human research participants would provide significant advantages to the world's biggest population. We believe that in the near future, the AP might be used as an extremely effective therapeutic agent for a range of ailments in order to treat human diseases. As a result, *Andrographis paniculata* may be developed as an alternate anti-diabetic medication. Further research may

be conducted to investigate the mechanism of action of *Andrographis paniculata* as an anti-diabetic in its extract, fractional component, and combination with a synthetic medicine or another herbal.

CONSENT

It is not applicable.

ETHICAL APPROVAL

Animal Ethic committee approval has been taken to carry out this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Pareek H, Sharma S, Khajja BS, Jain K, Jain GC. Evaluation of hypoglycemic and anti-hyperglycemic potential of *Tridax procumbens* (Linn.). BMC Complementary and Alternative Medicine. 2009;9:1-7.
2. Thévenod F. Pathophysiology of diabetes mellitus type 2: roles of obesity, insulin resistance and β -cell dysfunction. In Diabetes and Cancer. Karger Publishers. 2008;19:1-18.
3. Amos AF, McCarty DJ, Zimmet P. The rising global burden of diabetes and its complications: estimates and projections to the year 2010. Diabetic medicine. 1997;14(S5):S7-S85.
4. Hosseyni ES, Kashani HH, Asadi MH. Mode of action of medicinal plants on diabetic disorders. Life Sci J. 2012; 9(4):2776-83.
5. Saravanan G, Pari L. Hypoglycaemic and antihyperglycaemic effect of *Syzygium cumini* bark in streptozotocin-induced diabetic rats. J Pharmacol Toxicol. 2008; 3(1):1-10.
6. Alexiou P, Demopoulos VJ. A diverse series of substituted benzenesulfonamides as aldose reductase inhibitors with antioxidant activity: design, synthesis, and *in vitro* activity. Journal of medicinal chemistry. 2010;53(21):7756-7766.
7. Alarcon-Aguilara FJ, Roman-Ramos R, Perez-Gutierrez S, Aguilar-Contreras A, Contreras-Weber CC, Flores-Saenz, JL. Study of the anti-hyperglycemic effect of

- plants used as antidiabetics. *Journal of Ethnopharmacology*. 1998;61(2):101-110.
8. Thirumalai T, Therasa SV, Elumalai EK, David E. Hypoglycemic effect of *Brassica juncea* (seeds) on streptozotocin induced diabetic male albino rat. *Asian Pacific Journal of Tropical Biomedicine*. 2011;1(4):323-325.
 9. Ravi K, Ramachandran B, Subramanian, S. Effect of *Eugenia jambolana* seed kernel on antioxidant defense system in streptozotocin-induced diabetes in rats. *Life sciences*. 2004;75(22):2717-2731.
 10. Ajay SS. Hypoglycemic activity of *Coccinia Indica* (Cucurbitaceae) leaves. *International Journal of PharmTech Research*. 2009;1(3):892-893.
 11. Kumar R, Pate DK, Prasad SK, Sairam, K, Hemalatha S. Antidiabetic activity of alcoholic leaves extract of *Alangium lamarckii* Thwaites on streptozotocin–nicotinamide induced type 2 diabetic rats. *Asian Pacific Journal of Topical Medicine*. 2011;4(11):904-909.
 12. Kumar D, Kumar S, Kohli S, Arya R, Gupta J. Antidiabetic activity of methanolic bark extract of *Albizia odoratissima Benth.* in alloxan induced diabetic albino mice. *Asian Pacific Journal of Tropical Medicine*. 2011;4(11):900-903.
 13. Hu XZ, Xing XH, Zhang ZM, Wu RQ, Guo, Q, Cui SW, Wang Q. Antioxidant effects of *Artemis sphaerocephala Krasch.* Gum, on streptozotocin-induced type 2 diabetic rats. *Food Hydrocolloids*. 2011;25(2):207-213.
 14. Ibeh BO, Ezeaja MI. Preliminary studies of antidiabetic activity of the methanolic leaf extract of *Axonopus compressus* (P. Beauv) in alloxan-induced diabetic rats. *Journal of Ethnopharmacology*. 2011;138(3):713-716.
 15. Meliani N, Dib MEA, Allali H, Tabti B. Hypoglycaemic effect of *Berberis vulgaris L.* in normal and streptozotocin-induced diabetic rats. *Asian Pacific Journal of Tropical Biomedicine*. 2011;1(6):468-471.
 16. Kumar R, Patel DK, Prasad SK, Laloo D, Krishnamurthy S, Hemalatha S. Type 2 antidiabetic activity of bergenin from the roots of *Caesalpinia digyna* Rottler. *Fitoterapia*. 2012;83(2):395-401.
 17. Ohadoma SC, Michael HU. Effects of co—administration of methanol leaf extract of *Catharanthus roseus* on the hypoglycemic activity of metformin and glibenclamide in rats. *Asian Pacific Journal of Tropical Medicine*. 2011;4(6):475-477.
 18. Sefi M, Fetoui H, Lachkar N, Tahraoui A, Lyoussi B, Boudawara T, Zeghal N. *Centaurium erythrea* (Gentianaceae) leaf extract alleviates streptozotocin-induced oxidative stress and β -cell damage in rat pancreas. *Journal of Ethnopharmacology*. 2011;135(2):243-250.
 19. Sancheti S, Sancheti S, Seo SY. Antidiabetic and antiacetylcholinesterase effects of ethyl acetate fraction of *Chaenomeles sinensis* (Thouin) Koehne fruits in streptozotocin-induced diabetic rats. *Experimental and Toxicologic Pathology*. 2013;65(1-2):55-60.
 20. Akbar S. *Andrographis paniculata*: a review of pharmacological activities and clinical effects. *Alternative Medicine Review*. 2011;16(1):66-77.
 21. Kabir MH, Hasan N, Rahman MM, Rahman MA, Khan JA, Hoque NT, Rahmatullah M. A survey of medicinal plants used by the Deb barma clan of the Tripura tribe of Moulvibazar district, Bangladesh. *Journal of Ethnobiology and Ethnomedicine*. 2014;10:1-28.
 22. Li J, Huang W, Zhang H, Wang X, Zhou, H. Synthesis of andrographolide derivatives and their TNF- α and IL-6 expression inhibitory activities. *Bioorganic & Medicinal Chemistry Letters*. 2007;17(24):6891-6894.
 23. Kumar RA, Sridevi K, Kumar NV, Nanduri, S, Rajagopal S. Anticancer and immunostimulatory compounds from *Andrographis paniculata*. *Journal of Ethnopharmacology*. 2004;92(2-3):291-295.
 24. Gupta S, Choudhry MA, Yadava JNS, Srivastava V, Tandon JS. Antidiarrhoeal activity of diterpenes of *Andrographis paniculata* (Kal-Megh) against *Escherichia coli* enterotoxin in *in vivo* models. *International Journal of Crude Drug Research*. 1990;28(4):273-283.
 25. Sharma A, Singh RT, Sehgal V, Handa S. S. Antihepatotoxic activity of some plants used in herbal formulations. *Fitoterapia*. 1991;62(2):131-138.
 26. Nanduri S, Nyavanandi VK, Thunuguntla, SSR, Kasu S, Pallerla MK, Ram PS, Akella V. Synthesis and structure–activity relationships of andrographolide analogues as novel cytotoxic agents. *Bioorganic & Medicinal Chemistry Letters*. 2004;14(18):4711-4717.
 27. Subramanian R, Asmawi MZ, Sadikun A. *In vitro* α -glucosidase and α -amylase

- enzyme inhibitory effects of *Andrographis paniculata* extract and andrographolide. *Acta Biochimica Polonica*. 2008;55(2):391-398.
28. Chiou WF, Lin JJ, Chen CF. Andrographolide suppresses the expression of inducible nitric oxide synthase in macrophage and restores the vasoconstriction in rat aorta treated with lipopolysaccharide. *British Journal of Pharmacology*. 1998;125(2):327-334.
 29. Wiart C, Kumar K, Yusof MY, Hamimah, H, Fauzi ZM, Sulaiman M. Antiviral properties of ent-labdene diterpenes of *Andrographis paniculata* Nees, inhibitors of herpes simplex virus type 1. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*. 2005;19(12):1069-1070.
 30. Misra P, Pal NL, Guru PY, Katiyar JC, Srivastava V, Tandon JS. Antimalarial activity of *Andrographis paniculata* (Kalmegh) against Plasmodium berghei NK 65 in Mastomys natalensis. *International Journal of Pharmacognosy*. 1992;30(4):263-274.
 31. Kamdem RE, Sang S, Ho CT. Mechanism of the superoxide scavenging activity of neoandrographolide– a natural product from *Andrographis paniculata* nees. *Journal of Agricultural and Food Chemistry*. 2002;50(16):4662-4665.
 32. Zhang CY, Tan BKH. Mechanisms of cardiovascular activity of *Andrographis paniculata* in the anaesthetized rat. *Journal of Ethnopharmacology*. 1997; 56(2):97-101.
 33. Nanduri S, Nyavanandi VK, Thunuguntla, SSR, Kasu S, Pallerla MK, Ram PS, Akella V. Synthesis and structure–activity relationships of andrographolide analogues as novel cytotoxic agents. *Bioorganic & Medicinal Chemistry letters*. 2004;14(18):4711-4717.
 34. Handa SS, Sharma A. Hepatoprotective activity of andrographolide from *Andrographis paniculata* against carbontetrachloride. *The Indian Journal of Medical Research*. 1990;92:276-283.
 35. Calabrese C, Berman SH, Babish JG, Ma, X, Shinto L, Dorr M, Standish LJ. A phase I trial of andrographolide in HIV positive patients and normal volunteers. *Phytotherapy Research*. 2000; 14(5):333-338.
 36. Akbarsha MA, Murugaian P. Aspects of the male reproductive toxicity/male antifertility property of andrographolide in albino rats: effect on the testis and the cauda epididymidal spermatozoa. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*. 2000;14(6):432-435.
 37. Akbar S. *Andrographis paniculata*: a review of pharmacological activities and clinical effects. *Alternative Medicine Review*. 2011;16(1):66-77.
 38. Yu BC, Chang CK, Su CF, Cheng JT. Mediation of β -endorphin in andrographolide-induced plasma glucose-lowering action in type I diabetes-like animals. *Naunyn-Schmiedeberg's Archives of Pharmacology*. 2008;377:529-540.
 39. Subramanian R, Asmawi MZ, Sadikun A. Effect of andrographolide and ethanol extract of *Andrographis paniculata* on liver glycolytic, gluconeogenic, and lipogenic enzymes in a type 2 diabetic rat model. *Pharmaceutical Biology*. 2008;46 (10-11):772-780.
 40. Subramanian R, Asmawi MZ, Sadikun A. Effect of ethanolic extract of *Andrographis paniculata* (Burm. F.) nees on a combination of fat-fed diet and low dose streptozotocin induced chronic insulin resistance in rats. *Diabetologia Croatica*. 2008;37(1):13-22.
 41. Yu BC, Chen WC, Cheng JT. Antihyperglycemic effect of andrographolide in streptozotocin-induced diabetic rats. *Planta Medica*. 2003;69(12): 1075-1079.
 42. Curry DL, Bennett LL, Li CH. Stimulation of insulin secretion by beta-endorphins (I-27 & I-31). *Life Sciences*. 1987; 40(21):2053-2058.
 43. Liu IM, Chi TC, Chen YC, Lu FH, Cheng, JT. Activation of opioid μ -receptor by loperamide to lower plasma glucose in streptozotocin-induced diabetic rats. *Neuroscience Letters*. 1999;265(3):183-186.
 44. Cheng JT, Liu IM, Chi TC, Tzeng TF, Lu, FH, Chang CJ. Plasma glucose–lowering effect of tramadol in streptozotocin-induced diabetic rats. *Diabetes*. 2001; 50(12):2815-2821.
 45. Cheng JT, Liu IM, Kuo DH, Lin MT. Stimulatory effect of phenylephrine on the

- secretion of β -endorphin from rat adrenal medulla in vitro. *Autonomic Neuroscience*. 2001;93(1-2):31-35.
46. Liu IM, Chen WC, Cheng JT. Mediation of β -endorphin by isoferulic acid to lower plasma glucose in streptozotocin-induced diabetic rats. *Journal of Pharmacology and Experimental Therapeutics*. 2003; 307(3):1196-1204.
47. Hsu JH, Liou SS, Yu BC, Cheng JT, Wu, YC. Activation of α 1A-adrenoceptor by andrographolide to increase glucose uptake in cultured myoblast C2C12 cells. *Planta Medica*. 2004; 70(12):1230-1233.
48. Akowuah GA, Zhari I, Norhayati I, Mariam, A. HPLC and HPTLC densitometric determination of andrographolides and antioxidant potential of *Andrographis paniculata*. *Journal of Food Composition and Analysis*. 2006;19(2-3):118-126.
49. Kapil A, Koul IB, Banerjee SK, Gupta BD. Antihepatotoxic effects of major diterpenoid constituents of *Andrographis paniculata*. *Biochemical Pharmacology*. 1993;46(1):182-185.
50. Lee MJ, Rao YK, Chen K, Lee YC, Chung, YS, Tzeng YM. Andrographolide and 14-deoxy-11, 12-didehydroandrographolide from *Andrographis paniculata* attenuate high glucose-induced fibrosis and apoptosis in murine renal mesangial cell lines. *Journal of Ethnopharmacology*. 2010;132(2):497-505.
51. Sukhdev SH, Dev D, Rakesh KV. Compendium of medicinal and aromatic plants. New Delhi: ASIA. 2006;2:50-148.

© 2023 Dolon et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/103538>