



Comparative Evaluation and Co-relation of Difficult Airway and Temporomandibular Joint Disorder in Skeletal Class II Vertical Growth Pattern and Class-I Malocclusion – An Observational Study

Shruti Rathi ^{a≡ θ *} and Rizwan Gilani ^{b≡#}

^a *M.D.S Orthodontics and Dentofacial Orthopaedics, Sharad Pawar Dental College, Datta Meghe Institute of Medical Sciences (Deemed to be University), Wardha, India.*

^b *Department of Orthodontics and Dentofacial Orthopaedics, Sharad Pawar Dental College, Datta Meghe Institute of Medical Sciences (Deemed to be University), Wardha, India.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i60B34996

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/80041>

Study Protocol

Received 20 October 2021
Accepted 26 December 2021
Published 27 December 2021

ABSTRACT

Introduction: In the recent years it is observed that there has been increase in the no. of cases of Temporomandibular joint disorder and TMJ pain related problem, many of patients go undiagnosed and may transform into irreversible severe cases if the predisposing factors are present. There are different clinical and radiographical methods of diagnosing TMD cases, and one of the simplest ways to diagnose and categorize a Temporomandibular disorder is Helkimo index. The present study is encouraged to measure difficult airway, in Class II malocclusion with TMD.

Materials and Method: The observational study will be conducted in Orthodontics and Dentofacial Orthopaedics department, Sharad Pawar Dental College, Sawangi (M), Wardha in collaboration with department of Radiology, Acharya Vinoba Bhave Hospital. Sawangi (M), Wardha. Total 30 patient in age group of 20 to 50 years, will be selected from the patients coming to Out Patient Department (OPD) of Orthodontics and Dentofacial Orthopaedics, Wardha. MRI of patients with

[≡] Dr.;

[⊖] Post- Graduate;

[#] Reader;

*Corresponding author: E-mail: rathi09shruti@gmail.com;

skeletal class II vertical growth pattern and class I malocclusion having temporomandibular joint disorder will be taken and then airway is to be evaluated.

Expected Results: It is expected that individuals with skeletal class II vertical growth pattern and class I malocclusion with temporomandibular joint disorder may have a compromised airway.

Conclusion: No such study has been carried out to evaluate co-relation of difficult airway and temporomandibular joint disorder. Thus, this study aims to evaluate it in skeletal class II vertical growth pattern and class I malocclusion.

Keywords: Temporomandibular joint disorder; skeletal class II; Vertical growth pattern; Class-I malocclusion; compromised airway.

1. INTRODUCTION

"Temporomandibular joint disorder (TMD) is defined as pain and compromised movement of jaw joint and the surrounding muscles usually in front of ears and sometimes in the form of headache [1]."

In the recent years increase in the number of TMD cases and adults with pain in TMJ and related problem many of patients go undiagnosed and may convert into irreversible severe cases if the predisposing factors are present [2].

There are various clinical and radiographical methods for diagnosing TMD cases, and one of the easiest ways to diagnose and categorize a Temporomandibular disorder is Helkimo index [3]. It includes both anamnestic component and clinical examination of dysfunction which categorizes the TMD as mild moderate or severe [4]. There are various remedial way of treating TMD's, (such as orthopedic stabilization, physiotherapy, pharmacological mode and jaw exercise) [5,6].

Growth deformity in skeletal class II malocclusion caused by the bones usually related with retrusion of mandible in relation to facial structure [7]. The malformation causes functional disorders which involve the upper airway and the TMJ [8]. Various studied had been done to measure the airway; but, prevalence of Temporomandibular disorders in Class II malocclusion with difficult airway measurement are rare, which encouraged the present study [9,10]

1.1 Aim

Comparative evaluation and co-relation of difficult airway and Temporomandibular joint disorder in skeletal class II vertical growth pattern and class I malocclusion - An observational study.

1.2 Objectives

To evaluate grading of difficult airway in class I and skeletal class II vertical malocclusion. Not to be included. There are only 4 objectives.

1. To evaluate severity of Temporomandibular joint disorder in class I and skeletal class II vertical malocclusion.
2. To compare grading of difficult airway in skeletal class II vertical malocclusion with class I malocclusion.
3. To compare severity of Temporomandibular joint disorder in skeletal class II vertical with class I malocclusion
4. To co-relate grading of difficult airway and severity of Temporomandibular joint disorder in skeletal class II vertical with class I malocclusion.

2. MATERIALS AND METHODS

2.1 Study Design

This is an observational retrospective study to be conducted in the department of Orthodontics and Dentofacial Orthopaedics, Sharad Pawar Dental College, Sawangi (M), Wardha. The patient in the age group 20-50 will be taken. The age group is taken as adults since growth is completed and there will be no further changes in TMJ and mandible.

2.2 Sample Formula

Based on the prevalence of patients reporting to SPDC OPD with difficult airway and TMD, the sample size was calculated using single proportion. The sample size was calculated to be 15 in each group.

$$\text{Total sample size} = 2 \times 15 = 30$$

This calculator uses the following formula for the sample size n:

$$n=N*X(X+N-1),$$

where,

$$X=Za/22*p*(1-p)/MOE2,$$

and $Za/2$ is the critical value of the normal distribution at $a/2$ (e.g. for a coincidence level of 95%, a is 0.05 and the critical value is 1.96), MOE is the margin of error, $p= 0.95\%$ is the sample proportion, N is the population size.

Careful history, clinical examination and radiographic examination will be done. For each patient, the Helkimo index will be obtained which will classify them as mild, moderate and severe cases from which moderate cases will be selected for study. Based on Helkimo index patient will be classified for TMD and then for difficult airway.

2.3 Inclusion Criteria

1. Individuals who have a malocclusion of class I type.
2. Individuals who have a malocclusion of class II type with vertical growth pattern having Temporomandibular disorder.

2.4 Exclusion Criteria

1. Individuals with class II horizontal growth pattern.
2. Individuals with class III malocclusion
3. Individuals with non -TMD class II div 1 malocclusion.
4. Individuals with any systemic disease or any muscular dystrophies.
5. Syndromic Cases.
6. Severe skeletal asymmetry.

2.5 Method

The following study will be carried out in the Department of Orthodontics and Dentofacial Orthopedics, Sharad Pawar Dental College.

The study will include cases in total. The patients with class II div 1 will be selected from OPD and consent will be taken for the participation in the study. The total cases will be divided into 2 groups:

Group 1: Class I malocclusion

Group 2: Class II malocclusion with vertical growth pattern.

From the randomly selected cases in the OPD the cephalometric analysis of subjects is done.

Digital records of the patient (lateral cephalogram, photograph, models) will be taken and stored. The Helkimo index will be taken to assess the existence and severity of TMD in class II div 1 malocclusion patient which will classify them as mild, moderate and severe cases from which moderate to severe cases will be selected for study.

After the clinical and cephalometric analysis of patient with Temporomandibular disorder they will be subjected to MRI for airway evaluation.

Upper pharynx – upper pharyngeal width is measured from posterior most part of soft palate to the closest point of pharyngeal wall. Measurement of width is taken from anterior half of the soft palate outline. As it is the region for upper respiratory patency. Since a head film of nasopharynx is a 2D representation of a 3D structure.

Lower pharynx -lower pharyngeal width is considered from the junction of the posterior border of the tongue and inferior border of the mandible to the closest point on the posterior pharyngeal wall.

Chart 1. Classification of malocclusion

Class I malocclusion	Class II malocclusion with vertical growth pattern
ANB angle - 2°	ANB angle - 4°
FMA angle- $22-28^{\circ}$	FMA angle- $> 30^{\circ}$
Y-axis - 59.4°	Y-axis - $>59.4^{\circ}$
SN-MP - 21.9°	SN-MP - $>21.9^{\circ}$
Gonial angle - $128+7^{\circ}$	Gonial angle - $> 135^{\circ}$
Beta angle - $27- 33^{\circ}$	Beta angle - $<25^{\circ}$

Soft copy of MRI of individuals will be procured and converted to slices or cuts to JPG format. These slices will then be analyzed on AutoCAD 2010 software. Then the image will be scaled in millimeter scale. Using the scale option, the distance between the two points will be measured.

2.6 Statistical Analysis

The analysis be done with use of descriptive and inferential statistics (chi square test, student's paired t and unpaired t test, ANOVA test). Analysis will use software of SPSS 22.0 version and graph pad prism of 6.0 version and $P < 0.05$ will be taken as level of importance.

3. EXPECTED RESULT

It is expected that individuals with skeletal class II vertical growth pattern and class I malocclusion with temporomandibular joint disorder may have a compromised airway.

4. DISCUSSION

Patients seeking orthodontic treatment are mostly unaware of the underlying TMD's since they are asymptomatic. During the treatment or after the completion they encounter difficulties with TMJ and airway. Class II malocclusion with vertical growth pattern are the majority population for difficult airway.

This study will assist in early diagnosis of probability of patients having TMD with difficult airway. Hence it will provide better treatment planning for both improvisation of airway as well as preventing TMD.

Kirsi et al. in 2011 conducted a study to test the difference between the pharyngeal obstruction using MRI and lateral cephalogram. He took images of upper airway using lateral cephalogram and MRI and clinical observation of tonsil was also done. He found that there was significant positive relation in the values of MRI and lateral cephalogram. Therefore he concluded that MRI is a better diagnostic criteria for measurement of upper airway.

Nayana et al. in 2015 assessed the upper airway measurement in class II skeletal malocclusion to study the association between the measurements, position, length of the mandible. He studied 80 lateral cephalograms with class I

and class II malocclusion and measurements were used for cephalometric study. He concluded that in class II individuals the measurement of nasopharynx and oropharynx was compromise. Reduced upper airway measurements were found in class II malocclusion.

Few of the related studies on temporomandibular joint were reviewed [11,12]. Also studies on Class- I, II, and III malocclusion diagnosis and corrections were reviewed [13-15].

5. CONCLUSION

No such study has been carried out to evaluate co-relation of difficult airway and temporomandibular joint disorder. Thus, this study aims to evaluate it in vertical growth pattern for skeletal class II and class I malocclusion. This study will assist in early diagnosis of co-relation of difficult airway and Temporomandibular joint Disorder. It will indicate the potential risks factors that lead to Temporomandibular joint Disorders in asymptomatic cases.

CONSENT

Patient will be informed about the study and their consent will be taken.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval will be collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Dworkin SF, Huggins KH, LeResche L, Von Korff M, Howard J, Truelove E, et al. Epidemiology of signs and symptoms in temporomandibular disorders: Clinical signs in cases and controls. *J Am Dent Assoc* 1990;120(3):273-81.
2. Nordstrom D. ALF Appliance. Available:<http://bit.ly/1fXdioQ> Published February 12, 2008 Accessed August 10, 2013
3. Delz E. The ALF (Advanced Lightwire Functional appliance) creating facial beauty and balance. *IJO*. 2009;20(2).

4. Strokon D. Correction of dental and cranial sidebend with ALF. IJOM. 2010;21:3.
5. Joseph AA, Elbowaa J, Cisneros GJ, Eisig S B. A Cephalometric comparative study of the soft tissue airway dimensions in persons with hyperdivergent and normodivergent facial pattern. J Oral Maxillofac surg. 1998;56(2):135-9.
6. Sundrani A, Kamble RH, Ahuja MM, Bhola N, Shrivastav S. Surgical orthodontic correction of class II malocclusion with vertical maxillary excess and gummy smile- A case report. Journal of Evolution of Medical and Dental Sciences. 2020;9(1):67-71.
7. Dunn GF, Green LJ, Cunat JJ. Relationships between variation of mandibular morphology and variation of nasopharyngeal airway size in monozygotic twins. Angle Orthod. 1973;43:129-135.
8. John ZA, Shrivastav SS. A tri-laminated splint for management of anterior disk displacement. Medical Science. 2020;24(104):2096-102.
9. Pirilä-Parkkinen K, Löppönen H, Nieminen P, Tolonen U, Pääkkö E, Pirttiniemi P. Validity of upper airway assessment in children: A clinical, cephalometric, and MRI study. Angle Orthod. 2011;81(3):433-9.
10. Lateral Cephalogram and CBCT as a diagnostic aid for analysis of airway- Review Article. European Journal of Molecular & Clinical Medicine. 2020;7(7):1802-1808.
11. Dhannawat P, Shrivastav S, Ranjit K, Banerjee S. Different types of occlusal splint used in management of temporomandibular joint disorders- A review. European Journal of Molecular and Clinical Medicine. 2020;7(7):1809–15.
12. Naqvi WM, Fating TB. Temporomandibular joint dysfunction: A non-invasive approach towards the management of a long-standing condition. International Journal of Pharmaceutical Research. 2020;12:1131–34. Available:https://doi.org/10.31838/ijpr/2020.SP1.131.
13. Awasthi Eshan, Nitin Bhola, Ranjeet Kamble, Sunita Shrivastav, Abhilasha Goyal. Treatment of an adult with skeletal class III and a hemimandibular elongation a multidisciplinary approach. Journal of Clinical and Diagnostic Research. 2016;10(9):ZJ5–6. Available:https://doi.org/10.7860/JCDR/2016/20098.8468

© 2021 Rathi and Gilani; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/80041>*