



Impact of Foliar Feeding of Borax, Zinc Sulphate and Their Intraction on Post Harvest Losses and Economy of Guava (*Psidium guajava* L.) Cv. Gwalior 27

**Komal Yadav ^{a*}, Shashi S. Yadav ^b, Manju verma ^c,
Priyadarshani Arun Khambalkar ^b, Sunil Kumar Yadav ^d,
Shankar Lal Yadav ^b and Suman Chopra ^e**

^a Department of Horticulture, College of Agriculture, Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior, 474002, India.

^b Department of Soil Science, College of Agriculture, Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior, 474002, India.

^c Department of Horticulture, College of Agriculture, Nagaur, 341001, India.

^d Department of Horticulture, College of Agriculture, Shri Karan Narendra Agriculture University, Jobner, 303329, India.

^e Department of Plant Pathology, College of Agriculture, Shri Karan Narendra Agriculture University, Jobner, 303329, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i113153

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105394>

Original Research Article

Received: 16/06/2023

Accepted: 23/08/2023

Published: 05/10/2023

ABSTRACT

A field experiment entitled "Effect of foliar application of borax and zinc sulphate on growth, yield and quality of guava (*Psidium guajava* L.) cv. Gwalior-27" was carried out at Agrotechnology Park, Krishi Vigyan Kendra, College of Agriculture, Gwalior (M.P.) during 2016-2017. The experiment was

*Corresponding author: E-mail: dr.yadavkomal19@gmail.com;

consisting of 9 treatments having two levels of each Borax (0.2 per cent and 0.4 per cent) and Zinc sulphate (0.4 and 0.6 per cent). The experiment was laid out in Factorial randomized block design with three replications. Among different doses of foliar spray of nutrients, Zinc sulphate @ 0.6 (T₅) were found significantly superior over control with respect physiological loss in weight (g) and shelf life of guava. While the minimum spoilage % was recorded with treatment B₂Z₂ (Borax@ 0.4% + ZnSo₄@ 0.6%) at 5th, 10th and 15th day of storage respectively.

Keywords: Zinc sulphate; borax; guava; post harvest losses.

1. INTRODUCTION

The most widespread but significant fruit crop in India is the guava. After mango, banana, and citrus, it ranked fourth in terms of acreage and output. It belongs to the *Psidium* genus, which has more than 150 species. Due to its hardiness and tolerance to a wider range of environments, *Psidium guajava* L. was more economically utilized. Even with little care, it is effectively cultivated in a variety of soil types and climatic environments. Attention must be paid to organised fruit production if we are to harvest field produce of higher quality. It is important to emphasise certain procedures and approaches for improving the fruit's post-harvest quality. It is believed that Mrigbahar is the best time to purchase high-quality guava fruits because fruit flies are no longer an issue. The guava tree blooms three times a year, in February, June-July, and October, in eastern and southern India. The corresponding bahars are known by the names "Ambe," "Mrig," and "Hasta" bahar. Among these three bahars, "Mrigbahar" fruits, which are better in quality, flavour, and vitamin C content, ripen in the winter, from November to January.

Plants can get micronutrient applications by foliar and soil applications. Micronutrient application to the foliage is more efficient than application to the soil. According to Keram et al. [1], one of the eight required trace elements or micronutrients for agricultural plants' typical, healthy development and reproduction is zinc. However, there is little research on the use of micronutrients to increase fruit size and overall fruit quality in India and other countries. In recent years, it has become increasingly clear how important micronutrients are to increasing the output and improving the quality of fruit harvests [2]. In light of this, a study was conducted to evaluate the potential effects of micronutrients on the guava fruit's development, yield, and sensory characteristics.

The need for micronutrients (zinc, boron, iron, copper, chloride, molybdenum, and manganese) is minimal and is only partially satisfied by soil, chemical fertilisers, and other sources. Intensified agricultural practises, uneven fertiliser applications, particularly NPK, restricted use of organic manures, nutrient depletion, and a lack of replenishment are the main causes of micronutrient deficiencies. According to Jeyakumar and Balamohan [3], boron, iron, and mo deficiencies are the most common deficiencies affecting horticultural crops. Additionally, these micronutrients aid in the absorption of significant nutrients and are actively involved in all aspects of plant metabolism, including respiration, photosynthesis, chlorophyll formation, enzyme activity, hormone synthesis, nitrogen fixation and reduction, among others [4]. However, micronutrients can significantly boost horticultural crop productivity, enhance product quality, and lengthen the life of horticulture food after harvest [5]. By catalysing the effects of micronutrients, macronutrients are swiftly absorbed and used by plant tissue [6].

2. MATERIALS AND METHODS

The 2016–2017 growing season was the focus of the current study, "Effect of foliar application of borax and zinc sulphate on growth, yield, and quality of guava (*Psidium guajava* L.) cv. Gwalior–27." The current study was carried out at the Krishi Vigyan Kendra, College of Agriculture, Agrotechnology Park, Gwalior (M.P.). Guava variety "Gwalior 27" is 7 years old and was planted using a square planting scheme, 6 m apart. Throughout the trial period, all the plants were treated to the same cultural procedures in order to evaluate the impacts of various treatments. A unit of one plant was used in each of the three replications of each treatment in the experiment's Randomised Block Design layout. The treatments included two separate chemicals, namely Borax and Zinc Sulphate, in two different concentrations. Controlled amounts of ordinary

distilled water were sprayed on the plants. By sulphate and boric acid in the necessary quantity of water, the stock solution of various concentrations of zinc sulphate (neutralised with hydrated lime) and borax was created. Fruits were picked when their skins became a pale yellow. For various observations, randomly chosen branches from each treatment were tagged in all directions of the tree. Observations recorded to be physiological loss in weight, Shelf life of fruits and spoilage %. The data so obtained were analysed statically.

Table 1. List of treatments used for the study

Treatments	Details of the treatment
T ₁ (B ₀ Z ₀)	Control (Water spray)
T ₂ (B ₁)	Borax (0.2%)
T ₃ (B ₂)	Borax (0.4%)
T ₄ (Z ₁)	ZnSO ₄ (0.4%)
T ₅ (Z ₂)	ZnSO ₄ (0.6%)
T ₆ (B ₁ Z ₁)	Borax (0.2%) + ZnSO ₄ (0.4%)
T ₇ (B ₁ Z ₂)	Borax (0.2%) + ZnSO ₄ (0.6%)
T ₈ (B ₂ Z ₁)	Borax (0.4%) + ZnSO ₄ (0.4%)
T ₉ (B ₂ Z ₂)	Borax (0.4%) + ZnSO ₄ (0.6%)

3. RESULTS AND DISCUSSION

The PLW% is presented in Table 2 and the minimum physiological loss in weight was recorded 4.90 per cent at 4 days with foliar application of Zinc sulphate @ 0.6 (T₅). The maximum physiological loss in weight was recorded 7.10 per cent at 4 days under control (T₁). The maximum shelf life of fruit recorded 5.2 days with foliar spray of Zinc sulphate @ 0.6% (T₅) significantly superior to other treatments followed by T₃ and T₇. The minimum fruit storage stability is recorded 2.00 days under control (T₁).

The minimum weight loss in the fruits produced by the application of these treatments could be

dissolving the necessary quantity of zinc attributed to slower rate of respiration and transpiration from these fruits with reduced enzymatic activities and slower the biochemical changes occurring in the fruits obtained from the trees receiving these treatments. Kumar et al. [7,8,9] reported that, reduction in weight loss of guava fruits with foliar application of borax. The minimum PLW and prolonged shelf life of was observed due to foliar application of zinc sulphate [8]

The data pertaining to effect of spray of Borax, Zinc sulphate and their interaction on the fruit spoilage percentage are given in Table 3.

The minimum spoilage % were observed (3.68%) and (3.23%) on the 5th day of storage, (11.43%) and (11.12%) on the 10th day of storage, (20.96%) and (20.56%) on the 15th day storage were observed with treatments B₂(Borax @ 0.4%) and Z₂ (zinc sulphate @ 0.6%) respectively. however maximum decay per cent was recorded in the control [10,11].

The minimum spoilage % (4.11%), (11.11%) and (20.77%) was recorded with treatment B₂Z₂ (Borax@ 0.4% + ZnSo₄@ 0.6%) at 5th, 10th and 15th day of storage respectively, However, the maximum spoilage % (7.72%), (16.89%) and (28.44%) was recorded under control at 5th, 10th and 15th day of storage, respectively.

A critical examination of data presented in the Table 4 revealed that various treatments of Borax and Zinc sulphate increased the net return and B:C ratio. Maximum net returns (Rs 2, 43,280/ ha) was recorded in B₂Z₂ (Borax @ 0.4% & ZnSo₄@ 0.6 %) and the minimum (Rs 192674/ ha) in control. Whereas, minimum B: C ratio (1:8.00) was recorded with B₀Z₀

Table 2. Effect of foliar application of borax, zinc sulphate and their intraction on physiological loss in weight (%) and shelf life of guava

Treatment	Physiological loss in weight (%)	Shelf life of fruit
T ₁ -control	7.10	2.00
T ₂ -Borax (0.2%)	5.70	3.20
T ₃ -Borax(0.4%)	5.30	4.00
T ₄ -Zinc sulphate (0.4%)	5.10	3.50
T ₅ -Zinc sulphate(0.6%)	4.90	5.20
T ₆ -(Borax @0.2%+ZnSo4@0.4%)	6.00	3.00
T ₇ -(Borax @0.2%+ZnSo4@0.6%)	5.80	3.60
T ₈ -(Borax @0.4%+ZnSo4@0.4%)	5.60	3.40
T ₉ -(Borax @0.4%+ZnSo4@0.6%)	5.50	3.30
SEm±	0.077	0.120
CD at 5%	0.239	0.371

Table 3. Effect of foliar application of borax, zinc sulphate and their interaction on fruit spoilage % of guava

Treatments	Decay (%)		
	At 5 th day	At 10 th day	At 15 th day
Borax (H₃BO₃)			
B ₀	5.74	14.65	24.64
B ₁	4.92	12.37	22.46
B ₂	3.68	11.43	20.96
S.Em+	0.360	0.540	0.435
CD at 5%	1.04	1.52	1.23
ZnSO₄			
Z ₀	5.77	14.78	23.47
Z ₁	4.52	13.34	22.14
Z ₂	3.23	11.12	20.56
S.Em+	0.360	0.540	0.435
CD at 5%	1.04	1.52	1.23
B ₀ Z ₀	7.72	16.89	28.44
B ₁ Z ₁	5.33	12.68	24.34
B ₁ Z ₂	5.50	12.34	22.23
B ₂ Z ₁	4.44	13.23	22.33
B ₂ Z ₂	4.11	11.11	20.77
S.Em+	0.17	0.24	0.45
CD at 5%	0.55	0.62	1.25

Table 4. Economics of the different treatments (per ha.)

Treatments	Common expenditure/ (ha)	Treat. Cost (Rs)	Total Expenditure / ha	Gross Income/ ha	Net Income/ ha	B:C Ratio
T ₁	25000	0	25000.00	220200.00	192674.00	1:8.00
T ₂	25000	1500	26500.00	223600.00	197100.00	1:8.44
T ₃	25000	2155	27155.00	245000.00	217845.00	1:9.02
T ₄	25000	2526	25726.00	210600.00	185600.00	1:8.42
T ₅	25000	3830	28830.00	233400.00	204570.00	1:8.10
T ₆	25000	5621	30621.00	254800.00	224179.00	1:8.32
T ₇	25000	5100	30001.00	240600.00	210599.00	1:8.02
T ₈	25000	5801	28600.00	256400.00	227800.00	1:8.97
T ₉	25000	5675	27120.00	270400.00	243280.00	1:9.97

(Borax @ 0.2 % @ & no ZnSo4), while maximum (1:9.97) was in B₂Z₂ (Borax @ 0.4% & ZnSo4 @ 0.6 %). The net income and B:C ratio was calculated taking sale price of fruits @ 20/kg.

and Zinc sulphate, the treatment B₂Z₂ (Borax @ 0.4% & Zinc sulphate @ 0.6 %) were found to be the best treatments for almost Maximum net returns and maximum B: C ratio.

4. CONCLUSION

The minimum physiological loss in weight and maximum shelf life of fruit recorded with foliar spray of Zinc sulphate @ 0.6% (T₅) significantly superior to other treatments. The minimum spoilage % were observed on the 5th day of storage, 10th day of storage, and the 15th day storage were observed with treatments B₂ (Borax @ 0.4%) and Z₂ (zinc sulphate @ 0.6%) respectively. In the interaction effect of Borax

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Keram KS, Sharma BL, Kewat ML, Sharma GD. Effect of zinc fertilization on growth, yield and quality of wheat grown under agro-climatic condition of kymore

- plateau of madhyapradesh, India. The BioScan. 2014;9(4):1479-1483.
2. Anonymous; 2013. Available:<http://www.croponutrition.com/efu/micronutrients#overview>
 3. Jeyakumar P, Balamohan TN. Micronutrients for Horticultural Crop; 2013. Available:<http://agritech.tnau.ac.in>
 4. Das DK. Micronutrients: Their behaviours in soils and plants, Kalyani Publ Ludhiana. 2003;111-214.
 5. Fatima Sadia, Ruksana Nausheed, Syeda Maimuna Hussain, Israth Fatima, Nazneen Begum, and Riffath Siddi-qua. "Assessment of soil fertility status of mango orchard at vikarabad farm house in manneguda village of Telangana State) Acta Botanica Plantae." 2023: 11-17.
 6. Raja EM. Importance of micronutrients in the changing horticultural scenario. J Hort Sci. 2009;4(1):1-27.
 7. Phillips M. Economic benefits from using micronutrients for the farmer and the fertilizer producer. Int Sym Onmicr New Delhi, India. 2004;23-25.
 8. Mana PB, Waga B, Wang-Bara VYE, Mvondo S, Bourou, Palai O. "Evaluation of the agronomic and technological performance of three new cotton varieties in the cotton zone of Cameroon." Acta Botanica Plantae. *V02i01* 2023:28-39.
 9. Kumar P, Singh AK, Shankhdhar SC. Efficacy of soil and foliar application of macro and micronutrients on yield and quality of Mango Cv. Dashehari. Int J Curr Microbiol App Sci 2017;6(10):1855-1861.
 10. Goswami AK, Shukla HS, Kumar P, Mishard DS. Effect of preharvest application of micronutrients on quality of guava (*Psidium guajava* L.) Cv. Sardar. Hort Flora Res Spectrum 2012;1(1):60-63.
 11. Shreekant RD, Kumar U. Effect of foliar application of micronutrients on fruit set, yield attributes and yield of winter season Guava (*Psidium guajava* L.) Cv. L-49. Int J Pure App Biosci. 2017;5(5):1415-1419.

© 2023 Yadav et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/105394>