



Relationships between Different Growth and Yield Traits in Bottle Guord [*Lagenaria siceraria* (mol.) Standl] with Path Coefficient Analysis over Seasons under Salt Affected Soil

**Anoj Yadav^{a++*}, G. C. Yadav^{b#}, Ruchika Abha^{c++},
Vikash Singh^{b++} and Prabhakar Yadav^d**

^a Department of Vegetable Science, College of Horticulture and Forestry, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya 224229 (U.P.), India.

^b Department of Vegetable Science, College of Horticulture and Forestry, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya 224229 (U.P.), India.

^c Department of Horticulture, School of Agricultural Sciences and Technology, Babasaheb Bhimrao Ambedkar University Lucknow-226025, India.

^d Department of Agriculture, Rama University, Kanpur, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i102808

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/104396>

Original Research Article

Received: 20/06/2023

Accepted: 25/08/2023

Published: 28/08/2023

⁺⁺ Research Scholar;

[#] Professor;

*Corresponding author: E-mail: anojyadav005@gmail.com;

ABSTRACT

This study used 23 genotypes of bottle gourd—27 F₁ hybrids, 12 Parents (9 lines, and 3 testers)—during two seasons (Y₁ and Y₂) and pooled analyses to analyze the correlations between fruit production per plant (Kg), growth, and economic features. The observations were evaluated based on qualities that are related to growth and yield. It was discovered that the fruit yield per plant had exhibited a significant and positive phenotypic correlation with the length of the male and female flowers' pedicels, the number of primary branches per plant, the length of the vine, the number of nodes per vine, the internodal length, the duration of picking, the length of the peduncle, the length of the fruit, the average fruit's circumference, the average fruit's weight, the number of fruits per plant. The highest positive direct effect on fruit yield per plant was exerted by number of fruit per plant followed by average fruit weight at phenotypic level. Whereas, higher negative direct effects exerted by days to first fruit harvest. The soil type of experimental site was sandy loam with average fertility level and pH in the range of 7.5-8.5

Keywords: Correlation; fruit yield per plant; path analysis; bottle gourd.

1. INTRODUCTION

"Bottle gourd [*Lagenaria siceraria* (Mol.) Standl.] is one of the popular cucurbit vegetable crop with 2n = 2x = 22. It is an important cultivated annual cucurbitaceous crop grown throughout the country. Being warm season vegetable crop it thrives well in warm and humid climate but at present its off season cultivation has progressively stretched throughout the year in northern Indian plains. It is mainly grown for its fruits for culinary purposes and seeds which are good source of oil and protein" (Panse and Sukhatme [1]. "This delicious vegetable is also known by other names such as *bottle squash*, *calabash gourd*, *white flowered gourd*, *doodhi* and *lowki*. It is highly cross pollinated crop due to its monoecious and andromonoecious nature. Bottle gourd is the largest produced cucurbitaceous vegetables in the world preferred in both urban and rural population. In India, the total area covered under bottle gourd is 0.117 million ha with production of 2.18 million tonnes and its productivity is 18.6 tonnes per ha" [2].

2. MATERIALS AND METHODS

The research work was conducted during Zaid seasons of 2019-20 (Y₁) and 2020-21 (Y₂) to study heterosis over better-parent and standard variety using line × tester mating design at the Main Experiment Station (MES) of the Department of Vegetable Science, Acharya Narendra Deva University of Agriculture and Technology, Narendra Nagar, Kumarganj, Ayodhya (U.P.) India. The soil of this farm have more than 8 pH and alkaline in nature. The observations were recorded on twenty five characters.

The experimental materials for the present investigation comprised of nine promising and diverse inbred lines/varieties with three testers of bottle gourd selected on the basis of genetic variability from the germplasm stock maintained in the Department of Vegetable Science, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya (U.P.) India. The selected parental lines i.e.; NDBG-28 (L₁), NDBG-13 (L₂), NDBG-15 (L₃), Narendra Pooja (L₄), NDBG-104 (L₅), NDBG-Sel-1 (L₆), Narendra Kamna (L₇), NDBG-21(L₈), NDBG-22 (L₉) were crossed with three testers viz. Pusa Naveen (T₁), Narendra Prabha (T₂), Narendra Rashmi (T₃) to get 27 F₁ seed. Parental lines (9 lines and 3 testers) were also selfed/sibbed to get the true to type seeds. The present experiments were conducted in RBD with three replications to appraise the performance of 27 F₁ hybrids and their 12 parents (9 lines and 3 testers) for the study of heterobeltiosis and standard heterosis for twenty three fruit yield and quality attributing traits. The crop was sown in rows spaced at 3 meters apart with a plant to plant spacing of 0.50 meter. Sowing was done on 20 March, 2019-20 and 19 March, 2020-21. All the recommended agronomic package of practices and protection measures were followed to raise good crops. Observations were recorded on days to first male flower anthesis, days to first female flower anthesis, node number to first male flower appearance, node number to first female flower appearance, length of pedicel of male flower (cm), length of pedicel of female flower (cm), days to first harvest, primary branches per plant, vine length (m), number of node per vine, internodal length (cm), picking duration, peduncle length (cm), fruit length (cm), average fruit circumference (cm), average fruit weight (kg),

number of fruit per plant, fruit yield per plant (kg), total soluble solids (%), reducing sugars (%), non-reducing sugar (%), total sugars (%) and dry matter (g/100g). The data were subjected to analysis of variance for randomized block design as suggested by Panse and Sukhatme [1].

2.1 Statistical Analysis

As indicated by Al-Jibouri et al. [3], phenotypic and genotypic correlation coefficients have been estimated to investigate the association of various combinations of characteristics. A standard incomplete regression coefficient is the path coefficient. It provides a division of correlation coefficients into direct and indirect effects. Following Dewey and Lu [4], the path coefficient analysis of the traits that contribute to marketable green fruit yield per plant was done.

3. RESULTS AND DISCUSSION

3.1 Correlation Coefficients

Based on studies on correlation, using one character will improve all other correlated characters. Due to natural associations, many of the characters have positive or negative correlations with one another. Correlation tables' indirect correlation gets more complicated as more variables are taken into effect. The twenty-three characters under study's phenotypic and genotypic correlation coefficients are shown in Tables 1 and 2.

Fruit yield per plant had significant negative phenotypic correlation with days to first male flower anthesis, days to first female flower anthesis, days to first fruit harvest, and node number to first male flower appearance at phenotypic level during two seasons and also over seasons (pooled). Fruit yield per plant had positive and significant phenotypic correlation with number of fruits per plant, average fruit weight, vine length, number of primary branches per plant, and circumference of fruit. Many earlier researchers had also reported positive and significant correlation of most of the above traits with fruit yield per plant namely, [5,6,7,8,9,10] thereby, they also supported present findings.

Looked at these associations from findings of present research it appears that for improvement of bottle gourd, number of fruits per plant, average fruit weight, vine length, fruit

circumferences, primary branches per plant, days to first male flower anthesis, days to first female flower anthesis, days to first fruit harvest, inter nodal length and node number to first male flower appearance need to be given more consideration. A positive association of days to first male flower anthesis, days to first female flower anthesis and node number to first male flower appearance with days to first fruit harvest suggests that early flowering and flower appearance at lower node would be appropriate selection criteria to get early yield. The presence of positive correlation of number of fruits per plant with vine length and primary branches per plant revealed that longer vine length can be selected for harvesting more marketable fruits.

3.2 Path Coefficient Analysis

Finding out the direct and indirect effects of yield attributes, which are essential to choosing the best genotypes, is helpful. The estimated correlation coefficients merely show how individuals are related to one another; they do not reveal any information about the causal connections. The analysis of path coefficient, developed by Wright [11], offers a practical method for identifying both direct and indirect reasons of association. It enables a careful assessment of the precise forces at work to form a given correlation and quantifies the relative weight of each causal factor. The first study to show the value of route coefficient analysis in a breeding exertion employing relatives of crested wheat grass was Dewey and Lu in 1959. The degree of direct impacts exerted by independent variables and the impact exerted via other characters, which unavoidably arise as an inherent component of the growth pattern, govern the development of the dependent variable due to the reciprocal relationship. The complete correlation is insufficient in these complicated situations to explain the true association necessary for an efficient and successful manipulation of the characteristics. To determine the direct and indirect effects of various features on fruit yield, the route coefficient analysis was carried out using the phenotypic and genotypic correlation coefficients. Tables 1, 2, and 3 show the direct and indirect effects of numerous characteristics on fruit yield at phenotypic level. The number of fruits per plant and average fruit weight at the phenotypic level had the most positive direct effects on fruit output per plant, according to analysis of the path coefficient. The days until the first fruit harvest,

Table 1. Estimates of correlation coefficient at phenotypic level for growth, yield traits in bottle gourd during Zaid, 2020 (Y₁)

Traits																		Fruit yield/plant (kg)					
																	Number of fruits per plant						
Days to first male flower anthesis	1.00	0.98**	-0.09	0.059	-0.098	0.333**	0.997**	0.053	0.052	0.127	0.065	-0.160	0.272**	-	0.246**	0.123	0.125	0.063	-0.027	0.082	-0.16	0.12	0.12
Days to first female flower	1.00	-0.06	0.092	-0.108	0.217*	0.995**	0.069	0.065	0.152	0.073	-0.101	0.294**	-0.233*	0.241**	0.080	0.173	0.083	-0.071	-0.002	-0.13	0.10	0.10	
Node number to first male flower appearance	1.00	0.911**	0.272**	-0.030	0.055	-0.002	-0.029	-0.073	-0.014	-0.122	-0.030	-0.073	0.231*	0.279**	-0.035	-0.077	0.043	0.118	-0.23**	-0.35**	-0.42**		
Node number to first female flower appearance	1.000	0.207*	0.115	0.166	0.150	0.135	-0.010	-0.111	-	0.093	-	0.399**	0.277**	0.163	-0.053	0.019	0.067	-0.29**	-0.21*	-0.27**			
Length of pedicel of staminate flower	1.000	0.313**	-0.100	-0.181	-0.176	-	-0.041	-0.155	0.086	-0.168	-	0.357**	0.017	-0.200*	0.218*	0.278**	-0.160	-0.63**	-0.50**				
Length of pedicel of pistillate flower (cm)	1.000	0.259**	0.104	0.018	0.037	0.016	-	-0.053	-0.017	-	0.178	0.328**	0.210*	0.037	-0.071	-0.096	-0.169	-0.30**	-0.13				
Days to first harvest	1.000	0.023	-0.033	0.082	0.099	-	-0.073	0.211*	-	0.193*	0.165	0.133	0.296**	-	-0.005	0.031	-0.112	0.02	0.04				
Primary branches per plant	1.000	0.541**	0.444**	-0.053	0.062	0.301**	-	0.749**	0.575**	-0.120	0.187*	0.297**	-0.216*	-0.229*	-	0.41**	0.26**						
Vine length (m)	1.000	0.251**	-	-0.099	0.375**	0.521**	-	-	0.289**	0.515**	-	-	-	-	-0.110	0.54**	0.36**						
Number of node per vine	1.000	0.617**	0.597**	-	-	0.132	-0.075	0.255**	-0.112	0.648**	0.263**	-0.232*	-	-	-	0.357**	0.305**						
Internodal length(cm)	1.000	-0.096	-0.172	0.142	-	0.197*	-	0.159	0.245**	-	-	0.144	0.089	-0.196*	-0.14	-0.07							
Harvest duration	1.000	-0.144	-	-0.101	-	-0.161	-	-	0.590**	-	0.067	0.003	0.087	-	-0.006	0.06	0.11						
Peduncle length(cm)	1.000	0.604**	-	-0.162	0.320**	0.342**	-	0.160	-	-	0.252**	-	-	-	-0.014	0.23*	0.21*						
Fruits length (cm)	1.000	-	-	-	-	-	-	-	-	-	0.895**	0.835**	-0.133	0.10*	0.34**								
Average fruit circumference (cm)	1.000	0.560**	-	-0.050	0.669**	0.768**	0.768**	-	-	-	-	-	-	-	-	0.32**	0.20*						
Average fruit weight (kg)	1.000	-	-	-	0.585**	0.429**	-	-	-	-	0.409**	0.455**	0.445**	-	-	-	-						
Total soluble solids (%)	1.000	-	-	-	-	-	-	-	-	-	1.000	-0.116	-0.067	-0.162	-0.095	-0.56**	-0.54**						
Reducing sugar (%)	1.000	-	-	-	-	-	-	-	-	-	1.000	0.443**	-	-	-0.106	0.19*	-0.01						
Non-reducing sugar (%)	1.000	-	-	-	-	-	-	-	-	-	1.000	0.589**	0.747**	-	-	-	-						
Total sugars (%)	1.000	-	-	-	-	-	-	-	-	-	1.000	0.940**	0.994**	-0.077	0.29**	0.11							
Dry matter	1.000	-	-	-	-	-	-	-	-	-	1.000	1.211**	-0.163	-0.15	0.04								
Number of fruits per plant	1.000	-	-	-	-	-	-	-	-	-	1.000	-0.192*	-0.08	0.12									
Fruit yield/plant (kg)	1.000	-	-	-	-	-	-	-	-	-	1.000	-0.05	0.05	0.15									

*, ** Significant at 5 percent and 1 percent probability levels, respectively

Table 2. Estimates of correlation coefficient at phenotypic level for growth, yield traits in bottle gourd during Zaid, 2021 (Y₂)

Traits																							Fruit yield/plant (kg)	Number of fruits per plant
Days to first male flower anthesis	1.000	0.863**	-0.123	-0.150	-0.314**	0.076	0.764**	0.160	0.272**	0.257**	0.005	-0.125	0.245**	-0.043	0.112	-0.159	0.098	0.041	-0.040	-0.057	-0.070	0.388**	0.303**	
Days to first female flower	1.000	-0.233*	-0.214*	-0.301**	0.037	0.891**	0.237**	0.318**	0.255**	-0.036	-0.140	0.234*	-0.125	0.060	-0.272**	0.112	0.069	-0.111	0.025	-0.092	0.365**	0.187*		
Node number to first male flower appearance	1.000	0.256**	0.280**	0.066	-0.229*	-0.079	-0.094	-0.075	0.024	-0.081	-0.136	0.005	-0.053	0.127	-0.064	-0.104	0.100	-0.197*	-0.234*	-0.312**	-	0.424**	0.119	
Node number to first female flower appearance	1.000	0.078	0.016	-0.163	0.144	-0.006	0.277**	0.246**	-0.251**	0.042	0.022	-0.068	-0.111	-0.023	0.109	-0.202*	0.059	0.034	-0.008	-0.008	-0.008	-0.119		
Length of pedicel of staminate flower	1.000	0.163	-0.252**	-0.062	-0.159	-0.220*	-0.042	-0.062	0.059	-0.179	-0.020	0.408**	-0.035	-0.122	0.245**	-0.125	-0.134	-0.566**	-	0.376**	-			
Length of pedicel of pistillate flower (cm)	1.000	0.003	0.066	0.060	0.069	-0.005	-0.040	0.007	-0.261**	0.144	0.346**	0.171	0.038	-0.024	0.006	-0.049	-0.257**	-0.082	-	-	-			
Days to first harvest	1.000	0.211*	0.323**	0.259**	-0.035	-0.196*	0.150	-0.010	0.032	-0.282**	0.140	0.090	-0.103	0.080	-0.085	0.366**	0.200*	-	-	-	-	-		
Primary branches per plant	1.000	0.547**	0.368**	-0.075	0.071	0.281**	-0.201*	0.297**	-0.422**	0.098	0.260**	-0.229*	0.258**	-0.368**	0.476**	0.179	-	-	-	-	-	-		
Vine length (m)	1.000	0.235*	0.235*	-0.556**	-0.120	0.237*	0.054	0.140	-0.438**	0.208*	0.450**	-0.374**	0.448**	-0.144	0.528**	0.237*	-	-	-	-	-	-		
Number of node per vine	1.000	0.668**	-0.232*	0.087	-0.016	0.230*	-0.261**	0.225*	0.209*	-0.247**	0.234*	-0.215*	0.316**	-0.165	-	-	-	-	-	-	-	-		
Internodal length(cm)	1.000	-0.079	-0.065	-0.057	-0.107	0.111	0.016	-0.182*	0.085	-0.147	-0.102	-0.131	-0.046	-	-	-	-	-	-	-	-	-		
Harvest duration	1.000	-0.083	-0.084	-0.031	0.039	-0.118	0.073	-0.002	0.060	-0.028	0.011	0.116	-	-	-	-	-	-	-	-	-	-		
Pedicule length(cm)	1.000	-0.167	0.188*	-0.133	0.108	0.133	-0.114	0.187*	-0.028	0.166	0.103	-	-	-	-	-	-	-	-	-	-	-		
Fruits length (cm)	1.000	-0.379**	0.018	-0.073	-0.106	0.125	-0.105	0.013	0.060	-	-	-	-	-	-	-	-	-	-	-	-	-		
Average fruit circumference (cm)	1.000	-0.071	0.145	0.192*	-0.157	0.209*	-0.208*	0.206*	0.146	-	-	-	-	-	-	-	-	-	-	-	-	-		
Average fruit weight (kg)	1.000	0.032	-0.265**	0.265**	-0.305**	0.180	-0.703**	0.019	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Total soluble solids (%)	1.000	0.288**	-0.264**	0.266**	0.109	0.054	0.079	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Reducing sugar (%)	1.000	-0.820**	0.849**	0.058	0.236*	0.046	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Non-reducing sugar (%)	1.000	-0.712**	-0.124	-0.214*	0.027	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Total sugars (%)	1.000	0.034	0.239**	0.064	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Dry matter	1.000	0.008	0.257*	0.645**	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Number of fruits per plant	1.000	0.008	0.257*	0.645**	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Fruit yield/plant (kg)	1.000	0.008	0.257*	0.645**	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

*, ** Significant at 5 percent and 1 percent probability levels, respectively

Table 3. Estimates of correlation coefficient at phenotypic level for growth, yield traits in bottle gourd during Zaid, 2020-21 (pooled)

Traits																			Fruit yield/plant (kg)	Number of fruits per plant			
Days to first male flower anthesis	1.000	0.888**	-0.075	-0.068	-0.174**	0.163*	0.810**	0.122	0.227**	0.202**	-0.006	-0.082	0.230**	-0.081	0.119	-0.062	0.076	0.036	-0.018	0.008	-0.095	0.282**	0.207**
Days to first female flower appearance	1.000	-0.125	-0.099	-0.177**	0.098	0.916**	0.180*	0.268**	0.209**	-0.032	-0.062	0.232**	-0.106	0.071	-0.145*	0.072	0.059	-0.072	0.044	-0.093	0.272**	0.137*	
Node number to first male flower appearance	1.000	0.497**	0.243**	0.024	-0.091	-0.032	-0.018	-0.074	-0.034	-0.090	-0.072	0.009	0.056	0.130	-0.076	-0.047	0.066	0.004	-0.261**	-0.285**	0.373**	-0.151*	
Node number to first female flower appearance	1.000	0.098	0.038	-0.043	0.148*	0.074	0.147*	0.049	-0.246**	0.077	0.004	0.076	0.000	-0.014	0.058	-0.115	0.055	-0.107	-0.050	-0.243**	-0.074	-0.516**	-0.392**
Length of pedicel of staminate flower	1.000	0.166*	-0.152*	-0.087	-0.156*	-0.214**	-0.029	-0.102	0.061	-0.096	-0.077	0.353**	-0.056	-0.153*	0.214**	0.018	-0.128	-0.516**	-0.243**	-0.074	-0.426**	-0.121	
Length of pedicel of pistillate flower (cm)	1.000	0.093	0.090	0.080	0.069	-0.017	-0.028	-0.003	-0.229**	0.125	0.243**	0.179**	0.089	-0.046	0.030	-0.091	-0.243**	-0.074	-0.426**	-0.121	-0.377**	0.210*	
Days to first harvest	1.000	0.152*	0.240**	0.181**	-0.031	-0.078	0.149*	-0.044	0.053	-0.124	0.076	0.041	-0.044	0.093	-0.081	0.246**	0.121	-0.357**	0.437**	0.277**	0.505**	0.277**	
Primary branches per plant	1.000	0.528**	0.390**	-0.069	0.083	0.279*	-0.205**	0.333**	-0.251**	0.091	0.252**	-0.213**	0.012	-0.339**	0.072	-0.113	0.435**	0.181**	0.435**	0.050*	0.193**	0.193**	
Vine length (m)	1.000	0.231**	0.591**	-0.043	0.273**	0.014	0.220**	0.363**	0.181**	0.209**	0.273**	-0.339**	0.072	-0.233**	0.046	-0.233**	0.200**	-0.233**	0.317**	0.317**	0.193**	0.193**	
Number of node per vine	1.000	0.624**	-0.208**	0.102	-0.013	-0.013	0.173**	0.276**	0.209**	-0.173**	0.276**	-0.233**	0.046	-0.233**	0.046	-0.233**	0.200**	-0.233**	0.317**	0.317**	0.193**	0.193**	
Internodal length(cm)	1.000	-0.117	-0.103	-0.016	0.102	-0.013	-0.173**	0.276**	0.209**	-0.173**	0.276**	-0.233**	0.046	-0.233**	0.046	-0.233**	0.200**	-0.233**	0.317**	0.317**	0.193**	0.193**	
Harvest duration	1.000	-0.100	0.025	-0.059	-0.076	-0.191**	0.086	-0.011	0.094	-0.001	0.094	-0.138*	-0.138*	-0.138*	-0.138*	-0.138*	0.066	-0.024	0.191**	0.191**	0.135*	0.135*	
Pedicule length(cm)	1.000	-0.171**	0.201**	-0.116	0.112	0.134*	-0.117	0.066	-0.117	0.066	-0.117	-0.138*	-0.138*	-0.138*	-0.138*	-0.138*	0.066	-0.024	0.191**	0.191**	0.145*	0.145*	
Fruits length (cm)	1.000	-0.309**	-0.141*	-0.121	-0.180**	0.202**	0.161*	-0.055	0.121	0.212**	-0.207**	-0.069	-0.233**	-0.233**	-0.233**	-0.233**	0.231**	0.137*	0.054	0.184**	-0.529**	-0.243**	
Average fruit circumference (cm)	1.000	0.015	-0.181**	0.109	-0.187**	0.055	-0.529**	-0.055	-0.529**	-0.055	-0.529**	-0.055	-0.529**	-0.055	-0.529**	-0.055	-0.529**	-0.055	-0.529**	-0.055	-0.529**	-0.055	
Average fruit weight (kg)	1.000	0.275**	-0.283**	-0.076	0.047	0.058	0.069	1.000	0.275**	-0.283**	-0.076	0.047	0.058	0.069	1.000	0.275**	-0.283**	-0.076	0.047	0.058	0.069	1.000	
Total soluble solids (%)	1.000	-0.822**	-0.809	0.083	0.224**	0.070	1.000	0.262**	-0.142*	-0.167**	0.043	1.000	-0.067	0.036	0.096	1.000	0.262**	-0.142*	-0.167**	0.043	0.096	1.000	
Reducing sugar (%)	1.000	-0.822**	-0.809	0.083	0.224**	0.070	1.000	0.262**	-0.142*	-0.167**	0.043	1.000	-0.067	0.036	0.096	1.000	0.262**	-0.142*	-0.167**	0.043	0.096	1.000	
Non-reducing sugar (%)	1.000	-0.822**	-0.809	0.083	0.224**	0.070	1.000	0.262**	-0.142*	-0.167**	0.043	1.000	-0.067	0.036	0.096	1.000	0.262**	-0.142*	-0.167**	0.043	0.096	1.000	
Total sugars (%)	1.000	0.027	0.184**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	1.000	
Dry matter	1.000	0.027	0.184**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	1.000	
Number of fruits per plant	1.000	0.027	0.184**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	1.000	
Fruit yield/plant (kg)	1.000	0.027	0.184**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	-0.697**	1.000	

*, ** Significant at 5 percent and 1 percent probability levels, respectively

Table 4. Estimates of correlation coefficient at genotypic level for growth, yield traits in bottle gourd during Zaid, 2020 (Y1)

Traits																		Fruit yield/plant (kg)					
																		Number of fruits per plant					
																		Dry matter					
Days to first male flower anthesis	1.000	0.981**	-0.093	0.059	-0.098	0.333**	0.997*	0.053	0.052	0.127	0.065	-0.160	0.272**	-0.258**	0.246**	0.123	0.125	0.063	-0.027	0.082	-0.167	0.126	0.127
Days to first female flower	1.000	-0.067	0.092	-0.108	0.217*	0.995**	0.069	0.065	0.152	0.073	-0.101	0.294**	-0.233*	0.241**	0.080	0.173	0.083	-0.071	-0.002	-0.134	0.108	0.102	
Node number to first male flower appearance	1.000	0.911**	0.272**	-0.030	0.055	-0.002	-0.029	-0.073	-0.014	-0.122	-0.030	-0.073	0.231*	0.279**	-0.035	-0.077	0.043	0.118	-0.299**	-0.356**	-	0.419**	
Node number to first female flower appearance	1.000	0.207*	0.115	0.166	0.150	0.135	-0.010	-0.111	-0.276**	0.093	-0.405**	0.399**	0.277**	0.163	-0.053	0.019	0.067	-0.297**	-0.216*	-	0.277**	-	
Length of pedicel of staminate flower	1.000	0.313**	-0.100	-0.181	-0.176	-0.270**	-0.041	-0.155	0.086	-0.168	-0.297**	0.357**	0.017	-0.200*	0.218*	0.278**	-0.160	-0.635**	-0.508*	-	0.277**	-	
Length of pedicel of pistillate flower (cm)	1.000	0.259**	0.104	0.018	0.037	0.016	-0.053	-0.017	-0.685**	0.178	0.328**	0.210*	0.037	-0.071	-0.096	-0.169	-0.306**	-0.137	-	-	-	-	
Days to first harvest	1.000	0.023	-0.033	0.082	0.099	-0.073	0.211*	-0.266**	0.193*	0.165	0.133	0.025	-0.005	0.031	-0.112	0.028	0.048	-	-	-	-	-	-
Primary branches per plant	1.000	0.541**	0.444**	-0.053	0.062	0.301**	-0.749*	0.575**	-0.120	0.187*	0.297**	-0.216*	-0.229*	-0.383**	0.410**	0.263**	-	-	-	-	-	-	-
Vine length (m)	1.000	0.251**	-0.617**	-0.093	0.375**	-0.208*	0.521**	-0.293**	0.269**	0.515**	-0.357**	-0.305**	-0.110	0.544**	0.550**	-	-	-	-	-	-	-	-
Number of node per vine	1.000	0.597**	-0.045**	0.132	-0.075	0.255**	0.112	0.645**	0.263**	-0.232*	0.449**	-0.296**	-0.237**	-0.147	0.487***	0.276**	-	-	-	-	-	-	-
Internodal length(cm)	1.000	0.696	-0.172	0.192	-0.197*	0.159	0.245**	-0.252**	0.144	0.099	-0.196*	-0.147	-0.078	-	-	-	-	-	-	-	-	-	-
Harvest duration	1.000	-0.144	0.366**	-0.101	0.366**	-0.101	-0.161	-0.590**	0.027	0.003	0.007	-0.006	0.067	-0.067	-	-	-	-	-	-	-	-	-
Pedicule length(cm)	1.000	-0.604**	0.320**	-0.162	0.342**	0.160	-0.134	-0.184*	-0.014	0.230*	0.214*	-	-	-	-	-	-	-	-	-	-	-	-
Fruits length (cm)	1.000	-0.560**	-0.669**	-0.768**	0.895**	0.835**	-0.133	0.199*	0.342**	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Average fruit circumference (cm)	1.000	-0.050	0.585**	0.429**	-0.409**	-0.455**	0.445**	0.325**	0.202*	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Average fruit weight (kg)	1.000	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Total soluble solids (%)	1.000	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.541**
Reducing sugar (%)	1.000	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-0.004
Non-reducing sugar (%)	1.000	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.115
Total sugars (%)	1.000	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.041
Dry matter	1.000	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Number of fruits per plant	1.000	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Fruit yield/plant (kg)	1.000	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

*, ** Significant at 5 percent and 1 percent probability levels, respectively

Table 5. Estimates of correlation coefficient at genotypic level for growth, yield traits in bottle gourd during Zaid, 2021 (Y₂)

Traits																		Fruit yield/plant (kg)	Number of fruits per plant					
Days to first male flower anthesis	1.000	0.916**	-0.130	-0.177	-	0.102	0.870**	0.168	0.275**	0.251**	-0.011	-0.140	0.266**	-0.103	0.187*	-0.158	0.409**	0.053	-0.047	-0.065	-0.090	0.401**	0.342**	
Days to first female flower	1.000	-	-	-	0.374**	0.054	0.999**	0.274**	0.358**	0.275**	-0.059	-0.154	0.283**	-0.143	0.089	-	0.487**	0.086	-0.123	0.020	-0.111	0.388**	0.245**	
Node number to first male flower appearance	1.000	0.263**	0.270**	0.345**	0.356**	0.064	-	-0.074	-0.101	-0.070	0.042	-0.092	-0.153	0.039	-0.107	0.124	0.274**	-0.086	-0.114	0.102	-0.200*	-	-	-
Node number to first female flower appearance	1.000	0.098	0.041	-	-0.199*	0.142	-0.029	0.315**	0.295**	-	0.274**	-0.057	0.060	-0.022	-0.142	-0.177	0.163	-0.233*	0.072	0.014	0.004	0.325**	0.472**	-0.142
Length of pedicel of staminate flower	1.000	-	-	-	0.251**	-	-0.087	-0.184*	-	0.267**	-0.052	-0.078	0.051	-	-0.086	0.452**	-	-0.180	0.268**	-0.132	-0.142	-	0.630**	0.455**
Length of pedicel of pistillate flower (cm)	1.000	0.049	0.096	0.100	-	0.061	-0.044	-	0.053	0.034	-	0.350**	-	0.259**	0.409**	0.487**	0.058	-0.027	0.010	-0.080	-	0.292**	-0.102	-
Days to first harvest	1.000	-	-	-	0.250**	0.413**	0.292**	-0.093	-0.215*	0.187*	-0.064	0.030	-	-	0.333**	0.446**	0.109	-0.123	0.088	-0.128	0.431**	0.261**	-	
Primary branches per plant	1.000	-	-	-	-	0.561**	0.396**	-0.074	0.077	0.291**	-	-	0.633**	-	0.369**	0.293**	-	0.271**	-	-	0.491**	0.206*	-	
Vine length (m)	1.000	-	-	-	-	0.260**	-	-	1.000	-0.142	0.263**	0.057	0.270**	-	-	0.676**	0.531**	-	0.473**	-0.142	0.559**	0.256**	-	
Number of node per vine	1.000	-	-	-	-	0.562**	-	-	1.000	0.644**	-	0.111	-0.058	0.427**	-	0.483**	0.682**	0.242**	-	0.259**	-	0.342**	0.192*	-
Internodal length(cm)	1.000	-	-	-	-	0.239**	-	-	1.000	-0.059	-0.069	-0.104	0.187*	0.130	0.000	-0.228*	0.099	-0.161	-0.134	-0.019	0.015	0.149	-0.048	-
Harvest duration	1.000	-	-	-	-	0.000	-	-	1.000	-0.087	-0.099	-0.036	0.019	-	0.090	-0.013	0.056	-0.019	0.015	0.110	-	-	-	
Peduncle length(cm)	1.000	-	-	-	-	-	-	-	1.000	-	-	0.406**	-0.134	-	0.360**	0.322**	0.158	-0.118	0.203*	-0.028	0.166	0.122	-	-
Fruits length (cm)	1.000	-	-	-	-	-	-	-	1.000	-	0.247**	-	0.056**	0.100	-0.015	-0.165	-0.191*	0.188*	-0.159	0.049	0.097	0.129	-	-
Average fruit circumference (cm)	1.000	-	-	-	-	-	-	-	1.000	-	-	0.556**	-	1.040**	0.320**	-	0.324**	-	0.381**	0.180	-	-	-	
Average fruit weight (kg)	1.000	-	-	-	-	-	-	-	1.000	-	0.252**	-	1.000	0.018	-	0.281**	-	0.349**	0.206*	-	0.719**	-0.068	-	
Total soluble solids (%)	1.000	-	-	-	-	-	-	-	1.000	-	0.345**	-	1.000	0.718**	-	0.328**	0.706**	0.052	0.174	0.179	-	-	-	
Reducing sugar (%)	1.000	-	-	-	-	-	-	-	1.000	-	0.691**	-	1.000	-	0.943**	0.076	0.281**	0.036	-	-	-	-	-	
Non-reducing sugar (%)	1.000	-	-	-	-	-	-	-	1.000	-	0.929**	-	1.000	-	-	-0.131	-0.218*	0.031	-	-	-	-	-	
Total sugars (%)	1.000	-	-	-	-	-	-	-	1.000	-	0.737**	-	1.000	0.030	0.244**	0.064	-	-	-	-	-	-	-	
Dry matter	1.000	-	-	-	-	-	-	-	1.000	-	0.007	0.288**	1.000	0.007	-	0.207	-	0.693**	1.000	-	-	-	-	
Number of fruits per plant	1.000	-	-	-	-	-	-	-	1.000	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Fruit yield/plant (kg)	1.000	-	-	-	-	-	-	-	1.000	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

*, ** Significant at 5 percent and 1 percent probability levels, respectively

Table 6. Estimates of correlation coefficient at genotypic level for growth, yield traits in bottle gourd during Zaid, 2020-21 (pooled)

Traits																		Fruit yield/plant (kg)	Number of fruits per plant						
Days to first male flower anthesis	1.000	0.956**	-0.181**	-0.196**	-0.234**	0.183**	0.951**	0.300**	0.313**	0.413**	0.146*	-0.207**	0.477**	-0.131*	0.287**	-0.214**	0.708**	0.154*	-0.137*	0.029	-0.191**	0.306**	0.266**		
Days to first female flower	1.000	-0.186**	-0.178**	-0.228**	0.153**	1.008**	0.441**	0.413**	0.487**	0.135*	-0.165*	0.526**	-0.114*	0.223**	-0.335**	0.811**	0.075	-0.070	0.170*	-0.278**	0.326**	0.239**			
Node number to first male flower appearance	1.000	0.721**	0.434**	0.057	-0.164*	-0.080	-0.187**	-0.121	0.029	-0.212**	-0.221**	-0.089	0.134*	0.291**	-0.159*	-0.081	0.018	-0.037	-0.305**	-0.523**	-	0.586**			
Node number to first female flower appearance	1.000	0.197**	0.031	-0.132*	0.030	-0.256**	0.218**	0.382**	-0.372**	-0.014	-0.187**	0.306**	0.167*	0.105	0.374**	-0.589**	-0.440**	-0.198**	-0.322**	-	0.341**				
Length of pedicel of staminate flower						1.000	0.504**	-0.305**	-0.283**	-0.380**	-0.329**	0.018	-0.118	0.060	-0.332**	-0.183**	0.633**	-0.192**	-0.235**	0.225**	-0.155*	-0.077	-0.821**	-0.669**	
Length of pedicel of pistillate flower (cm)						1.000	0.153*	0.111	-0.085	-0.026	0.087	0.131*	0.041	-0.571**	0.264**	0.495**	0.599**	0.205**	-0.179**	-0.113	-0.101	-0.465**	-	0.290**	
Days to first harvest						1.000	0.429**	0.416**	0.503**	0.138*	-0.216**	0.444**	-0.082	0.181**	-0.368**	0.796**	0.070	-0.075	0.201**	-0.293**	0.323**	0.240**			
Primary branches per plant						1.000	0.670**	0.779**	0.210**	0.187**	0.460**	-0.293**	0.764**	-0.417**	0.507**	0.489**	-0.397**	-0.200**	-0.675**	0.592**	0.389**				
Vine length (m)						1.000	0.468**	-0.395**	-0.029	0.279**	0.101	0.383**	-0.489**	0.355**	0.385**	-0.088	0.194**	-0.111	0.667**	0.597**					
Number of node per vine						1.000	0.612**	-0.219**	0.399**	0.397**	0.147*	0.523**	-0.375*	0.704**	0.463**	-0.489**	-0.043	-0.352**	0.535**	0.333**	0.403**				
Internodal length(cm)						1.000	-0.184**	0.169*	-0.200**	0.184**	0.057	0.364**	0.066	-0.374**	-0.127	-0.329**	-0.044	-	0.186**						
Harvest duration											1.000	-0.077	-0.009	0.177**	-0.199**	-0.439**	0.231**	-0.043	0.235**	-0.270**	0.188**	0.142**			
Pedicule length(cm)											1.000	-0.557**	0.610**	-0.144*	0.396**	0.292**	-0.304**	-0.207**	-0.070	0.361**	0.329**				
Fruits length (cm)											1.000	-0.471**	-0.359**	-0.508**	-0.331**	0.343**	0.253**	0.071	0.223**	0.216**					
Average fruit circumference (cm)											1.000	-0.020	0.746**	0.704**	0.740**	-0.726**	-0.228**	-0.348**	0.391**	0.403**					
Average fruit weight (kg)											1.000	0.134*	-0.077	-0.003	-0.208**	0.155*	-0.755**	-							
Total soluble solids (%)											1.000	0.320**	-0.383**	-0.167*	0.094	0.185**	0.144**								
Reducing sugar (%)											1.000	-0.967**	-0.471**	0.110	0.382**	0.308**									
Non-reducing sugar (%)											1.000	0.245**	0.066	-0.271**	0.066	-0.271**	-								
Total sugars (%)												1.000	-0.269**	-0.067	-						0.202**				
Dry matter													1.000	-0.078	0.106						0.208**				
Number of fruits per plant													1.000	0.903**											
Fruit yield/plant (kg)														1.000											

*, ** Significant at 5 percent and 1 percent probability levels, respectively

Table 7. Estimates of direct and indirect effects of different growth and yield traits on fruit yield per plant of bottle gourd at genotypic level during Zaid (Y_1)

Traits	Fruit yield/plant (kg)	Number of fruits per plant	Dry matter
Days to first male flower anthesis	-0.180	0.010	0.016
Days to first female flower	-0.165	0.021	0.032
Node number to first male flower appearance	0.023	-0.005	-0.121
Node number to first female flower appearance	0.032	-0.006	-0.042
Length of pedicel of staminate flower	0.067	-0.007	-0.042
Length of pedicel of pistillate flower (cm)	-0.018	0.001	-0.008
Days to first male flower appearance	-0.036	0.022	0.021
Days to first female flower	-0.055	0.022	-0.011
Node number to first male flower	0.070	-0.021	-0.013
Node number to first female flower	0.070	-0.021	-0.013
Length of pedicel of staminate flower	-0.051	-0.020	-0.006
Length of pedicel of pistillate flower	-0.015	-0.008	-0.008
Days to first harvest	-0.157	0.021	0.033
Primary branches per plant	-0.030	0.006	0.009
Vine length (m)	-0.050	0.007	0.012
Number of node per vine	-0.045	0.006	0.008
Internodal length(cm)	0.002	-0.001	-0.005
Harvest duration	0.025	-0.003	0.011
Peduncle length(cm)	-0.048	0.006	0.019
Fruits length (cm)	0.019	-0.003	-0.005
Average fruit circumference (cm)	-0.034	0.002	0.013
Average fruit weight (kg)	0.029	-0.006	-0.015
Total soluble solids (%)	-0.074	0.010	0.010
Reducing sugar (%)	-0.010	0.002	0.014
Non-reducing sugar (%)	0.009	-0.003	-0.012
Total sugars (%)	0.012	0.000	0.024
Dry matter	0.016	-0.002	0.032
Number of fruits per plant	-0.072	0.008	0.039
Fruit yield/plant (kg)	0.001	0.001	0.038

* , ** Significant at 5 percent and 1 percent probability levels, respectively

Table 8. Estimates of direct and indirect effects of different growth and yield traits on fruit yield per plant of bottle gourd at genotypic level during Zaid , 2021 (Y₂)

Traits	Fruit yield/plant (kg)	Number of fruits per plant	Dry matter	Total sugars (%)	Non-reducing sugar (%)	Reducing sugar (%)	Total soluble solids (%)	Average fruit weight (kg)	Average fruit circumference (cm)														
Days to first male flower anthesis	-0.180	0.019	0.016	-0.036	0.023	-0.021	0.146	-0.024	-0.235	0.288	0.015	-0.028	0.017	0.014	0.009	-0.160	0.003	-0.001	-0.014	-0.008	-0.003	0.502	0.342**
Days to first female flower	-0.165	0.021	0.032	-0.055	0.022	-0.011	0.167	-0.038	-0.305	0.316	0.076	-0.031	0.018	0.019	0.004	-0.277	0.004	-0.002	-0.037	0.002	-0.003	0.487	0.245**
Node number to first male flower appearance	0.023	-0.005	-0.121	0.070	-0.021	-0.013	-0.046	0.010	0.086	-0.080	-0.055	-0.019	-0.010	-0.005	-0.005	0.125	-0.001	0.002	0.031	-0.025	-0.008	-0.408	-
Node number to first female flower appearance	0.032	-0.006	-0.042	0.202	-0.006	-0.008	-0.033	-0.020	0.025	0.362	-0.381	-0.059	0.004	-0.008	-0.001	-0.143	-0.001	-0.003	-0.070	0.009	0.000	0.006	0.472**
Length of pedicel of staminate flower	0.067	-0.007	-0.042	0.020	-0.061	-0.051	-0.070	0.012	0.157	-0.306	0.067	-0.016	0.003	0.047	-0.004	0.457	-0.003	0.003	0.080	-0.016	-0.004	-0.790	-
Length of pedicel of pistillate flower (cm)	-0.018	0.001	-0.008	0.008	-0.015	-0.020	0.008	-0.014	-0.085	0.070	0.056	-0.011	0.002	0.053	0.012	0.413	0.004	-0.001	-0.008	0.001	-0.002	-0.366	-0.102
Days to first harvest	-0.157	0.021	0.033	-0.040	0.025	-0.010	0.168	-0.035	-0.353	0.335	0.120	-0.044	0.012	0.009	0.001	-0.336	0.003	-0.002	-0.037	0.011	-0.004	0.540	0.261**
Primary branches per plant	-0.030	0.006	0.009	0.029	0.005	-0.020	0.042	-0.140	-0.479	0.455	0.096	0.016	0.018	0.048	0.000	-0.443	0.003	-0.005	-0.011	0.033	-0.011	0.615	0.206*
Vine length (m)	-0.050	0.007	0.012	-0.006	0.011	-0.020	0.069	-0.079	-0.853	0.295	0.724	-0.029	0.017	-0.008	0.013	-0.488	0.005	-0.003	-0.116	0.058	-0.004	0.701	0.256**
Number of node per vine	-0.045	0.006	0.008	0.064	0.016	-0.012	0.049	-0.055	-0.222	1.147	-0.830	-0.048	0.007	0.008	0.020	-0.297	0.006	-0.004	-0.078	0.032	-0.007	0.429	0.192*
Internodal length(cm)	0.002	-0.001	-0.005	0.060	0.003	0.009	-0.016	0.010	0.479	0.739	-1.289	-0.012	-0.004	0.014	0.009	0.131	0.000	0.004	0.030	-0.020	-0.004	-0.187	-0.048
Harvest duration	0.025	-0.003	0.011	-0.059	0.005	0.011	-0.036	-0.011	0.121	-0.274	0.076	0.202	-0.013	0.013	-0.002	0.019	-0.003	-0.004	0.007	-0.001	0.019	0.110	-
Pedicule length(cm)	-0.048	0.006	0.010	0.171	-0.003	-0.007	0.031	-0.041	-0.255	0.127	0.089	-0.118	0.063	0.033	0.019	-0.135	0.003	-0.003	-0.036	-0.025	-0.001	0.209	0.122
Fruits length (cm)	0.019	-0.003	-0.005	0.012	0.021	0.079	-0.011	0.049	-0.049	0.067	0.135	-0.020	-0.016	-0.135	-0.026	-0.015	-0.001	0.003	0.056	-0.020	0.001	0.121	0.129
Average fruit circumference (cm)	-0.034	0.002	0.013	-0.005	0.005	-0.053	0.005	-0.089	-0.230	0.490	-0.241	-0.007	0.026	0.075	0.004	-0.255	0.008	-0.006	-0.082	0.000	-0.010	0.478	0.180
Average fruit weight (kg)	0.023	-0.006	-0.015	-0.029	-0.028	-0.083	-0.056	0.06	0.412	-0.337	-0.167	0.004	-0.009	0.02	-0.012	1.017	0.000	0.006	0.084	-0.040	0.006	-0.901	-0.068
Total soluble solids (%)	-0.074	0.010	0.010	-0.036	0.021	-0.069	0.058	-0.052	-0.077	0.783	0.000	-0.073	0.021	0.022	0.044	0.019	0.008	-0.012	-0.206	0.007	0.002	0.218	0.179
Reducing sugar (%)	-0.010	0.002	0.014	0.030	0.011	-0.012	0.018	-0.041	-0.453	0.277	0.294	0.018	0.010	0.026	0.015	-0.348	0.000	-0.017	-0.278	0.116	0.002	0.352	0.036
Non-reducing sugar (%)	0.009	-0.003	-0.012	-0.047	-0.016	0.006	-0.021	0.033	0.332	-0.269	-0.128	-0.003	-0.008	-0.025	-0.013	0.284	-0.006	0.016	0.299	-0.091	-0.004	-0.273	0.031
Total sugars (%)	0.012	0.000	0.024	0.015	0.008	-0.002	0.015	-0.038	-0.403	0.297	0.208	0.011	0.013	0.022	0.015	-0.331	0.006	-0.016	-0.220	0.123	0.001	0.305	0.064
Dry matter	0.016	-0.002	0.032	0.003	0.009	0.016	-0.021	0.055	0.121	-0.294	0.173	-0.004	-0.002	-0.007	-0.016	0.208	0.000	-0.001	-0.039	0.004	0.028	0.009	0.288**
Number of fruits per plant	-0.072	0.008	0.039	0.001	0.038	0.059	0.072	-0.069	-0.477	0.393	0.192	0.003	0.011	-0.013	0.018	-0.726	0.001	-0.005	-0.065	0.030	0.000	1.254	0.693**
Fruit yield/plant (kg)																							

*, ** Significant at 5 percent and 1 percent probability levels, respectively

Table 9. Estimates of direct and indirect effects of different growth and yield traits on fruit yield per plant of bottle gourd at genotypic level during Zaid, 2020-21 (pooled)

Traits	Fruit yield/plant (kg)	Dry matter	Number of fruits per plant																				
Total sugars (%)																							
Total soluble solids (%)																							
Average fruit weight (kg)																							
Average fruit circumference (cm)																							
Fruits length (cm)																							
Peduncle length(cm)																							
Harvest duration																							
Internodal length(cm)																							
Vine length (m)																							
Number of node per vine																							
Primary branches per plant																							
Length of pedicel of staminate flower																							
Length of pedicel of pistillate flower (cm)																							
Days to first harvest																							
Node number to first male flower appearance																							
Node number to first female flower appearance																							
Days to first male flower anthesis	0.263	2.192	-0.082	0.217	0.128	-0.187	-1.744	0.662	1.242	-2.162	0.639	0.005	-0.210	-0.002	-0.123	0.065	-0.111	-0.059	0.249	-0.016	-0.246	-0.452	0.266**
Days to first female flower	0.251	2.293	-0.085	0.196	0.125	-0.157	-1.848	0.972	1.635	-2.550	0.592	0.004	-0.232	-0.002	-0.096	0.102	-0.128	-0.029	0.127	-0.093	-0.358	-0.482	0.239**
Node number to first male flower appearance	-0.048	-0.426	0.456	-0.796	-0.238	-0.058	0.300	-0.176	-0.740	0.632	0.127	0.005	0.097	-0.002	-0.058	-0.089	0.025	0.031	-0.032	0.020	-0.393	0.774	-0.586**
Node number to first female flower appearance	-0.052	-0.407	0.329	-1.104	-0.108	-0.032	0.242	0.065	-1.013	-1.141	1.680	0.008	0.006	-0.003	-0.131	-0.051	-0.016	-0.144	1.068	0.241	-0.254	0.476	-0.341**
Length of pedicel of staminate flower	-0.061	-0.522	0.198	-0.217	-0.548	-0.516	0.558	-0.623	-1.508	1.721	0.080	0.003	-0.027	-0.006	0.079	-0.193	0.030	0.091	-0.407	0.085	-0.099	1.215	-0.669**
Length of pedicel of pistillate flower (cm)	0.048	0.352	0.026	-0.034	-0.276	-1.024	-0.280	0.244	-0.336	0.135	0.383	-0.003	-0.018	-0.011	-0.114	-0.151	-0.094	-0.079	0.324	0.062	-0.130	0.688	-0.290**
Days to first harvest	0.250	2.312	-0.075	0.146	0.167	-0.156	-1.833	0.946	1.650	-2.633	0.606	0.005	-0.196	-0.002	-0.078	0.112	-0.125	-0.027	0.135	-0.110	-0.376	-0.478	0.240**
Primary branches per plant	0.079	1.011	-0.036	-0.033	0.155	-0.113	-0.786	2.206	2.655	-4.078	0.925	-0.004	-0.203	-0.005	-0.329	0.127	-0.080	-0.189	0.721	0.110	-0.868	-0.875	0.389**
Vine length (m)	0.082	0.946	-0.085	0.282	0.208	0.087	-0.763	1.478	3.963	-2.450	-1.736	0.001	-0.123	0.002	-0.164	0.149	-0.056	-0.149	0.160	-0.106	-0.143	-0.987	0.597**
Number of node per vine	0.108	1.117	-0.055	-0.240	0.180	0.026	-0.922	1.719	1.855	-5.235	2.689	0.005	-0.176	-0.003	-0.225	0.115	-0.111	-0.179	0.886	0.023	-0.453	-0.792	0.333**
Internodal length(cm)	0.038	0.309	0.013	-0.422	-0.010	-0.089	-0.253	0.464	-1.565	-3.204	4.395	0.004	-0.074	-0.004	-0.079	-0.018	-0.057	-0.025	0.678	0.070	-0.423	0.066	-0.186**
Harvest duration	-0.054	-0.379	-0.097	0.410	0.064	-0.134	0.395	0.413	-0.116	1.146	-0.807	-0.023	0.034	0.000	-0.076	0.061	0.069	-0.089	0.078	-0.129	-0.347	-0.278	0.142**
Peduncle length(cm)	0.125	1.205	-0.101	0.016	-0.033	-0.042	-0.814	1.015	1.104	-2.086	0.741	0.002	-0.441	-0.010	-0.262	0.044	-0.062	-0.113	0.552	0.113	-0.090	-0.534	0.329**
Fruits length (cm)	-0.034	-0.260	-0.040	0.206	0.182	0.585	0.150	-0.647	0.399	0.770	-0.878	0.000	0.246	0.019	0.202	0.110	0.080	0.128	-0.622	-0.139	0.091	-0.330	0.216**
Average fruit circumference (cm)	0.075	0.510	0.061	-0.337	0.100	-0.271	-0.331	1.688	1.517	-2.737	0.809	-0.004	-0.269	-0.009	-0.430	0.006	-0.117	-0.272	1.315	0.125	-0.447	-0.579	0.403**
Average fruit weight (kg)	-0.056	-0.768	0.133	-0.185	-0.347	-0.507	0.674	-0.921	-1.936	1.963	0.252	0.005	0.063	-0.007	0.009	-0.305	-0.021	0.030	0.005	0.114	0.200	1.117	-0.491**
Total soluble solids (%)	0.186	1.861	-0.073	-0.115	0.105	-0.614	-1.460	1.117	1.406	-3.685	1.600	0.010	-0.174	-0.009	-0.321	-0.041	-0.157	-0.124	0.694	0.092	0.121	-0.274	0.144**
Reducing sugar (%)	0.040	0.172	-0.037	-0.413	0.129	-0.210	-0.129	1.079	1.526	-2.423	0.288	-0.005	-0.129	-0.006	-0.302	0.024	-0.050	-0.386	1.753	0.094	-0.142	-0.564	0.308**
Non-reducing sugar (%)	-0.036	-0.161	0.008	0.650	-0.123	0.183	0.137	-0.877	-0.349	2.559	-1.643	0.001	0.134	0.006	0.312	0.001	0.060	0.373	-1.813	-0.134	0.110	0.401	-0.202**
Total sugars (%)	0.008	0.390	-0.017	0.485	0.085	0.115	-0.368	-0.442	0.767	0.223	-0.559	-0.005	0.091	0.005	0.098	0.063	0.026	0.068	-0.445	-0.548	-0.346	0.099	-0.208**
Dry matter	-0.050	-0.637	-0.139	0.218	0.042	0.103	0.536	-1.489	-0.441	1.845	-1.444	0.006	0.031	0.001	0.149	-0.047	-0.015	0.043	-0.155	0.147	1.287	0.116	0.106
Number of fruits per plant	0.080	0.747	-0.239	0.355	0.450	0.476	-0.592	1.305	2.643	-2.802	-0.195	-0.004	-0.159	0.004	-0.168	0.231	-0.029	-0.147	0.491	0.037	-0.100	-1.479	0.903**
Fruit yield/plant (kg)																							

*, ** Significant at 5 percent and 1 percent probability levels, respectively

Table 10. Estimates of direct and indirect effects of different growth and yield traits on fruit yield per plant of bottle gourd at phenotypic level during Zaid 2020 (Y_1)

Traits	Fruit yield/plant (kg)	Number of fruits per plant	Dry matter	Total sugars (%)	Reducing sugar (%)	Non-reducing sugar (%)																	
Days to first male flower anthesis	-0.056	-0.044	0.014	-0.010	0.007	-0.006	0.066	-0.004	-0.145	0.174	-0.004	-0.013	0.015	0.000	-0.001	-0.150	-0.002	0.000	-0.007	-0.007	-0.004	0.479	0.086
Days to first female flower	-0.048	-0.052	0.027	-0.014	0.006	-0.003	0.077	-0.006	-0.169	0.173	0.026	-0.015	0.014	0.000	-0.001	-0.256	-0.002	0.000	-0.021	0.003	-0.005	0.451	0.063
Node number to first male flower appearance	0.007	0.012	-0.116	0.017	-0.006	-0.005	-0.020	0.002	0.050	-0.051	-0.018	-0.009	-0.008	0.000	0.001	0.119	0.001	0.000	0.019	-0.022	-0.012	-0.385	-
Node number to first female flower appearance	0.008	0.011	-0.030	0.066	-0.002	-0.001	-0.014	-0.004	0.003	0.188	-0.179	-0.027	0.003	0.000	0.001	-0.104	0.000	0.000	-0.038	0.007	0.002	-0.010	-0.215*
Length of pedicel of staminate flower	0.018	0.016	-0.033	0.005	-0.021	-0.012	-0.022	0.002	0.085	-0.149	0.030	-0.007	0.004	0.000	0.000	0.382	0.001	-0.001	0.046	-0.014	-0.007	-0.699	-
Length of pedicel of pistillate flower	-0.004	-0.002	-0.008	0.001	-0.003	-0.075	0.000	-0.002	-0.032	0.047	0.004	-0.004	0.000	0.000	-0.002	0.325	-0.003	0.000	-0.005	0.001	-0.002	-0.317	-0.088
(cm)																							
Days to first harvest	-0.043	-0.046	0.027	-0.011	0.005	0.000	0.086	-0.006	-0.172	0.176	0.026	-0.021	0.009	0.000	0.000	-0.265	-0.003	0.000	-0.019	0.009	-0.004	0.451	0.016
Primary branches per plant	-0.009	-0.012	0.009	0.009	0.001	-0.005	0.018	-0.027	-0.291	0.249	0.055	0.008	0.017	0.000	-0.004	-0.396	-0.002	0.001	-0.043	0.029	-0.019	0.588	0.237**
Vine length (m)	-0.015	-0.016	0.011	0.000	0.003	-0.005	0.028	-0.015	-0.533	0.159	0.406	-0.013	0.014	0.000	-0.002	-0.411	-0.004	0.002	-0.070	0.051	-0.007	0.652	0.322*
Number of node per vine	-0.014	-0.013	0.009	0.018	0.005	-0.005	0.022	-0.010	-0.125	0.677	-0.488	-0.025	0.005	0.000	-0.003	-0.245	-0.004	0.001	-0.046	0.027	-0.011	0.390	0.222*
Internodal length(cm)	0.000	0.002	-0.003	0.016	0.001	0.000	-0.003	0.002	0.296	0.452	-0.731	-0.008	-0.004	0.000	-0.001	0.104	0.000	-0.001	0.016	-0.017	-0.005	-0.162	-0.086
Harvest duration	0.007	0.007	0.009	-0.016	0.001	0.003	-0.017	-0.002	0.064	-0.157	0.058	0.107	-0.005	0.000	0.000	0.037	0.002	0.000	0.000	0.007	-0.001	0.013	0.125
Harvest duration	0.007	0.007	0.009	-0.016	0.001	0.003	-0.017	-0.002	0.064	-0.157	0.058	0.107	-0.005	0.000	0.000	0.037	0.002	0.000	0.000	0.007	-0.001	0.013	0.125
Pedicule length(cm)	-0.014	-0.012	0.016	0.003	-0.001	-0.001	0.013	-0.008	-0.126	0.059	0.048	-0.009	0.060	0.000	-0.002	-0.125	-0.002	0.001	-0.021	0.021	-0.001	0.205	0.186*
Fruits length (cm)	0.002	0.006	-0.001	0.001	0.004	0.020	-0.001	0.005	-0.029	-0.011	0.042	-0.009	-0.010	-0.001	0.005	0.017	0.001	0.000	0.023	-0.012	0.001	0.074	0.237*
Average fruit circumference (cm)	-0.006	-0.003	0.006	-0.004	0.000	-0.011	0.003	-0.008	-0.074	0.156	-0.078	-0.003	0.011	0.000	-0.012	-0.067	-0.003	0.001	-0.029	0.024	-0.010	0.254	0.138
Average fruit weight (kg)	0.009	0.014	-0.015	-0.007	-0.009	-0.026	-0.024	0.011	0.233	-0.177	-0.081	0.004	-0.008	0.000	0.001	0.938	-0.001	-0.001	0.049	-0.035	0.009	-0.868	-
Total soluble solids (%)	-0.006	-0.006	0.007	-0.002	0.001	-0.013	0.012	-0.003	-0.111	0.153	-0.012	-0.013	0.007	0.000	-0.002	0.030	-0.019	0.001	-0.049	0.030	0.006	0.066	0.059
Reducing sugar (%)	-0.002	-0.004	0.012	0.007	0.003	-0.003	0.008	-0.007	-0.245	0.142	0.133	0.008	0.008	0.000	-0.002	-0.249	-0.006	0.004	-0.152	0.096	0.003	0.292	0.110
Non-reducing sugar (%)	0.002	0.006	-0.012	-0.013	-0.005	0.002	-0.009	0.006	0.199	-0.167	-0.062	0.000	-0.007	0.000	0.002	0.249	0.005	-0.003	0.186	-0.081	-0.006	-0.264	0.050
Total sugars (%)	0.003	-0.001	0.023	0.004	0.003	-0.001	0.007	-0.007	-0.238	0.159	0.107	0.006	0.011	0.000	-0.003	-0.286	-0.005	0.003	-0.132	0.113	0.002	0.295	0.118
Dry matter	0.004	0.005	0.027	0.002	0.003	0.004	-0.007	0.010	0.077	-0.146	0.075	-0.003	-0.002	0.000	0.002	0.169	-0.002	0.000	-0.023	0.004	0.050	0.010	0.103
Number of fruits per plant	-0.022	-0.019	0.036	-0.001	0.012	0.019	0.032	-0.013	-0.281	0.214	0.096	0.001	0.010	0.000	-0.002	-0.660	-0.001	0.001	-0.040	0.027	0.000	1.234	0.762**
Fruit yield/plant (kg)																							

*, ** Significant at 5 percent and 1 percent probability levels, respectively

Table 11. Estimates of direct and indirect effects of different growth and yield traits on fruit yield per plant of bottle gourd at phenotypic level during Zaid 2021 (Y_2)

Traits	Fruit yield/plant (kg)	Number of fruits per plant
Dry matter	Total sugars (%)	Non-reducing sugar (%)
Days to first male flower anthesis	-0.056	-0.044
Days to first female flower	-0.048	-0.052
Node number to first male flower appearance	0.007	0.012
Node number to first female flower appearance	-0.116	-0.014
Length of pedicel of staminate flower	0.008	0.011
Length of pedicel of pistillate flower	0.018	0.016
Length of pedicel of pistillate flower (cm)	-0.004	-0.002
Days to first harvest	-0.043	-0.046
Primary branches per plant	-0.009	-0.012
Vine length (m)	0.009	0.009
Number of node per vine	-0.015	-0.016
Internodal length(cm)	-0.014	-0.013
Harvest duration	0.000	0.002
Pedicule length(cm)	-0.003	-0.003
Number of nodes per vine	0.007	0.007
Vine length (m)	0.001	0.001
Primary branches per plant	-0.001	-0.001
Days to first harvest	0.000	0.000
Length of pedicel of pistillate flower (cm)	-0.001	-0.001
Length of pedicel of staminate flower	0.000	0.000
Node number to first female flower appearance	-0.016	-0.014
Node number to first male flower appearance	0.014	-0.010
Days to first female flower	-0.007	-0.006
Days to first male flower	-0.048	-0.052
Days to first male flower anthesis	-0.056	-0.044
Total soluble solids (%)	-0.006	-0.006
Reducing sugar (%)	-0.002	-0.004
Non-reducing sugar (%)	0.002	0.006
Total sugars (%)	0.003	-0.001
Dry matter	0.004	0.005
Number of fruits per plant	-0.022	-0.019
Fruit yield/plant (kg)	0.036	-0.001

*; ** Significant at 5 percent and 1 percent probability levels, respectively

Table 12. Estimates of direct and indirect effects of different growth and yield traits on fruit yield per plant of bottle gourd at phenotypic level during Zaid 2020-21 (pooled)

* , ** Significant at 5 percent and 1 percent probability levels, respectively

however, have a detrimental direct effect. Number of primary branches per plant as well as the number of fruits per plant showed a positive relationship with fruit yield per plant at the phenotypic level. However, negative indirect effects via average fruit weight had an influence on this relationship. The number of fruits per plant had a positive indirect effect on circumference of fruit, which was positively correlated with fruit output per plant. Fruit output per plant shown a favorable link with fruit circumference via fruit production per plant. The connection between fruit output per plant and fruit production per plant was positive. As determined by the quantity of fruits per plant, fruit production per plant showed a favorable connection. The number of fruits per plant and average fruit weight showed a favorable correlation between vine length and fruit yield per plant. Average fruit weight, which exhibited a strong positive direct effect on fruit yield per plant at the phenotypic level, was the primary contributor to the significant positive correlation. Plant fruit output significantly correlated negatively with the number of days before the first male flower anthesis. When this link was broken down, it became clear that the indirect impacts caused by the quantity of fruits produced by each plant were mostly to blame. Days to first female flower with fruit yield per plant revealed a significant inverse relationship with days to first female flower. It was mostly because of the indirect effect of the number of fruits per plant that the node number to the first male flower appearance showed a substantial negative association with fruit production per plant. The relationship between fruit yield per plant and days before the first fruit harvest was negatively significant. When this negative correlation was broken down, it became clear that the amount of fruits produced by each plant was mostly to blame for this association. Internodal length revealed a negative, statistically significant correlation with fruit output per plant at the phenotypic level via the indirect effect of the number of fruits per plant. Positive direct effect of various traits on fruit yield has also been reported by earlier workers viz., for average fruit weight [5,6,7,12] for number of fruits per plant [8,9,13,10] which substantiate the present findings. The path coefficient analysis showed that the genotypic level direct and indirect effects were significantly different from those at the phenotypic level, which may be because the environment influences the various traits under study to various degrees. Component variance analysis and correlation tests conducted in all

three seasons and habitats confirmed these findings. Last but not least, the path coefficient analysis showed that increasing the number of fruits per plant and average fruit weight would improve the bottle gourd plant's overall output per plant [14-17].

4. CONCLUSION

The study well characterized the relationships between different growth and yield traits in bottle gourd with path coefficient analysis over seasons under salt affected soil. The presence of a positive link between the number of fruits per plant and the number of primary branches per plant suggested that greater vine length can be chosen to harvest more marketable fruits.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Panse VG, Shukhatme PV. Statistical methods for agriculture workers (2nd Eds.). Indian Council of Agricultural Research, New Delhi. 1967;328.
2. Anonymous; 2018. Available:<http://www.agriculturalproductsindia.com/vegetables/vegetables-bottle-gourd-html>
3. Al-Jibouri HA, Miller PA, Robinson HF. Genetic and environmental variances and covariance in upland cotton cross of interspecific origin. Agron. J. 1958;50:633-637.
4. Dewey DR Lu KH. A. Correlation and path analysis of components of crested wheat grass seed production. Agronomy Journal. 1959;51:515-518.
5. Ghuge MB, Syamal MM. Karcho S. Heterosis in bottle gourd [*Lagenaria siceraria* (Mol.) Standl]. Indian J. Agric. Res. 2016;50(5):466-470.
6. Gautam DK, Yadav GC, Kumar P, Kumar Vimlesh, Singh M. Estimation of heterosis for growth, yield and quality traits in bottle gourd [*Lagenaria Siceraria* (Mol.) Standl]. Int. J. Curr. Microbiol. App. Sci. 2017; 6(8):789-802.
7. Malviya AV, Bhandari DR, Tank RV, Patel AI, Patel UV. Combining ability and gene action studies for fruit yield and its components in bottle gourd [*Lagenaria*

8. *siceraria* (Mol.) Standl]. Trend. Biosci. 2017;10(2):758-762.
9. Niva D, Patel JN, Maibam U. Combining ability studies in bottle gourd (*Lagenaria siceraria* (Mol.) Standl). Int. J. Agril. Sci. Res. 2018;7(5):33-38.
10. Quamruzzaman A, Ahmad S. Genetic analysis of some yield components of bottle gourd [*Lagenaria siceraria* (Mol.) Standl]. SAARC J. Agric. 2020;8(1):1-9.
11. Geeta ON, Patel JB, Jyoti GJ. Studies on heterosis and inbreeding depression in bottle gourd [*Lagenaria siceraria* (Mol.) Standl]. Ind. J. Pure App. Biosci. 2021; 9(1):132-139.
12. Wright S. Systems of mating. Genetica. 1921;16:111-179.
13. Padmakshi T. Heterosis and combining ability for yield attributing traits in bottle gourd (*Lagenaria siceraria* (Mol.) Standl). Ph. D. (Horti) Thesis, Indira Gandhi Vishwavidyalaya, Raipur; 2017. Available:<http://krishikosh.egranthac>.
14. Kumar V, Mishra DP, Yadav GC, Babu U. Determining relationships between different growth and yield traits in pumpkin with path coefficient analysis. The Pharma Innovation Journal 2017;6(12):18-26
15. Anoj Yadav, GC Yadav. Studies on heritability and genetic advance for the quantitative traits in bottle gourd [*Lagenaria siceraria* (Mol.) Standl] over seasons under salt affected soil. J. Pharm. Innov. 2022;11(2):1630-1632.
16. Anoj Yadav, GC Yadav. Evaluation of performance for the quantitative traits in bottle gourd [*Lagenaria siceraria* (Mol.) Standl] over seasons under salt affected soil. J. Pharm. Innov. 2022;11(6):346-355.
17. Anoj Yadav, Yadav GC, Abha R, Singh V, Yadav P. Estimation of Heterosis for Yield and Quality Traits in Bottle Gourd [*Lagenaria siceraria* (mol.) Standl.] Over Seasons under Salt Affected Soil. Int. j. plant soil sci. 2023;35(18)1887-1903.

© 2023 Yadav et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104396>