



Efficacy of Bio-Rational Insecticides in Controlling Thrips in Summer Country Bean

Most. Sukhy Begum ^a, M. M. Rahman ^a, M. S. Hossain ^a,
Sabera Yasmin ^a, Ruhul Amin ^a, Md. Shahidul Islam Khan ^a
and Md. Emam Hossain ^{a*}

^a Department of Entomology, Sher-e-Bangla Agricultural University, Dhaka-1207, Bangladesh.

Authors' contributions

This work was carried out in collaboration among all authors. Authors MMR and MSH planned and designed the research. Author MSB conducted the work on the field, collected the data and managed literature searches. Authors MSIK and MEH performed the statistical analysis and wrote the manuscript. Authors SY and RA edited and reviewed the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARRB/2023/v38i1030612

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111220>

Original Research Article

Received: 24/10/2023
Accepted: 29/12/2023
Published: 30/12/2023

ABSTRACT

Aims: To study the efficacy of biorational insecticides in controlling thrips in summer country bean.
Study design: The study used a single component Randomized Complete Block Design (RCBD) with three replications.
Place and Duration of Study: The experiment was conducted in the central farm of Sher-e-Bangla Agricultural University, Dhaka, Bangladesh from March to May 2021.
Methodology: The experiment consisted of seven treatments and was laid in a single factor Randomized Complete Block Design (RCBD) with three replications. The experiment consists of the following management practices: T₁=Spraying Neem oil @ 6 ml/L of water at 7 days interval,

*Corresponding author: E-mail: emamhossain72@gmail.com;

T₂=Spraying Neem leaf extract @ 200 mg/L of water at 7 days interval, T₃= Mehgoni leaf extract @ 200 mg/L of water at 7 days interval, T₄=Tobacco leaf powder @ 30 mg/L of water at 7 days interval, T₅= Spraying Spinosad @ 0.4 ml/L of water at 7 days interval, T₆=Spraying Biotrin @1.4 ml/L of water at 7 days interval and T₇= Untreated control.

Results: Among the seven (7) treatments the lowest number of thrips (2.50 plant⁻¹) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval), while highest number of thrips (16.67 plant⁻¹) was observed in T₇ (Untreated control). The highest percentage of thrips reduction over control (85.50%) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval) treatment. The present study revealed that the lowest number of thrips as insect pest and increased plant, leaf, inflorescence and pod infestation, different yield contributing characters and yield of country bean might be obtained by using the treatment T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval) besides T₆ (Spraying Biotrin @ 1.4 ml/L of water at 7 days interval) performed as second best treatment whereas, T₇ (Control) and T₃ (Mehgoni leaf extract @ 200 mg/L of water at 7 days interval) may protect the country bean at lowest rate in summer.

Conclusion: It can be concluded that, spraying of Spinosad @ 0.4ml/L of water at 7 days interval as a bio-rational management practice for thrips population control for country bean variety BARI Sheem-4 in summer reduced the infestation of thrips at the most satisfactory level.

Keywords: Biorational insecticides; thrips; country bean; leaf extract.

1. INTRODUCTION

Country bean (*Lablab purpureus*) is an important vegetable-cum-pulse, food-secure and nutritious crop and well known as "Sheem". It is reported to be originated in India [1]. then spread to other parts of the world. It is grown in a significant acreage after brinjal and tomato in Bangladesh. In Bangladesh, country bean has been cultivated in 51578 acres of land with a production of 144050 metric ton [2]. Country bean (*Lablab purpureus*) is normally grown during the rabi or winter season. Around 12,000 ha lands are cultivated and 50,000 metric tons of pods are produced every year. Country bean is rich in protein and can be grown easily on roof tops or trellises. Insect pests and diseases have caused a sharp decline in Bangladesh's country bean production, despite the crop's importance [3,4]. From the time of seedling to harvest, almost 55 different types of mites and insect pests target country beans [5]. Thrips, Aphids, bean bugs, green semiloopers, hooded hoppers, leaf miners, leaf weevils, pod borers, shot borers, shoot weevils, epilachna beetles, and mites are among the insect pests that seriously harm the nation's bean crop economically [6]. Thrips is one of the most important insect pests of country bean at flowering and harvesting stage [7]. Thrips damages the pods of beans. Infestation begins in the flowers and pods become twisted, deformed with reddish-brown russet marks as the thrips feed. Large infestation causes poor pod set, stunted plants and leaves and flowers to wilt [7]. Farmers in Bangladesh mostly use chemical pesticides due to their affordability, availability, and efficacy [8,9]. The ecology and the

biodiversity of the land is severely harmed due to indiscriminate use of pesticide without consideration of their negative consequences [10].

In most of the cases, the farmers either forgot the instructions or did not care to follow those instructions and went on using insecticides at their own choice or experience. Some farmers believed that excess use of insecticide could solve the insect pests' problem. As a result, harmful impact of insecticides on man, animal, wildlife, beneficial insects and environment is imposing a serious threat. Previously many researchers have also used and evaluated different synthetic chemicals against different insect pests, especially against sucking insects like thrips of country bean. These conventional chemical control measures failed to adequately control thrips which results in severe yield losses [11].

Products derived from the natural sources are known as bio-rational insecticide such as plant extracts, insect pathogens etc. are efficacious against target pest but safe to major natural enemies. As a result, bio-rational insecticides are becoming popular day-by-day considering the above situations present experiment was conducted to find a suitable way to safe production of country bean with profitable yield. For controlling the insect pest successfully and to save biological control agents, it is necessary to use environmentally safe insecticides judiciously. In this experiment, an effort was taken to find out the most effective bio-insecticide in controlling thrips of country bean, under the circumstances

of the necessity to find out some eco-friendly alternative methods for thrips management.

2. MATERIALS AND METHODS

The present research was conducted at the research field of Sher-e-Bangla Agricultural University, Sher-e-Bangla Nagar, Dhaka during the period from March to May 2021. Summer-country bean variety BARI Sheem-4 was used as experimental materials for the study and the seed of the variety of this experiment collected from Bangladesh Agricultural Research Institute (BARI), Joydebpur, Gazipur. The experiment comprised of seven treatments including an untreated control applied at 7 days interval. The details of the treatments are given below: T₁=Spraying Neem oil @ 3 ml/L of water, T₂=Spraying Neem leaf extract @ 200 mg/L of water, T₃= Mehgoni leaf extract @ 200 mg/L of water, T₄=Tobacco leaf powder @ 30 mg/L of water, T₅= Spraying Spinosad @ 0.4 ml/L of water T₆=Spraying Biotrin @1.4 ml/L of water T₇=Untreated control. The experiment was laid out in Randomized Complete Block Design (RCBD) with three replications using variety BARI Sheem-4. Each block was divided into seven equal plots. Thus, there were 21 (3 × 7) unit plots altogether in the experiment. The size of each unit plot was 3 m × 2 m. The distance between block to block and that of the plot to plot was 1.0 m and 1.0 m, respectively. The treatments of the experiment randomly allotted into the experimental plot. The fertilizers were applied as per fertilizers recommendation guide [12]. The applied manures were mixed properly with the soil in the plot using a spade. The dose and method of application of fertilizers are as

Urea 30 kg/ha. TSP 70 kg/ha. And MP 35 kg/ha. Seeds were sown on 6th March 2021 at the rate of 45 kg ha⁻¹ in the furrow and the furrows were covered with soils soon after seeding. The line to line (furrow to furrow) distance was maintained treatment arrangements with continuous sowing of seeds in the line. Plant to plant distance was 6 cm. Data were collected on different parameters viz. number of thrips incidence plant⁻¹, number of infested plant plot⁻¹, number of infested leaf plant⁻¹, number of infested inflorescence plant⁻¹, number of infested pod inflorescence⁻¹. The data obtained from experiment on various parameters were statistically analyzed in MSTAT-C computer program [13]. The mean values for all the parameters were calculated and the analysis of variance for the characters was accomplished by the Least Significant Differences (LSD) test at 5 % levels of probability [14].

3. RESULTS AND DISCUSSION

3.1 Incidence of Thrips in Summer Country Bean Under Different Bio-rational Insecticides

The average population of thrips in summer country bean under different bio-rational insecticides has been shown in Fig. 1. The data (Table 1) expressed that the lowest number of thrips (2.50 plant⁻¹) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval), while highest number of thrips (16.67 plant⁻¹) was observed in T₇ (Untreated control). The highest percentage of thrips reduction over control (85.50%) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval) treatment.




Fig. 1. Average population of thrips and efficiency of different bio-rational insecticides country bean in summer

Table 1. Efficiency of different bio-rational insecticides against leaf infestation in controlling Thrips of country bean in summer

Treatments	Number observed plants plot ⁻¹	Number healthy plants plot ⁻¹	Number infested plants plot ⁻¹
T ₁ (Neem oil @ 3 ml/L of water)	10	5.30 ab	4.67 ab
T ₂ (Neem leaf extract @ 200 mg/L of water)	10	5.10 ab	4.90 ab
T ₃ (Mehgoni leaf extract @ 200 mg/L of water)	10	4.50 bc	5.50 ab
T ₄ (Tobacco leaf powder @ 30 mg/L of water)	10	4.70 b	5.30 ab
T ₅ (Spinosad @ 0.4 ml/L of water)	10	6.00 a	4.00 bc
T ₆ (Biotrin @1.4 ml/L of water)	10	5.67 ab	3.33 c
T ₇ (Untreated control)	10	4.00 c	6.00 a
LSD (0.05)	-	1.29	1.08
Level of Sig.	-	*	**
CV %	-	8.41	11.05

[In a column, means followed by the same letter(s) are not significantly different at 5% level of probability by Least Significant Differences (LSD)]

Table 2. Efficiency of different bio-rational insecticides against inflorescence infestation in controlling Thrips of country bean in summer at different flowering stage

Treatments	Number infested inflorescence plant ⁻¹	Number infested inflorescence plant ⁻¹	Number infested inflorescence plant ⁻¹
	Early flowering stage	Mid flowering stage	Late flowering stage
T ₁ (Neem oil @ 3 ml/L of water)	6.67 de	9.67 d	5.66 cd
T ₂ (Neem leaf extract @ 200 mg/L of water)	7.00 d	10.33 cd	6.00 bc
T ₃ (Mehgoni leaf extract @ 200 mg/L of water)	9.33 b	12.10 b	7.33 b
T ₄ (Tobacco leaf powder @ 30 mg/L of water)	7.73 c	10.80 c	6.50 bc
T ₅ (Spinosad @ 0.4 ml/L of water)	5.00 ef	8.00 e	4.00 e
T ₆ (Biotrin @1.4 ml/L of water)	5.67 e	8.50 de	4.67 d
T ₇ (Untreated control)	11.75 a	14.33 a	10.75 a
LSD (0.05)	1.29	1.17	1.77
Level of Sig.	**	**	*
CV %	8.70	5.04	10.33

[In a column, means followed by the same letter(s) are not significantly different at 5% level of probability by Least Significant Differences (LSD)]

3.2 Number of Infested Plants Plot⁻¹

Plant infestation caused by thrips on summer country bean was significantly varied in different effect of bio-rational insecticides (Table 1). The result demonstrated that the lowest percent plant infestation (33.30%) was found in T₆ (Spraying Biotrin @1.4 ml/L of water at 7 days interval), and the highest percent plant infestation (60%) was found in T₇ (Untreated control). The lowest number of infested plants (3.33 plot⁻¹) was observed in T₆ (Spraying Biotrin @1.4 ml/L of water at 7 days interval), the highest number of infested plants (6 plot⁻¹) was observed in T₇ (Untreated control). The lowest number of healthy plants (4 plot⁻¹) was observed in T₇ (Untreated control), the highest number of healthy plants (6 plot⁻¹) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval). Similar result was also observed by [15].

3.3 Number of Infested Inflorescence Plant⁻¹ at Different Flowering Stages

Inflorescence infestation at early flowering stage caused by thrips on summer country bean was significantly varied in different effect of bio-rational insecticides (Table 2). The lowest number of infested inflorescence (5.00 plant⁻¹) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval), the highest number of infested inflorescence (11.75 plant⁻¹) was observed in T₇ (Untreated control). At mid flowering stage, the lowest number of infested inflorescence (8.00 plant⁻¹) was observed in T₅ (Spraying Spinosad @ 0.4ml/L of water at 7 days interval), the highest number of infested inflorescence (14.33 plant⁻¹) was observed in T₇ (Untreated control). The lowest number of infested inflorescence (4.00 plant⁻¹) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval), the highest number of infested inflorescence (10.75 plant⁻¹) was observed in T₇ (Untreated control). These results support the findings of Atakan [16].

3.4 Number of Infested Flower Inflorescence⁻¹ at Different Flowering Stages

Flower infestation at early flowering stage caused by thrips on summer country bean was significantly varied in different effect of bio-rational insecticides (Table 3). The lowest number of infested flower (2.00 inflorescence⁻¹)

was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval), the highest number of infested flower (5 inflorescence⁻¹) was observed in T₇ (Untreated control).

At mid flowering stage, the lowest number of infested flower (3.00 inflorescence⁻¹) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval), the highest number of infested flower (5 inflorescence⁻¹) was observed in T₇ (Untreated control) (Table 3). The result demonstrated that the lowest percent flower infestation (20%) was found in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval), the highest percent flower infestation (47.62%) was found in T₇ (Untreated control) (Table 3).

At later stage, the lowest number of infested flower (1.67 inflorescence⁻¹) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval), the highest number of infested flower (3.75 inflorescence⁻¹) was observed in T₃ (Mehgoni leaf extract @ 200 mg/L of water at 7 days interval) (Table 3). Botanical insecticides are naturally occurring chemicals extracted from plants. They are easily biodegradable, maintain biological diversity of predators and reduce environmental degradation and human health hazards [17]. Studies by Nderitu et al. [18]. Palumbo et al. [19]. Indicated that botanical pesticides are not as effective as synthetic pesticides, however neem-based pesticides can be used in an IPM program to neem-based use of synthetic insecticides and production costs.

3.5 Number of Infested Pod Inflorescence⁻¹ at Different Fruiting Stages

Pod infestation at early flowering stage caused by thrips on summer country bean was significantly varied in different effect of bio-rational insecticides (Table 4). The result demonstrated that the lowest percent pod infestation (23.30%) was found in T₅ (Spraying Spinosad @0.4ml/L of water at 7 days interval), the highest percent pod infestation (62.50%) was found in T₇ (Untreated control). The lowest number of infested pod (2.33 inflorescence⁻¹) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval), the highest number of infested pod (5 inflorescence⁻¹) was observed in T₇ (Untreated control). The lowest number of healthy pod (4 inflorescence⁻¹) was observed in T₇ (Untreated control), the highest number of healthy pod (7.67 inflorescence⁻¹) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L

Table 3. Efficiency of different bio-rational insecticides against flower infestation in controlling Thrips of country bean in summer at different flowering stage

Treatments	Number of infested flower inflorescence ⁻¹	Number of infested flower inflorescence ⁻¹	Number of infested flower inflorescence ⁻¹
	Early	Mid	Late
T ₁ (Neem oil @ 3 ml/L of water)	3.80 b	4.00 ab	2.67 bc
T ₂ (Neem leaf extract @ 200 mg/L of water)	4.33 ab	4.33 ab	3.33 ab
T ₃ (Mehgoni leaf extract @ 200 mg/L of water)	4.67 ab	4.75 ab	3.75 a
T ₄ (Tobacco leaf powder @ 30 mg/L of water)	4.33 ab	4.67 ab	3.33 ab
T ₅ (Spinosad @ 0.4 ml/L of water)	2.00 c	3.00 bc	1.67 c
T ₆ (Biotrin @1.4 ml/L of water)	2.67 bc	3.33 b	2.33 bc
T ₇ (Untreated control)	5.00 a	5.00 a	3.67 ab
LSD (0.05)	1.04	1.05	1.24
Level of Sig.	**	*	**
CV %	6.43	4.61	3.78

[In a column, means followed by the same letter(s) are not significantly different at 5% level of probability by Least Significant Differences (LSD)]

Table 4. Efficiency of different bio-rational insecticides against pod infestation in controlling Thrips of country bean in summer at different fruiting stages

Treatments	Number of infested pod inflorescence ⁻¹	Number of infested pod inflorescence ⁻¹	Number of infested pod inflorescence ⁻¹
	Early	Mid	Late
T ₁ (Neem oil @ 3 ml/L of water)	3.00 bc	3.67 ab	2.33 ab
T ₂ (Neem leaf extract @ 200 mg/L of water)	3.67 b	3.80 ab	2.67 ab
T ₃ (Mehgoni leaf extract @ 200 mg/L of water)	4.00 ab	3.33 ab	3.00 a
T ₄ (Tobacco leaf powder @ 30 mg/L of water)	3.80 ab	3.00 b	2.67 ab
T ₅ (Spinosad @ 0.4 ml/L of water)	2.33 c	2.33 bc	2.00 bc
T ₆ (Biotrin @1.4 ml/L of water)	2.67 bc	2.67 bc	2.50 ab
T ₇ (Untreated control)	5.00 a	5.33 a	3.00 a
LSD (0.05)	1.08	1.05	0.93
Level of Sig.	**	**	*
CV %	3.19	6.33	4.05

[In a column, means followed by the same letter(s) are not significantly different at 5% level of probability by Least Significant Differences (LSD)]




Fig. 2. Efficiency of different bio-rational insecticides against yield of country bean in controlling Thrips in summer

Pod infestation at mid flowering stage caused by thrips on summer country bean was significantly varied in different effect of bio-rational insecticides (Table 4). The result demonstrated that the lowest percent pod infestation (22.56%) was found in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval), the highest percent pod infestation (66.62%) was found in T₇ (Untreated control). The lowest number of infested pod (2.33 inflorescence⁻¹) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval), the highest number of infested pod (5.33 inflorescence⁻¹) was observed in T₇ (Untreated control). The lowest number of healthy pod (3.67 inflorescence⁻¹) was observed in T₇ (Untreated control), the highest number of healthy pod (8 inflorescence⁻¹) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval).

Pod infestation at late flowering stage caused by thrips on summer country bean was significantly varied in different effect of bio-rational insecticides (Table 4). The result demonstrated that the lowest percent pod infestation (21.05%) was found in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval), the highest percent pod infestation (50%) was found in T₇ (Untreated control). The lowest number of infested pod (2 inflorescence⁻¹) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval), the highest number of infested pod (3 inflorescence⁻¹) was observed in T₇ (Untreated

control) and T₃ (Mehgoni leaf extract @ 200 mg/L of water at 7 days interval). The lowest number of healthy pod (2 inflorescence⁻¹) was observed in T₇ (Untreated control), the highest number of healthy pod (7.5 inflorescence⁻¹) was observed in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval).

3.6 Pod Yield (Ton/ha.)

Infestation caused by thrips on summer country bean was significantly varied in different effect of bio-rational insecticides (Fig. 2). The result demonstrated that the lowest pod yield (7.25 t/ha) was found in T₃ (Mehgoni leaf extract @ 200 mg/L of water at 7 days interval), the highest pod yield (16 t/ha) was found in T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval). (Fig. 2)

4. CONCLUSION

The present study revealed that the lowest number of thrips as insect pest and increased plant, leaf, inflorescence and pod infestation, different yield contributing characters and yield of country bean might be obtained by using the treatment T₅ (Spraying Spinosad @ 0.4 ml/L of water at 7 days interval) besides T₆ (Spraying Biotrin @ 1.4 ml/L of water at 7 days interval) performed as second best treatment whereas, T₇ (Control) and T₃ (Mehgoni leaf extract @ 200 mg/L of water at 7 days interval)

may protect the country bean at lowest rate in summer. The highest (3.08) benefit cost ratio was obtained due to application of T₅(Spraying Spinosad @ 0.4ml/L of water at 7 days interval) while the lowest (2.26) in T₇(control). From the above description, it can be concluded that, spraying of Spinosad @ 0.4 ml/L of water at 7 days interval as a bio-rational management practice for thrips population control for country bean variety BARI Sheem-4 in summer reduced the infestation of Thrips at the most satisfactory level.

ACKNOWLEDGEMENTS

The National Science and Technology (NST) Fellowship, Ministry of Science and Technology, Government of the People's Republic of Bangladesh, supported this research. The authors are appreciative for the financial assistance they have received.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Sibiko KW, Ayuya OI, Gido EO, Mwangi JK. An analysis of economic efficiency in bean production: Acta entomology and zoology evidence from eastern Uganda. *J Econ Sust Dev.* 2013;4(13):1-10.
- BBS. Statistical Yearbook of Bangladesh. Bangladesh bureau of statistics. Statistics Division, ministry of planning, Govt. of Peoples Republic of Bangladesh. Dhaka. Bangladesh. 2020;71.
- Khan AU, Choudhury MAR, Islam MS, Maleque MA. Abundance and fluctuation patterns of insect pests in country bean. *J Sylhet Agric Univ.* 2018;5:167-172.
- Mollah MMI, Rahman MM, Khatun S, Mala M. Insect pest complex of year round country bean (*Lablab purpureus* L.) during summer season. *Scirea J Agric.* 2017;1:186-196.
- Chopkar PS, Desai VS, Samrit RM, Uparkar AL, Choudhari RJ, Shelke SB. Effect of border crops on pest population in Lablab bean (*Lablab purpureus* L.). *J Entomol Zool Stud.* 2020;8:1407-1412.
- Chowdhury MGF, Rahman MA, Miaruddin M, Khan MHH, Rahman MM. Assessment of pesticides and ripening chemicals used in selected vegetables at different locations of Bangladesh. *Bangladesh J Agric Res.* 2019;44:261-279.
- Singh BP, Singh B. Response of French bean to phosphorus and boron in cid alfisols in Meghalaya. *J Indian Society Soil Sci.* 2019;38(4):769-771.
- Pham TT, Van Geluwe S, Nguyen VA, Van der Bruggen B. Management challenges for sustainable use of pesticides for tropical crops in (South-East) Asia to avoid environmental pollution. *J Mater Cycles Waste Manag.* 2012;14:379-387.
- Siddique MA, Azad AK. Prioritization of Research for Horticultural Crops; Final Report of a Study; BARC: Dhaka, Bangladesh, 2010;56.
- Muriithi BW, Affognon HD, Diiro GM, Kingori SW, Tanga CM, Nderitu PW, Mohamed SA, Ekesi S. Impact assessment of Integrated Pest Management (IPM) strategy for suppression of mango-infesting fruit flies in Kenya. *Crop Prot.* 2016;81:20-29.
- Islam MA. Integrated pest (Insects) management of vegetables. Consultancy report, AVRDC - USAID Bangladesh Project, Horticulture Research Center, BARI, Gazipur. 1998;1999:1701.
- BARI (Bangladesh Agriculture Research Institute). *Krishi projukti hatboi* (in Bangla). 4th ed., Bangladesh Agril Res Inst, Gazipur, Bangladesh. 2006;209-211.
- Russell DF. MSTAT-C package programme. Dept. Crop Soil Sci. Michigan State Univ. USA. 1986;59-60.
- Gomez KA, Gomez AA. Statistical procedures for agricultural research. A Wiley Int. Sci. Publ. John Wiley and Sons. New York, Brisbane, Singapore. 1984;139-240
- Hossain MA. Development of insecticide application schedule for management of flower thrips and pod borer in mungbean (*Vigna radiata* L.). *Bangladesh J Agril Res.* 2013;38(1):19-28.
- Atakan E. Thrips (*Thysanoptera*) species occurring in winter vegetable crops in Çukurova region of Turkey. *Acta Phytopatho Entomol Hungarica.* 2008;43(2):227-234.
- Asogwa EU, Ndubuaku TCN, Ugwu JA, Awe OO. Prospects of botanical pesticides from neem, *Azadirachta indica* for routine protection of cocoa farms against the brown cocoa mirid – *Sahlbergella singularis*. *Nigeria J Medi Plants Rech.* 2010;4(1):001-006.

18. Nderitu JH, Wambua EM, Olubayo F, Kasina JM, Waturu CN. Management of thrips (*Thysanoptera: Thripidae*) Infestation on French beans (*Phaseolus vulgaris* L.) in Kenya by combination of insecticides and varietal Resistance. J Entomol. 2007; 4:469-473.
19. Palumbo J, Mullis C, Reyes F, Amaya A, Ledesma L, Cary L. Management of western flower thrips in head lettuce with conventional and botanical insecticides university of Arizona college of agriculture. Vegetable Report; 2000. Available:<http://ag.arizona.edu/pubs/crops/az1177>

© 2023 Begum et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/111220>