



Effect of Germination Time on the Properties of Flours from Red and Brown Varieties of Pigeon Pea (*Cajanus cajan*) Seeds

Eze, S. I.^a, Okoye, J. I.^{b*} and Onwuama, N. N.^c

^a Department of Home Economics, Federal College of Education, Eha-Amufu, Enugu State, Nigeria.

^b Department of Food Science and Technology, Enugu State University of Science and Technology, Enugu, Nigeria.

^c Department of Nutrition and Dietetics, University of Nigeria Teaching Hospital, Ituku-Ozalla, Enugu State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJBCRR/2024/v33i3864

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/108857>

Original Research Article

Received: 23/09/2023

Accepted: 27/11/2023

Published: 18/03/2024

ABSTRACT

This study was designed to evaluate the effect of germination time on the proximate composition, functional and pasting properties of flours from red and brown varieties of pigeon pea seeds. The red and brown pigeon pea seeds were cleaned and germinated for 48, 72 and 96 h, respectively and processed into flours, while the flours produced from the ungerminated red and brown pigeon pea seeds served as control samples. The flours obtained were analyzed for proximate, functional and pasting properties using standard methods. The proximate composition of the samples showed that the flours produced from the red pigeon pea seeds had a range of 6.24 – 9.34% moisture, 10.37 – 20.95% crude protein, 1.21 – 1.12% fat, 2.32 – 3.32% ash, 3.14 – 4.23% crude fibre, 76.75-

*Corresponding author: E-mail: okoyejoseph6@gmail.com;

61.07% carbohydrate and 329.29 – 338.10KJ/100g energy, while that of the brown pigeon pea flours had a range of 6.44-9.77% moisture, 10.47-23.69% crude protein, 1.23-1.13% fat, 2.42-3.13% ash, 3.18-4.31% crude fibre, 76.26-57.99% carbohydrate and 358.18-336.83KJ/100g energy. The functional properties of the red pigeon pea flours revealed that the bulk density, water absorption capacity, oil absorption capacity, swelling capacity, gelation capacity and foam capacity ranged from 0.58-0.22 ml/g, 1.36-1.58 %, 1.40-1.61 %, 2.13-2.38 ml/g, 8.24-11.13 ml/g and 12.28-12.01 %, while that of the brown variety ranged from 0.63-0.36 ml/g, 1.47-1.69 %, 1.46-1.67 %, 1.67-2.46 ml/g, 8.88-11.41 ml/g and 13.33-13.06 %, respectively. The pasting properties of the red pigeon pea flours also showed that the peak, trough, breakdown, set back and final viscosities as well as the peak time and pasting temperature ranged from 3.10 – 5.556cP, 3.35 – 4.86cP, 2.65 – 4.57cP, 5.23 – 7.75cP, 8.82 -10.02cP, 6.79 – 7.34min and 69.79 – 92.71°C, while that of the brown variety ranged from 3.59 – 6.21cP, 4.19 – 5.61cP, 3.77 – 5.23cP, 5.59 – 8.25cP, 9.60 – 11.08cP, 7.07 – 7.99min and 80.11 – 99.21°C, respectively. The study, however, revealed that the flours produced from both the red and brown pigeon pea seeds could be generally used as nutrient dense and functional ingredients in the preparation of a wide range of food products especially in Nigeria and other developing countries of the world where there is acute shortage of protein in order to ensure food and nutritional security.

Keywords: *Pigeon pea seeds; varieties; proximate composition; germination; functional properties; pasting properties.*

1. INTRODUCTION

Legumes are nutritious foods which serve as good substitutes for animal protein. The term legume is applied broadly to cover all plants of pea and bean family which are botanically referred to as *Leguminosae* or *Fabaceae* [1]. Legumes are known to be the third largest family among the flowering plants and they consist of approximately 650 genera and 20,000 species [2].

Pigeon pea is one of the legumes that is commonly referred to as 'otili' in the south-west of Nigeria, where the seeds are boiled and eaten by the rural populace. It is also called 'fio-fio' in the Igbo speaking areas of Nigeria. Although it is a perennial herbaceous leguminous plant, it is cultivated at subsistence level in Nigeria with the existence of different varieties. Pigeon pea is mainly classified and consumed as lesser known and underutilized legume in Nigeria and other sub-Sahara African countries compared to other legumes like cowpea, soybean and kidney bean despite its numerous nutritional benefits which include the improvement of the level of malnutrition and boosting of food security [3,4]. It can also serve as one of the good sources of plant protein, vitamins, minerals and plant sterols. It is one of the important pulses in India where it contributes about 20% to the total production of all pulses.

"Pigeon pea seeds are made up of 85% cotyledons, 14% seed coat and about 10%

embryo which contains a variety of dietary nutrients" [5]. "It also contains 20-22% protein, 1.2% fat, 65% carbohydrate and 3.8% ash" [6]. It equally contains crude fibre, magnesium, manganese and copper [5].

Germination refers to a series of changes in morphology, physiology and biochemistry. It is characterized by the growth of the embryo in the grain that is manifested by the growth of rootlets and increase in the length of the shoot (acropire), which would lead to the modification of the contents of the endosperm. These changes are likely to affect the nutrient and antinutrient contents as well as the functional properties of the flour produced after germination.

Germination is one of the most common and effective processes for improving the quality of legumes and germinated legumes are widely consumed all over the world. The process is influenced by the varietal differences and external factors such as soaking and germination time" [7]. During germination, some of the food reserve materials in legume seeds are degraded and used for respiration and synthesis of new cells in the developing embryo, thereby causing significant changes in nutritional, functional and sensory characteristics of the legume flours and other products. Steve [8] and Valdez Aquino et al. [9] also noted that "germination is a process that is not expensive but is technologically effective in improving the nutritional quality of beans".

Germination also increases the nutrient content, digestibility and availability of free amino acids, dietary fibre and bioactive components of lupine seeds (*Lapinus angustifolius* L). Germination equally increases the water absorption capacity of seeds as well as the protein solubility of minced bean. It also increases the capacity of millet flour to bind oil but decreases its water absorption capacity. In addition, germination has been reported to increase the protein and crude fibre contents of legume products.

Functional properties refer to the intrinsic physicochemical characteristics which may affect the behaviour of food systems during processing, storage and consumption such as solubility, foamability, gelation and emulsification properties [10].

Functional properties constitute the major criteria used for the adoption and acceptability of protein in food systems during processing, preservation and preparation. Functional properties include bulk density, protein solubility, gelation capacity, wettability, water and oil absorption capacities, emulsification and foaming properties among others. Although pigeon pea is known to be a rich source of many substances, the effect of germination on the nutrient composition, functional and pasting properties of red and brown pigeon pea flours have not been widely reported in recent times. For proper utilization and acceptability of the legume flour, it is desirable to study its rheological properties. This is because these properties contribute immensely in improving the shelf life as well as the sensory properties of foods during production, preparation and processing [11].

Pasting properties are among the rheological characteristics of starch that are used to determine the behaviour of starch in a food system especially when an aqueous suspension of starch is heated above a critical temperature during processing [3]. Some of the important pasting properties that are commonly determined during the processing of starchy and leguminous food products are peak, trough, setback, breakdown and final viscosities, in addition to other parameters which include peak time and pasting temperature. The complex nature of food composition often makes the measurement and interpretation of rheological properties difficult. To be able to successfully introduce a new supplement in any food item, it is necessary to know whether it possesses suitable functional

and pasting properties for food applications and consumer acceptability. The aim of this study therefore was to determine the effect of germination time on the proximate, functional and pasting profiles of the flours produced from red and brown varieties of pigeon pea seeds.

2. MATERIALS AND METHODS

2.1 Procurement of Raw Materials

Mature red and brown varieties of pigeon pea (*Cajanus cajan*) seeds used for this study were purchased from Ogbete Main Market, Enugu, Enugu State, Nigeria. The choice of location for the purchase was informed by the availability of fresh and wholesome red and brown varieties of pigeon pea seeds in the market.

2.2 Pre-preparation of the Seed Samples

The seeds were sorted manually to remove the stones, damaged and immature seeds after which the seeds were divided into four (4) equal lots of 500g each case. The first lot of each variety was processed raw, while the other lots were subjected to germination treatment at different steeping and holding periods, respectively.

2.3 Preparation of Raw Pigeon Pea Flour

The raw pigeon pea flour was prepared according to the method described by Ugwu and Oranye [12] with slight modifications. Five hundred grams (500g) of each variety of sorted pigeon pea seeds were sterilized with 1.5% Sodium hypochlorite solution followed by soaking in 75% ethanol for 20 min. The seeds were rinsed thoroughly with tap water, spread on trays and dried in a hot air oven (Model DHG 9101 ISA) at 60°C for 8 h with occasional stirring of the seeds at intervals of 30 min to ensure uniform drying. The dried seeds were dehulled by cracking them in the attrition mill followed by winnowing to remove the hulls. Thereafter, the dried and dehulled seeds were milled into flour using the attrition mill and sieved through a 500 micron mesh sieve. The flour produced was packaged in an airtight plastic container, labelled and kept in a refrigerator until needed for further use.

2.4 Preparation of Malted Pigeon Pea Flour Samples

The malted pigeon pea flours were prepared according to the method described by Ugwu and Oranye [12] with slight modifications. Five

hundred grams (500g) of each variety of sorted pigeon pea seeds were sterilized with 1.5% Sodium hypochlorite solution followed by soaking in 75% ethanol for 20 min. The seeds were rinsed thoroughly with tap water and steeped for 6 h in tap water at room temperature ($30\pm 2^{\circ}\text{C}$). The hydrated seeds were spread evenly on wet jute bags and allowed to germinate for 48 h. The seeds were sprinkled with water at intervals of 6 h to hasten sprouting. The ungerminated seeds were discarded. The sprouted seeds were collected and then dried in a hot air oven (Model DHG 9101 ISA) at 60°C for 6 h. The dried seeds were dehulled manually by rubbing them in-between palms to remove the hulls along with the rootlets. The dehulled seeds were milled into flour using the attrition mill and sieved through a 500 micron mesh sieve. The flour produced was packaged in an airtight plastic container, labelled and kept in a refrigerator until needed for further use. The same procedure was followed in the preparation of other samples of malted pigeon flours in each case except that steeping and germination treatments were carried out at intervals of 8 h for 72 h and 10 h for 96 h, respectively for both the third and fourth lots of red and brown varieties of pigeon pea seeds.

2.5 Proximate Analysis

The moisture, crude protein, ash, fat and crude fibre contents of the samples were determined on dry weight basis according to the standard analytical methods of AOAC [13]. The carbohydrate was calculated by difference. % Carbohydrate = $100 - (\% \text{ Moisture} + \% \text{ Crude Protein} + \% \text{ Fat} + \% \text{ Ash} + \% \text{ Crude Fibre})$. The energy content was calculated by multiplying the percentage values of protein, fat and carbohydrate by the Atwater factors of 4, 9 and 4, respectively [13]. All determinations were carried out in triplicate samples.

2.6 Evaluation of Functional Properties

The water and oil absorption capacities were determined on dry weight basis according to the methods of AOAC [13]. The foam and swelling capacities were determined according to the methods described by Onwuka [14]. The bulk density was determined according to the method described by Iwe et al. [15]. The gelation capacity was determined according to the method described by Butt and Batool [16]. All determinations were carried out in triplicate samples.

2.7 Evaluation of Pasting Properties

The pasting characteristics of the samples were determined using Rapid Visco Analyser (RVA) (Model 4500, Perten Instrument, Sweden) according to the method of AOAC [13]. Two grams (2g) of each sample was weighed into a dried empty canister and then 25mL of distilled water was dispensed into the canister containing the sample. The slurry was thoroughly mixed in each case and each canister was fitted into the Rapid Visco Analyzer. The slurry was individually heated from 50°C to 95°C with a holding time of 2 min followed by cooling to 50°C with 2 min holding time. The rate of heating and cooling was maintained at a constant rate of 11.25°C per min. The peak, trough, breakdown, final and setback viscosities as well as the peak time and pasting temperature of each sample were read from the pasting profile with the aid of a thermocytline for windows software that was connected to a computer.

2.8 Statistical Analysis

The data generated were subjected to one-way analysis of variance (ANOVA) using Statistical Package for Social Sciences (SPSS, Version 20) software. Significant means were separated using Turkey's least significant difference (LSD) test at $p < 0.05$.

3. RESULTS AND DISCUSSION

3.1 Proximate Composition of Raw and Germinated Red and Brown Pigeon Pea Flours

The proximate composition of raw and germinated red and brown pigeon pea flour samples are presented in Table 1.

The moisture content of the samples ranged from 6.24 to 9.34% for the red variety and 6.44 to 9.77% for the brown variety, while that of the raw samples ranged from 6.24 to 6.44% with the red raw flour sample having the least moisture content (6.24%). The samples steeped for 10 h and germinated for 96 h had the highest moisture contents of 9.34% and 9.77%, respectively for the red and brown pigeon pea flours. The increase could be attributed to the absorption of large quantity of water by the seeds due to prolonged soaking in water during processing. The moisture contents (6.24-9.34% and 6.44-9.77%) obtained for both samples were lower than the recommended moisture content limit of 14% that is compatible for the proper storage of legume flours [17]. "High moisture content of

Table 1. Proximate composition (%) of raw and germinated red and brown pigeon pea flours

Parameters	Samples							
	A 1 h (0 h)	B 6 h (48 h)	C 8 h (72 h)	D 10 h (96 h)	E 1 h (0 h)	F 6 h (48 h)	G 8 h (72 h)	H 10 h (96 h)
Moisture	6.24 ^g ±0.01	7.24 ^g ±0.01	8.11 ^d ±0.01	9.34 ^b ±0.01	6.44 ^f ±0.01	7.32 ^e ±0.01	8.55 ^c ±0.01	9.77 ^a ±0.01
Crude Protein	10.37 ^g ±0.01	14.51 ^f ±0.01	17.64 ^c ±0.01	20.95 ^b ±0.05	10.47 ^e ±0.01	15.10 ^d ±0.01	17.77 ^c ±0.01	23.69 ^a ±0.01
Fat	1.21 ^b ±0.01	1.18 ^c ±0.01	1.15 ^c ±0.00	1.12 ^d ±0.01	1.23 ^a ±0.01	1.20 ^b ±0.00	1.17 ^c ±0.01	1.13 ^d ±0.01
Ash	2.32 ^f ±0.02	2.41 ^e ±0.01	2.77 ^c ±0.01	3.32 ^a ±0.01	2.42 ^e ±0.01	2.56 ^d ±0.01	2.72 ^c ±0.02	3.13 ^b ±0.01
Crude fibre	3.14 ^g ±0.02	3.53 ^f ±0.01	3.88 ^d ±0.01	4.23 ^b ±0.01	3.18 ^g ±0.01	3.64 ^e ±0.01	3.95 ^c ±0.01	4.31 ^a ±0.01
Carbohydrate	76.75 ^a ±0.05	71.16 ^b ±0.02	66.47 ^d ±0.01	61.07 ^f ±0.01	76.26 ^b ±0.02	70.20 ^c ±0.02	65.87 ^e ±0.02	57.99 ^g ±0.06
Energy (KJ/100g)	359.29 ^g ±0.01	353.22 ^b ±0.12	346.76 ^d ±0.01	338.10 ^d ±0.09	358.18 ^a ±0.05	351.96 ^c ±0.11	345.005 ^e ±0.12	336.83 ^f ±0.07

Values are mean ± SD of triplicate determinations. Means within the same row with the same superscripts are not significantly different ($p < 0.05$)

A-Red (raw) pigeon pea flour without steeping and germination treatment, B-Red pigeon pea flour steeped for 6 h and germinated for 48 h, C-Red pigeon pea flour steeped for 8 h and germinated for 72 h, D-Red pigeon pea flour steeped for 10 h and germinated for 96 h, E- Brown (raw) pigeon pea flour without steeping and germination treatment, F- Brown pigeon pea flour steeped for 6 h and germinated for 48 h, G-Brown pigeon pea flour steeped for 8 h and germinated for 72 h, H-Brown pigeon pea flour steeped for 10 h and germinated for 96 h

flour samples have been reported to accelerate enzymatic and microbial activities which in turn lead to loss of nutrients and inferior sensory characteristics of such products” [18]. Therefore, the lower the moisture content of a food material, the higher its shelf stability. Nsa and Ukachukwu [19] reported that high moisture content generally affects the storage stability of legume and other flour products. The result obtained in this study indicate that the processed pigeon pea flours would have good keeping qualities with proper packaging and storage.

The crude protein content of the flour samples was significantly ($p < 0.05$) higher in the germinated pigeon pea flours and ranged from 10.37 to 20.95% for the red variety, while that of the brown variety ranged from 10.47 to 23.69% compared to the raw samples which ranged from 10.37 to 10.47%. This development is in consonance with the findings of Duenas et al. [20] who reported that germination has the ability to increase the protein content and availability of free amino acids of lupine seeds. Hamad and Field [21] also reported significant improvement in the protein content of sorghum and other grains during malting. “Dietary proteins are needed for the synthesis of new cells, enzymes and hormones required for the development of the body. Specific proteins help in blood clotting, fluid balance as well as in hormone and enzyme production [22].

The fat content of the flours ranged from 1.21 to 1.12% for the red variety and 1.23 to 1.13% for the brown variety. The fat content of the samples were generally reduced by steeping and germination treatments compared to the raw samples. The decrease in fat content could be due to increase in the activities of lipase enzymes which were activated during malting [23]. The decrease in fat contents of both the red and brown germinated pigeon pea flours could be nutritionally beneficial particularly in the preparation of a variety of food products for people suffering from obesity and hypercholesterolemia. Fat is important in human diets because it is a high energy-yielding nutrient. Fat provides body backup fuel and acts as an insulator to the body during cold. It also provides satiety, delicious flavours and aroma during food preparation and processing.

The ash content of the flour samples ranged from 2.32 to 3.32% for the red variety, while that of the brown variety ranged from 2.42 to 3.13%. The ash contents of the flours processed from both

varieties increased significantly ($p < 0.05$) by steeping and germination treatments. Mbaeyi and Onweluzo [24] reported a decrease in ash content of sprouted sorghum flour and suggested that the reduction could be due to the removal of vegetative part of the seeds during drying and milling which led to decrease in ash content.

The crude fibre content of the samples ranged from 3.14 to 4.31% with the brown variety steeped for 10 h and germinated for 96 h having the highest value (4.31%). The crude fibre content of the germinated pigeon pea flours increased with the increase in the period of steeping and germination treatments. This increase could be attributed to the extraction of more cell wall materials from the outer layers of the seeds during germination which in turn support the development of fibrous materials in the seeds [25]. Fibre plays a significant role in the digestion and absorption of foods through the promotion of increased excretion of bile acids, sterols and fats which have been implicated in the etiology of certain ailments in humans [22].

The carbohydrate content of the samples was significantly ($p < 0.05$) lower in the germinated flour samples compared to the control samples. The carbohydrate content ranged from 76.75 to 61.07% for the red variety, while that of the brown variety ranged from 57.99 to 76.26%. Similar decrease in carbohydrate content has been reported by Akubor and Chukwu [23] for germinated millet flour. The decrease in carbohydrate content could be due to the reduction in alpha amylase activity which tends to break down complex carbohydrates into simpler and more absorbable sugars by the growing seedlings during the early stage of germination. The decrease in the carbohydrate contents of both the red and brown germinated pigeon pea flours has some beneficial effects especially when the flours are intended for use in the formulation of food products for people suffering from diabetes mellitus and cardiovascular diseases.

The energy content of the samples ranged from 329.29 to 353.22KJ/100g for the red variety and 336.83 to 358.18KJ/100g for the brown variety. The raw flour sample produced from the brown variety had the highest energy content of 358.18KJ/100g compared to the red variety which had the energy value of 329.29KJ/100g. The result showed that the energy content of the samples was generally higher than the energy value of 325.46 KJ/100g reported by Adebowale

et al. (2011) for fermented pigeon pea flour. Generally, the steeping and germination of red and brown varieties of pigeon pea seeds for 10 and 96 h, respectively greatly improved the protein, ash and crude fibre contents of the flour samples, while their fat, carbohydrate and energy contents were drastically reduced.

3.2 Functional Properties of Raw and Germinated Red and Brown Pigeon Pea Flours

The functional properties of raw and germinated pigeon pea flour samples are presented in Table 2.

The bulk density of the processed samples ranged between 0.22 to 0.48 ml/g for the red variety and 0.36 to 0.58ml/g for the brown variety, while that of the raw samples ranged from 0.58 to 0.63 ml/g with the brown variety having the highest value (0.63 ml/g). The reduction in the bulk density observed in all the germinated samples could be due to solubilization of starch polymers during processing as a result of steeping and germination treatments [26]. The observation is in agreement with the report of Arukwe et al. [3] who stated that germination resulted in the reduction of the bulk density of the flours and other flour products. The bulk density is important in the selection of packaging materials and proper handling of food materials during processing and preparation of foods [27].

The water absorption capacity of the processed samples ranged between 1.44 to 1.58% for the red variety and 1.54 to 1.69% for the brown variety, whereas the water absorption capacity of the raw samples ranged from 1.36 to 1.47% with the brown variety having the highest value (1.47%). The increase in the water absorption capacity observed in all the germinated samples could be due to ability of the protein present in both varieties of the seed to absorb and retain water during processing as a result of steeping and germination treatments. The water absorption capacity of the samples was significantly ($p < 0.05$) higher in samples steeped for 10 h and germinated for 96 h compared to the samples steeped for 6 h and germinated for 48 h and the ones steeped for 8 h and germinated for 72 h, respectively for both the flours produced from the red and brown varieties of pigeon pea seeds. The observation is in agreement with the findings of Leonel et al. [28] who reported that germination increased the water absorption

capacity of the cereal and legume flours. The increase in water absorption capacity of both the processed red and brown pigeon pea flours suggest that they can be used as thickeners in the preparation of liquid and semi liquid foods because of their ability to absorb and retain water due to increase in hydrophilicity of the seed proteins [29,30]. The flours will also have higher flavour retention capacity and better mouthfeel than the flours made from the raw samples.

The oil absorption capacity of the processed samples ranged between 1.48 to 1.61% for the red variety and 1.46 to 1.67% for the brown variety, while that of the raw samples ranged from 1.40 to 1.46% for brown and red varieties. The increase in oil absorption capacity of the germinated samples could be attributed to the presence of more hydrophobic proteins than the non – polar amino acid side chains which have superior binding effects to oil during processing as a result of steeping and germination treatments [31]. The oil absorption capacity was significantly ($p < 0.05$) higher in samples steeped for 10 h and germinated for 96 h compared to the samples steeped for 6 h and germinated for 48 h and the ones steeped for 8 h and germinated for 72 h, respectively for both the red and brown pigeon pea flours. The result is in agreement with the report of Falmata et al. [32] who reported significant increase in oil absorption capacity of cereal and legume flours as a result of sprouting. The oil absorption capacity of the flour is known to be of great importance in enhancing the sensory characteristics such as flavour and mouthfeel of the foods. Flours with excellent oil absorption capacity may be useful in the preparation of pastries and doughnuts. The flours could be used also as flavour retainers in food formulations.

The swelling capacity of the processed samples ranged between 2.22 to 2.38ml/g for the red variety and 2.21 to 2.46ml/g for the brown variety, while that of raw samples ranged from 1.67 – 2.13ml/g with the brown variety having the highest value (2.13ml/g). The increase in the swelling capacity of the germinated pigeon pea flours might be due to modifications of the starch granules during sprouting which resulted in higher water uptake by the granules. The result is in agreement with the report of Falmata et al. [32] who stated that sprouting of legumes and cereals resulted in increase in the swelling capacities of their flours. The high swelling capacity of the flour is an indication of their suitability for use as disintegrants in the

pharmaceutical industry [33]. Flours with good swelling capacity are primarily used for thickening of soups, sauces, and gravies.

The gelation capacity of the processed samples ranged from 9.12 to 11.13ml/g for the red variety and 9.37 to 11.41ml/g for the brown variety, while that of the raw samples ranged from 8.24 – 8.88ml/g with the red and brown pigeon pea flours having the least (8.24ml/g) and highest (8.88ml/g) values, respectively. The variation in gelation capacity of germinated pigeon pea flours could be associated with the relative ratios of different constituents such as protein, carbohydrate and fat that made up the seeds [25]. Flours with good gelation capacity could be useful in the preparation of puddings and snacks especially where thickening and gelling are needed.

The foam capacity of the processed samples ranged between 12.01 to 12.19% for the red variety and 13.06 to 13.25% for the brown variety, whereas the foam capacity of the raw samples ranged from 12.28 to 13.33% with the brown variety having the highest value (13.33%). The foam capacity of the flour samples was found to decrease as the time of steeping and germination treatments increased during processing. This development is in consonance with the findings of Akubor and Obiegbunam [34] who reported that germination reduced the foam capacity of millet flour but its foam stability was slightly increased. Foams are used to improve the texture, consistency and appearance of foods during processing and preparation. Flours with good foam capacity could be used in the preparation of baked products and pastries.

Generally, steeping and germination greatly enhanced the functional properties of the pigeon pea flours with the exception of bulk density and foam capacity. This implies that the flours can be used in food systems like bread, cake and biscuit products. The ability of the flours to absorb water and oil may help to enhance the sensory qualities of bakery products and pastries such as flavour and mouthfeel.

3.3 Pasting Properties of Raw and Germinated Red and Brown Pigeon Pea Flours

The pasting properties of raw and germinated pigeon pea flour samples are presented in Table 3. The peak viscosity of the processed samples ranged between 4.10 to 5.56cP for the red

variety and 4.73 to 6.21cP for the brown variety, while the peak viscosity of the raw samples ranged from 3.10 to 3.59cP with the brown variety having the highest value (3.59cP). The samples steeped for 10 h and germinated for 96 h had the highest values of 5.56cP and 6.21cP, respectively for the red and brown pigeon pea flours. There were significant differences ($p<0.05$) in the peak viscosity of the flour samples. The increase in the peak viscosity of the germinated samples could be attributed to the increase in the activities of hydrolytic enzymes which degraded the starch molecules into simpler sugars during processing as a result of steeping and germination treatments [3]. Germination has been reported to increase the peak viscosity of food materials [26].

The trough viscosity of the processed samples ranged between 3.35 to 4.10cP for the red variety and 4.19 to 5.09cP for the brown variety, whereas that of the raw samples ranged from 4.86 to 5.61cP with the red pigeon pea flour having the least value (4.86cP). There were significant differences ($p<0.05$) in the trough viscosity of the flour samples. The reduction in the trough viscosity of the germinated samples could be attributed to the production of flours with low retrogradation tendencies from red and brown pigeon pea seeds coupled with the increase in the activities of hydrolytic enzymes as a result of steeping and germination treatments during processing [26]. Trough is the minimum viscosity which measures the ability of paste to withstand breakdown during cooling [35].

The breakdown viscosity of the processed samples ranged between 3.28 to 4.57cP for the red variety and 4.20 to 5.23cP for the brown variety, while the breakdown viscosity of the raw samples ranged from 2.65 to 3.77cP with the red pigeon pea flour having the least value (2.67cP). The samples steeped for 10 h and germinated for 96 h had the highest values of 4.57cP and 5.23cP, respectively for the red and brown pigeon pea flours. There were significant differences ($p<0.05$) in the breakdown viscosities of the flour samples. The increase in the breakdown viscosity observed in the germinated samples could be attributed to treatment effects which resulted in the degradation of starch molecules into simpler sugars by the activities of enzymes during processing. The breakdown viscosity is essentially a measure of the degree of paste stability or starch granule disintegration during heating [36]. The ability of starch to withstand heating at high temperature and shear

Table 2. Functional properties of raw and germinated red and brown pigeon pea flours

Parameters	A 1 h (0 h)	B 6 h (48 h)	C 8 h (72 h)	D 10 h (96 h)	E 1 h (0 h)	F 6 h (48 h)	G 8 h (72 h)	H 10 h (96 h)
Bulk Density (ml/g)	0.58 ^b ±0.01	0.48 ^c ±0.01	0.33 ^e ±0.01	0.22 ^f ±0.01	0.63 ^a ±0.01	0.58 ^b ±0.01	0.43 ^d ±0.01	0.36 ^e ±0.01
Water Absorption Capacity (%)	1.36 ^d ±0.01	1.44 ^c ±0.01	1.51 ^b ±0.01	1.58 ^a ±0.01	1.47 ^d ±0.01	1.54 ^c ±0.01	1.61 ^b ±0.01	1.69 ^a ±0.01
Oil Absorption capacity (%)	1.40 ^d ±0.01	1.48 ^c ±0.01	1.55 ^b ±0.01	1.61 ^a ±0.01	1.46 ^d ±0.01	1.54 ^c ±0.01	1.61 ^b ±0.01	1.67 ^a ±0.01
Swelling capacity (ml/g)	2.13 ^d ±0.01	2.22 ^c ±0.01	2.29 ^b ±0.01	2.38 ^a ±0.01	1.67 ^d ±0.05	2.21 ^c ±0.01	2.38 ^b ±0.01	2.46 ^a ±0.01
Gelation capacity (ml/g)	8.24 ^d ±0.02	9.12 ^c ±0.02	10.07 ^b ±0.02	11.13 ^a ±0.01	8.88 ^d ±0.01	9.37 ^c ±0.02	10.77 ^b ±0.01	11.41 ^a ±0.01
Foam capacity (%)	12.28 ^a ±0.01	12.19 ^b ±0.01	12.08 ^c ±0.01	12.01 ^d ±0.01	13.33 ^a ±0.01	13.25 ^b ±0.01	13.18 ^c ±0.01	13.06 ^d ±0.01

Values are mean ± SD of triplicate determinations. Means within the same row with the same superscripts were not significantly different ($p < 0.05$)

A-Red (raw) pigeon pea flour without steeping and germination treatment, B-Red pigeon pea flour steeped for 6 h and germinated for 48 h, C-Red pigeon pea flour steeped for 8 h and germinated for 72 h, D-Red pigeon pea flour steeped for 10 h and germinated for 96 h, E- Brown (raw) pigeon pea flour without steeping and germination treatment, F- Brown pigeon pea flour steeped for 6 h and germinated for 48 h, G-Brown pigeon pea flour steeped for 8 h and germinated for 72 h, H-Brown pigeon pea flour steeped for 10 h and germinated for 96 h

Table 3. Pasting properties of raw and germinated red and brown pigeon pea flours

Parameters	Samples							
	A 1 h (0 h)	B 6 h (48 h)	C 8 h (72 h)	D 10 h (96 h)	E 1 h (0 h)	F 6 h (48 h)	G 8 h (72 h)	H 10 h (96 h)
Peak viscosity (cP)	3.10 ^h ±0.38	4.01 ^f ±0.69	4.57 ^d ±0.72	5.56 ^b ±0.01	3.59 ^g ±0.70	4.73 ^e ±0.02	5.22 ^c ±0.01	6.21 ^a ±0.01
Trough viscosity (cP)	4.86 ^c ±0.01	4.10 ^e ±0.01	3.90 ^g ±0.70	3.35 ^h ±0.01	5.61 ^a ±0.01	5.09 ^b ±0.47	4.69 ^d ±0.49	4.19 ^f ±0.01
Breakdown viscosity (cP)	2.65 ^h ±0.66	3.28 ^g ±0.70	3.99 ^e ±0.01	4.57 ^c ±0.00	3.77 ^f ±0.01	4.20 ^d ±0.01	4.77 ^b ±0.47	5.23 ^a ±0.47
Setback Viscosity (cP)	7.75 ^b ±0.36	7.10 ^d ±0.93	6.10 ^f ±0.39	5.23 ^h ±0.60	8.25 ^a ±0.01	7.28 ^c ±0.62	6.57 ^e ±0.01	5.59 ^g ±0.24
Final viscosity (cP)	10.02 ^c ±0.01	9.77 ^e ±0.52	9.40 ^g ±0.57	8.82 ^h ±0.57	11.08 ^a ±0.57	10.42 ^b ±0.55	9.45 ^d ±0.61	9.60 ^f ±0.59
Peak time (min)	7.34 ^c ±0.01	7.21 ^d ±0.01	7.06 ^e ±0.01	6.79 ^f ±0.01	7.99 ^a ±0.01	7.76 ^b ±0.01	7.37 ^c ±0.01	7.07 ^e ±0.01
Pasting temperature (°C)	92.71 ^c ±0.01	88.33 ^d ±0.63	77.59 ^g ±0.01	69.79 ^h ±0.01	99.21 ^a ±0.41	93.66 ^b ±0.01	86.93 ^e ±0.53	80.11 ^f ±0.01

Values are mean ± SD of triplicate determinations. Means within the same row with the same superscripts were not significantly different ($p < 0.05$)

A-Red (raw) pigeon pea flour without steeping and germination treatment, B-Red pigeon pea flour steeped for 6 h and germinated for 48 h, C-Red pigeon pea flour steeped for 8 h and germinated for 72 h, D-Red pigeon pea flour steeped for 10 h and germinated for 96 h, E- Brown (raw) pigeon pea flour without steeping and germination treatment, F- Brown pigeon pea flour steeped for 6 h and germinated for 48 h, G-Brown pigeon pea flour steeped for 8 h and germinated for 72 h, H-Brown pigeon pea flour steeped for 10 h and germinated for 96 h.

stress is an important factor in many processes. Elofsson et al. [37] stated that the formation of gels by the proteins in foods results from a two-step process involving, first, the partial denaturation of individual proteins to allow more access to the reactive side groups within the protein molecules and secondly, the aggregation of these proteins by means of reactive side groups into a continuous three dimensional network structure that is capable of retaining significant amount of water and also exhibiting same structural rigidity. This phenomenon is of great importance in foods since it contributes significantly ($p < 0.05$) to the textural and rheological properties of various food products.

The setback viscosity of the processed samples ranged between 5.23 to 7.10cP for the red variety and 5.59 to 7.28cP for the brown variety, whereas that of the raw samples ranged from 7.75 to 8.25cP with the brown variety having the highest value (8.25cP). There were significant differences ($p < 0.05$) in the setback viscosity of the flour samples. The reduction in the setback viscosity of the germinated samples could be attributed to the increase in the activities of hydrolytic enzymes which degraded the starch molecules into simpler sugars as a result of steeping and germination treatments during processing [3]. The reduction in setback viscosity of the flour samples is an indication of low rate of retrogradation and syneresis tendency of the pastes prepared from the flours. Setback viscosity has been correlated with the texture of the food products [38].

The final viscosity of the processed samples ranged between 8.82 to 9.77cP for the red variety and 9.60 to 10.42cP for the brown variety, while the final viscosity of the raw samples ranged from 10.02 to 11.08cP for the red and brown pigeon pea flours, respectively. There were significant differences ($p < 0.05$) in the final viscosity of the flour samples. The reduction in the final viscosity of the germinated samples could be attributed to treatment effects during processing. The final viscosity is an indication of the stability of paste after cooking. Akubor [1] stated that flours with low final viscosity values are more stable than flours with high final viscosity values. Therefore, the low final viscosity values recorded by the germinated pigeon pea flours could be an added advantage in the preparation of a wide range of food products.

The peak time of the processed samples ranged between 6.79 to 7.21 min for the red variety and 7.07 to 7.76 min for the brown variety, whereas that of the raw samples ranged from 7.34 to 7.99 min with the brown variety having the highest value (7.99 min). There were significant differences ($p < 0.05$) in the peak time of the flour samples. The reduction in the peak time of the germinated samples could be attributed to the increase in the activities of hydrolytic enzymes which degraded the starch molecules of red and brown pigeon pea seeds into simpler sugars during processing as a result of steeping and germination treatments [3]. This showed that the samples steeped for 10 h and germinated for 96 h would require less time to form paste compared to the other samples steeped for 6 and 8 h and germinated for 48 and 72 h, respectively for the red and brown pigeon pea flours. The peak time is an indication of the actual time taken by a food product to attain its peak viscosity, and the lower the peak time, the faster the cooking time of such a food product [36].

The pasting temperature of the processed samples ranged between 69.79 to 88.33°C for the red variety and 80.11 to 93.66°C for the brown variety, while the pasting temperature of the raw samples ranged from 92.71 to 99.21°C with the red variety having the least value (92.71°C). There were significant differences ($p < 0.05$) in the pasting temperature of the flour samples. The differences in the pasting temperature of the processed samples could be due to treatment effects which might have made the samples to exhibit different stages of gelatinization [39]. The pasting temperature provides an indication of the minimum temperature required to cook a given sample and this also has some implications on energy usage during cooking [40]. This means that the germinated pigeon pea flour samples could cook faster and save more energy (fuel) than the raw samples during cooking.

Generally, the good pasting properties of the processed flours would be beneficial in the formulation of infant foods and in meat processing industry.

4. CONCLUSION

This present study evaluated the effect of germination on the proximate composition, functional and pasting properties of red and brown pigeon pea flours. The proximate composition of the samples revealed that

steeping and germination treatments relatively increased the protein, crude dietary fibre and ash contents of red and brown pigeon pea flours, while their carbohydrate, fat and energy contents decreased drastically. The functional properties of the samples showed that the water absorption, oil absorption, swelling and gelation capacities were enhanced by steeping and germination treatments during processing, while the bulk density and foam capacity of the samples decreased. The pasting properties revealed that the peak, breakdown and final viscosities increased significantly ($p < 0.05$) with increase in the time of steeping and germination during processing, while the trough and setback viscosities as well as the peak time and pasting temperature of the germinated flour samples decreased. Generally, the processed red and brown pigeon pea flours have good functional and pasting properties which suggest that they could be used in the preparation of baked products, supplementary foods and meat extenders. They could be also used as thickening agents in both liquid and semi-liquid food products.

ACKNOWLEDGEMENTS

The facilities provided by the Centre for Research and Innovation, National Root Crops Research Institute, Umudike Abia State, Nigeria to perform much of this work is appreciated. The authors would also like to thank the Management of Global International Company Ltd, Port Harcourt, Rivers State, Nigeria for financing this research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Akubor PI. Effect of processing methods on the chemical composition and functional properties of pigeon pea seed. *Asian Journal of Advances in Agricultural Research*. 2017;2(2):1-8.
2. Adebowale OJ, Maliki K. Effect of fermentation period on the chemical composition and functional properties of pigeon pea (*Cajanus cajan*) seed flour. *International Food Research Journal*. 2011;18(4):1329-1333.
3. Arukwe DC, Nwanekezi EC, Agomuo JK. Effects of combined processing methods on the functional and pasting properties of pigeon pea (*Cajanus cajan*) flour. *International Journal of Science and Qualitative Analysis*. 2017;3(3):23-30.
4. Nwanekezi EC, Ehirim FN, Arukwu DC. Combined effects of different processing methods on vitamin and anti-nutrient contents of pigeon pea (*Cajanus cajan*) flour. *IOSR Journal of Environment, Toxicology and Food Technology*. 2017; 11:73-81.
5. Faris DG, Singh U. Pigeon Pea: Nutrition and Products. In: The Pigeon pea (Nene Y.L; Hall S.D and Sheila V.K. Eds). CAB International, UK. 1990;401-434.
6. FAO. State of food and agriculture. Rome, Italy: Food and agriculture organization in developing countries. 2010;200-205.
7. Ridge I. The Regulation of Plant Growth. In: Ridge, I. (Ed.), *Plant Physiology*. Hodder and Stoughton, London, U.K. 1991;282-333.
8. Steve IO. Influence of germination and fermentation on chemical composition, protein quality and physical properties of wheat flour (*Triticum aestivum*). *Journal of Cereals and Oil Seeds*. 2012;3(3): 35- 47.
9. Valdez-Anguino DM, Herrera-Cabrera E, Davila-Oritz G, Oomah BD, Cardador-Martinez A, Jimenez-Martinez C. Content and distribution of nutritional and non-nutritional compounds during germination of three Mexican faba bean (*Vicia faba*). *International Journal of Research in Agriculture and Food Sciences*. 2015;2(9): 2311-2476.
10. Giami SY, Bekebain DA. Proximate composition and functional properties of raw and processed full-fat fluted pumpkin (*Telfairia occidentalis*) seed flour. *Journal of Food Science and Agriculture*. 1992; 59(3):321-325.
11. Adebowale KO, Lawal OS. Comparative study of the functional properties of bambara groundnut (*Voanbzeia subterranean*), jack bean (*Cavanalia ensiformis*) and mucuna bean (*Mucuna pruriens*). *Food Research International*. 2004;37:355-365.
12. Ugwu FM, Oranye NA. Effects of some processing methods on the toxic components of African breadfruit (*Treculia africana*). *African Journal of Biotechnology*. 2006;6:234-246.
13. AOAC. Official methods of analysis. 18th edn. Association of official analytical

- chemists. Washington D.C. USA. 2010; 214-225.
14. Onwuka GI. Food analysis and instrumentation. Theory and practice. Naphthali publishers, Lagos, Nigeria. 2005;48-56.
 15. Iwe MO, Onyeukwu U, Agiripa AN. Proximate, functional and pasting properties of FARO 44 rice, African yam bean and brown cowpea composite flours. Cogent Food and Agriculture. 2016;2:114-122.
 16. Butt MS, Batool R. Nutritional and functional properties of some promising legume protein isolates. Pakistan Journal of Nutrition. 2010;9(4):374-379.
 17. Nwosu JN, Ojukwu M, Ogueke CC, Ahaotu I, Owuamanam C. The anti-nutritional properties and ease of dehulling on the proximate composition of pigeon pea (*Cajanus cajan*) as affected by malting. International Journal of Life Sciences. 2013;2(2):60-67.
 18. Butt MS, Sharif M, Nasir M, Iftikhar S, Rehman K. Preparation and quality evaluation of cookies containing sweet potato flour. Industrial Journal of Plant Science. 2004;3:69-76.
 19. Nsa EE, Ukachukwu SN. Effect of thermal processing methods on the proximate composition, gross energy, mineral and ricin content of undecorticated castor oil seed (*Ricinus communis*). Global Journal of Agricultural Science. 2009;8(2):244-252.
 20. Duenas M, Hernandez J, Estrella I, Fernandez D. Germination as a process to increase the polyphenol content and antioxidant activity of lupin seeds (*Lupinus augusti folius L.*). Food Chemistry. 2009;117:599-607.
 21. Hamad AM, Fields MC. Reduction of protein quality and available lysine of germinated and ungerminated cereals. Journal of Food Science. 1997;44:456-459.
 22. Okaka JC, Akobundu ENT, Okaka ANT. Food and Human Nutrition. An Integrated Approach. Ojano Academic Publishers, Enugu, Nigeria. 2006;373- 395.
 23. Akubor PI, Chukwu JK. Proximate composition and selected functional properties of fermented and unfermented African oil bean (*Pentaclethra macrophylla*) seed flour. Plant Foods for Human Nutrition. 1999;54:227-238.
 24. Mbaeyi IE, Onweluzo JC. Effect of sprouting and pre-gelatinization on the physicochemical properties of sorghum-pigeon pea composite blend used for the production of breakfast cereal. Journal of Tropical Agriculture, Food, Environment and Extension. 2010;9(1):8-17.
 25. Akubor PI, Isolokwu PC, Ugane O, Onimawo IA. Proximate composition and functional properties of African bread fruit kernel and wheat flour blends. Food Research International. 2000;33:707-712.
 26. Uche SN, Charity UN, Abbas O, Aliyu M, Francis GB, Oche O. Proximate, anti-nutrients and mineral composition of raw and processed (boiled and roasted) *Sphenostylis stenocarpa* seeds from Southern Kaduna, Northwest Nigeria. International Scholarly Research Notices. 2014;20(4):280-287.
 27. Malomo O, Ogunmoyela OA, Olewajoba BSO, Kukoyi I. Effect of germinated soy flour on the sensory acceptability of soy-wheat composite bread. Asian Journal of Natural and Applied Sciences. 2012;2:48-53.
 28. Leonel M, Santos TB, Mischan MM. Physical characteristics of extruded cassava starch. Brazilian Journal of Chemical Engineering. 2009;20:129-137.
 29. Osundahunsi OF, Fagbemi TN, Kesselman E, Shimoni E. Comparison of the physicochemical properties and pasting characteristics of flour and starch from red and white sweet potato cultivars. Journal of Agriculture and Food Chemistry. 2003;51:2232-2236.
 30. Fasasi OS, Eleyinmi AF, Oyarekua MA. Effect of some traditional processing operations on the functional properties of African bread fruit (*Treculia africana*) seed flour. LWT Food Science and Technology. 2007;40:153-159.
 31. Onimawo AI, Akubor PI. Food Chemistry with Biochemical Approach. Ambik Press Ltd, Benin City, Edo State Nigeria. 2005;222-228.
 32. Falmata AS, Modu S, Badau HD, Babagana M, Bintu BP. Formulation and evaluation of complementary weaning food prepared from single and combined sprouted and fermented local red sorghum (*Sorghum bicolor*) variety blended with cowpea (*Vigna unguiculata*) and groundnut (*Arachis hypogea*). International Journal of Biotechnology and Food Science. 2014; 2(8):149-155.

33. Omojola A, Isah O, Adewumi M, Ogunsola O, Attah S. Evaluation of the effects of various additives on the acceptability of kilishi. *Tropical Journal of Animal Science*. 2004;6:10-43.
34. Akubor PI, Obiegbonam JE. Functional properties of ungerminated and germinated milled flour. *Journal of Food Science and Technology*. 1999;36(3):241-243.
35. Bolarinwa IF, Olaniyan SA, Adebayo LO, Ademola AA. Malted sorghum-soy composite flour: preparation, chemical and physicochemical properties. *Journal of Food Processing and Technology*. 2015;6(8):1-7.
36. Usman MA, Bolade MK, James S. Functional properties of weaning food blends from selected sorghum (*Sorghum bicolor* (L.) Moench) varieties and soybean (*Glycine max*). *African Journal of Food Science*. 2016;10(8):112-121.
37. Elofsson C, Dejmek P, Poulso M, Burlria H. Characterization of a cold whey concentrate. *International Journal of Dairy Science*. 1997;7:601-607.
38. Machiyo W, Tomoko M, Kikuchi T, Hirochi K, Naofumi M. Application of pre-germinated brown rice for bread making. *Cereal Chemistry*. 2004;3:293-317.
39. Newport Scientific. Applications manual for the rapid visco analyser using thermocline for windows. Newport Scientific Pty Ltd.1/2 Apollo Street, Warriewood NSW 2102, Australia. 1996;2-26.
40. Ragaei S, Abdel-Aal EM. Pasting properties of starch and protein in selected cereals and quality of their food products. *Food Chemistry*. 2006;95:9-18.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/108857>