



# Molecular Characterization of *Anaplasma spp* Isolated from Cattle Slaughtered at Central Abattoir Bauchi, Bauchi State, Nigeria

Hadiza Mudi <sup>a,b\*</sup>, Iliyasu Mahmud Yerima <sup>b</sup>,  
Samaila Adamu Babayo <sup>b</sup>, Panda Sam Mao <sup>b</sup>,  
Obishakin Emmanuel Feyi <sup>c</sup>, Shuaibu Hafsat Jagab <sup>c</sup>  
and Olaolu Olushola Samuel <sup>c</sup>

<sup>a</sup> Department of General studies, Aliko Dangote College of Nursing Sciences, Bauchi state, Nigeria.

<sup>b</sup> Biological sciences Department, Abubakar Tafawa Balewa University, Bauchi, Nigeria.

<sup>c</sup> Biotechnology Centre, National Veterinary Research Institute Vom, Plateau state, Nigeria.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113886>

Original Research Article

Received: 10/01/2024

Accepted: 14/03/2024

Published: 18/03/2024

## ABSTRACT

**Background:** This study was conducted to molecularly characterize *Anaplasma species* isolated from cattle slaughtered at the Bauchi central abattoir. Bovine anaplasmosis is a disease of cattle caused by *Anaplasma* haemoparasite. It has significant effects on the animal's health but has been overshadowed by other haemoparasitic diseases like Trypanosomiasis and Babebiosis. The discovery of new species with zoonotic potentials in recent years has given it more significance.

\*Corresponding author: E-mail: hadizamudi@gmail.com;

**Materials and Methods:** Blood samples were collected from the severed jugular vein of the animals at the point of slaughter and collected samples were taken to the laboratory for analysis. DNA extraction was carried out using the Quick -DNA™ Miniprep kit using the Quick protocol. Extracted DNA was amplified with specific primers targeting the 16S rRNA (≈430 bp) genes in the *Anaplasma* parasite. PCR was performed using BioinGentech Veterinary PCR kits (Concepcion, Chile) and amplified DNA fragments of the 16sRNA gene from the *Anaplasma* spp. isolates were directly sequenced using the Sanger method.

**Results:** Blood samples positive for *Anaplasma* specie by microscopy were further examined using molecular method to confirm the isolated parasite by targeting the 16SrRNA gene. 52% were found to be positive for 16srRNA with bands corresponding to 430bp of the Anaplasma primer and the amplified genes were directly sequenced using the Sanger method. Analyzed sequences were compared by BLASTn with those present in the GenBank database and *Anaplasma* specie obtained from this study clustered into four groups following phylogenetic analysis. *Anaplasma* species from Bauchi state clustered with other *Anaplasma* sequences and based on multiple alignment, the sequence presented nucleotide differences between them suggesting that different strains are circulating in the state based on the analyzed data. The DNA sequencing result of the 16SrRNA also revealed that *Anaplasma platys* and *Anaplasma phagocytophilum* are circulating among cattle herds in the state and this is of public health implication because *A. phagocytophilum* is zoonotic.

**Conclusion:** Five nucleotide sequences were obtained from this study and deposited in Genbank under the following accession numbers: OQ538141 for *Anaplasma platys*, OQ538142 for *Anaplama marginale*, OQ538143 for *Anaplasma phagocytophilum* and OQ538144 to OQ538145 for uncultured *Anaplasma* species.

**Keywords:** *Anaplasma*; cattle; abattoir; Bauchi; Nigeria.

## 1. INTRODUCTION

*Anaplasma* species is a haemoparasitic organism (protozoan) and is the causative agent of anaplasmosis. Up to 17 different tick vector species (including *Dermacentor*, *Rhipicephalus*, *Ixodes*, *Hyalomma*, and *Argas*) have been reported to transmit *Anaplasma* spp." [1]. "Not all of these are likely significant vectors in the field, and it has been shown that strains of *A. marginale* also coevolve with particular tick strains. *Rhipicephalus* spp. are major vectors in Australia and Africa, and *Dermacentor* spp have been incriminated as the main vectors in the USA" [2]. According to a study [3], "bovine anaplasmosis primarily caused by *A. marginale*, is considered as one of the most prevalent and costly tick-borne diseases of cattle globally". "It is reported as being endemic in most of the cattle-farming areas in Africa" [4].

"There is a huge economic impact of anaplasmosis from different countries which is corroborated by the widespread occurrence of the *Anaplasma* spp. in cattle" [5]. "The disease is becoming a serious concern for the animal breeding system, as the infection puts an additional burden on veterinary care by reducing the body weight of animals, decreasing milk

production, and frequently causing abortions leading to death" [6]. The red blood cells of cattle and wild ruminants are chosen by *A. marginale* and *A. centrale* as a site of infection, while small ruminants presenting the same cells are infected by *Anaplasma bovis* causing anaplasmosis, targets small mammals and ruminants, which results in the infection of monocytes. According to [7], "the prevalence rate is dependent on the type of species infected and the diagnostic method used". A study on molecular characterization conducted in some states by [8], revealed "the presence of mixed haplotypes in both *A. platys* and *A. marginale* in cattle in Nigeria".

"Traditionally, the identification, description and classification of *Anaplasma* species are based on the morphological characteristics using microscopic examination of Giemsa-stained thin or thick blood smears with or without serology. However, recent studies have demonstrated that *Anaplasma* and related piroplasmids are not monophyletic and can co-infect hosts; therefore, they require the use of molecular tools for the identification and characterization of these cryptic and polyphyletic species" [9,10]. It is reported that the small subunit ribosomal RNA (16S rRNA) gene is widely used for detection and

phylogenetic analysis of protozoan organisms due to its high level of conservation.

There is a need for the accurate identification of the causative agents of anaplasmosis considering the widespread distribution and the huge economic losses associated with the disease in order to design and facilitate implementation of effective control measures.

Molecular techniques have become very useful tools in parasite identification since they target the genes of the organism thus giving more accurate and definite information.

“The detection and characterization of *Anaplasma spp.* are performed using molecular tools in which sensitive and specific nucleic-acid based techniques targeting several genes such as the 16S rRNA, heat-shock protein (*groEL*), major surface protein-1a (*Msp1a*), *Msp2*, *Msp4* and *Msp5* have been developed to detect and characterize *Anaplasma spp.* infection in animals” [3].

PCR is considered to be the gold standard diagnostic approach for anaplasmosis [11], but it has not been used preferentially as a diagnostic tool in most *Anaplasma*-related epidemiological studies in Nigeria. This molecular based study reveals the first 16S ribosomal RNA based evidence of the Anaplasmosis in Bauchi state, Nigeria. This study employed the use of *Anaplasma/Ehrlichia* universal primers to determine the molecular profiles of *Anaplasma spp.* infecting cattle in the study area and analyzed the phylogenetic relationship and genetic diversity among the identified isolates and those from different countries. To the best of our knowledge, this is the first phylogenetic analysis of *Anaplasma spp.* circulating among cattle population in the area.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

The study was conducted between the months of May 2021 to January 2022 at the Bauchi central abattoir. The Bauchi Central abattoir is the largest abattoir in the state. It is about 6 kilometres away from the state capital and situated along Gombe Adamawa road in North Eastern Nigeria. The animals to be slaughtered are bought by butchers from nearby villages and towns markets. The abattoir has an average daily sacrifice of 50 cattle. The state lies between

Latitude 9.3° and 12.3° North of the Equator and Longitude 8.5° and 11° East of the Greenwich meridian. The state covers a total land area of 49,259.01sqkm which is about 5.3% of Nigeria's total land mass. It has a population of 4, 676,465 people based on the 2006 population census. Bauchi state is one of the states in the Northern part of Nigeria that spans two distinctive vegetation zones namely the Sudan savanna and the Sahel savanna.

### 2.2 Sample Collection

Samples were collected at the point of slaughter from the severed jugular vein into the Ethylene Diamine Tetra Acetate (EDTA) bottles which were immediately labelled and placed in a box containing ice packs. Each sample was assigned a Laboratory Identification Number. The blood samples were collected between the months of May 2021 to January 2022 at the Bauchi central abattoir. Both sexes of cattle were considered for the purpose of this research. The blood samples were transported to the National Veterinary Research Institute Vom for analysis. Samples were subjected to routine microscopy and positive samples obtained were stored at -20°C pending further analysis. The labelled blood samples were stored at -20°C pending DNA extraction.

### 2.3 Microscopic Examination

Blood samples were subjected to routine microscopy. *Anaplasma spp.* were detected using, stained thin blood film Giemsa [12]. The smears were examined at x100 objective lens (oil immersion) for presence of parasites and identification. A minimum of 50 fields was searched per slide in each of the procedures adopted. Positive samples obtained were labelled and stored at -20°C pending further analysis which is the DNA extraction.

### 2.4 DNA Extraction

DNA extraction was carried out using the Quick - DNA™ Miniprep kit catalogue Nos. D3024 & D3025 using the Quick protocol. For optimal performance, beta-mercapto-ethanol was added to the Genomic lysis buffer to form a final dilution of 0.5% (V/v) i.e. 250µl per 50ml. To lyse the sample, 4 volumes of Genomic Lysing buffer (800µl) was added to 200µl of the sample (whole blood) in a volume of 4:1. Samples were vortexed for 4-6 seconds (to break down the cells) and then allowed to stand at room temperature for 5-10 minutes. The mixture was

**Table 1. Oligonucleotide sequence of the primers used for Amplification of 16sRNA gene**

Genes	Primers	Oligonucleotide sequence	Amplicon size	Reference
16S rRNA	F R	GGTTTAATTCGATGCAACGCGA CGTATTACACCGTGGCATG	430	Simuunza, 2009 [13]

then transferred to a zymo-spin™ column in a collection tube and centrifuged at 12,000xg for 1 minute and then the collection tube with the flow through was discarded. Zymo-spin™ column was then transferred to a new collection tube and 200µl of DNA Pre-wash buffer was added to the spin column and was centrifuged at 12,000xg for 1 minute. A 500µl of g-DNA Wash buffer was added to the spin column and it was centrifuged at 12,000xg for 1 minute. The spin column was then transferred to a clean microcentrifuge tube. 50µl DNA Elution buffer was added to the spin column and was added incubated for 5 minutes at room temperature and then centrifuged at top speed of 16,000xg for 30 seconds to elute the DNA. All procedures were performed at room temperature (15-30°C) and all centrifugation steps were performed at 10,000xg-16,000xg using Eppendorf centrifuge 5417R. The eluted DNA was stored at -20°C pending to next analysis.

## 2.5 PCR Amplification of 16S rRNA Gene

Extracted DNA from the blood sample was amplified with specific primers targeting forward and reverse primers targeting the 16S rRNA (≈430 bp) genes in the *Anaplasma* parasite as described by [13]. The Oligonucleotide sequence of the primer is shown in Table 1. PCR was performed using BioinGentech Veterinary PCR kits (Concepcion, Chile) according to the instructions of the manufacturer. The primary PCR was performed with *Anaplasma/Ehrlichia* universal primers kit.

The PCR amplification followed the cycling conditions by initial denaturation at 94°C for 2 minutes, 30 cycles (94 °C for 30seconds, 57 °C for 30 seconds, 72 °C for 60 seconds) and final extension at 72 °C for 5 minutes.

## 2.6 Quantification of PCR Products by Agarose Gel Electrophoresis

The buffer used for elution in the subsequent steps were equilibrated at room temperature (15-25 °C) and the water bath was heated to 56 °C, 2% agarose gels was used for electrophoresis [14].

## 2.7 Genes Sequencing Analysis

Amplified DNA fragments of the 16sRNA gene from the *Anaplasma spp.* isolates were directly sequenced using the Sanger method at a commercial facility (Macrogen BV, Neitherlands) using the PCR primers, and the analyzed sequences were compared by BLASTn with those present in the GenBank database.

## 2.8 Nucleotide and Phylogenetic analysis

The 16S rRNA nucleotide sequences obtained in this study were edited manually using the Bioedit version 7.0.5.3 [15] and were compared to reference sequences available in the GenBank using the BLASTn algorithm (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

“Ten reference 16S rRNA sequences of different *Anaplasma spp.* were retrieved from GenBank and added to alignments of the sequences of the study. Phylogenetic analysis was inferred using the Neighbour-Joining method” [16]. “The bootstrap consensus tree inferred from 1000 replicates is taken to represent the evolutionary history of the taxa analyzed” [17].

## 3. RESULTS

### 3.1 Prevalence of *Anaplasma spp* Based on Microscopy

A total of 163 (42.4%) out of the 384 blood samples randomly collected from the cattle and examined were found to be positive for *Anaplasma marginale* based on microscopy. Fifty out of the 163 samples that were positive by microscopy were further examined using molecular method to confirm the *Anaplasma specie* isolated targeting the 16SrRNA gene of the parasite.

### 3.2 Molecular Characteristics of the *Anaplasma species* Detected from Cattle in the Study Area

Polymerase Chain Reaction (PCR) was used to characterize the pathogens observed in the microscopy. DNA extracted from the blood

samples were analysed and 52% of the samples were found to be positive for 16SrRNA in the *Anaplasma species* with band corresponding to 430 bp, as shown in the reference primers (Fig. 1).

### 3.3 Phylogenetic Analysis

Phylogenetic shows that the 16S rRNA nucleotide sequences of *Anaplasma specie*

obtained from this study clustered into four groups as shown in Fig. 2. “Branches corresponding to partitions reproduced in less than 50% bootstrap replicates are collapsed. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches” [17]. Evolutionary analyses were conducted in MEGA11 [18].




Fig. 1. Agarose gel image of PCR products (16SrRNA) of *Anaplasma species* with amplicon size of 430bp




Fig. 2. Phylogenetic tree based on the 16S rRNA nucleotide sequence of *Anaplasma spp.* of cattle from Bauchi state, Nigeria. The tree was inferred using the Neighbour-Joining method [16]. The evolutionary distances were computed using the Kimura 2-parameter method [19] and are in the units of the number of base substitutions per site. This analysis involved 16 nucleotide sequences. All positions containing gaps and missing data were eliminated. There were a total of 64 positions in the final dataset. Sequenced obtained in this study were identified by a black circle (●). Nucleotide sequence of *Ehrlichia canis* (MT066094.1) from Egypt was used as an outgroup

**Table 2. *Anaplasma* spp detected by BLASTn analysis of 16SrRNA gene sequences of the isolates from Bauchi state**

Isolate	Accession no. (this study)	Matching sequence	Accession no. highest match	E-value	% Identity
T7	OQ538141.1	<i>Anaplasma platys</i>	KU586158.1	0.0	>90%
T8	OQ538142.1	<i>Anaplasma marginale</i>	MK680807.1	0.0	>90%
T9	OQ538143.1	<i>Anaplasma phagocytophilum</i>	OP788189.1, OQ152552.1	0.0	>90%
T10, T11	OQ538144.1, OQ538145.1	Uncultured <i>Anaplasma</i> species	ON408402.1, MW019806.1	0.0	>90%

### 3.4 Nucleotide Sequence Accession Numbers for *Anaplasma* Detected in Cattle in Bauchi State

The partial 16SrRNA gene sequences obtained from this study were deposited in the Genbank. The gene sequences were deposited under accession numbers OQ538141 for *Anaplasma platys*, OQ538142 for *Anaplasma marginale*, OQ538143 for *Anaplasma phagocytophilum* and OQ538144 to OQ538145 for Uncultured *Anaplasma* species described as isolates T7, T8, T9, T10 and T11 respectively (Table 2).

## 4. DISCUSSION

A total of 50 blood samples from the cattle positive for *Anaplasma* species through microscopy were screened by PCR for the presence of *Anaplasma* parasite using primers targeting the 16SrRNA gene. *Anaplasma* organism was detected in 26 blood samples from the cattle when primers targeting the 16SrRNA was used. The overall positive rate was 52%. This high prevalence rate is consistent with the work done by [20] who reported a prevalence rate of 49.1% for *Anaplasma* genus by conventional PCR on cattle blood samples collected from 5 states of Nigeria.

The 16S rRNA nucleotide sequences of *Anaplasma* spp. obtained in this study were clustered into four groups following phylogenetic analysis. The sequence OQ538141.1 clustered with the sequence of *Anaplasma platys* from China in the GenBank with a high bootstrap value (>60%). Similarly, another sample, OQ538142.1 clustered with the sequence of *Anaplasma marginale* (MK680807.1) from Pakistan. Two sequences from this study,

OQ538144.1 and OQ538145.1 were placed in the same clade with Uncultured *Anaplasma* spp. from African countries, Mozambique (ON408402.1) and Kenya (MW019806.1). One sequence OQ538143.1, from this study formed a distinct clade and was placed between two *Anaplasma phagocytophilum* sequences (OP788189.1, OQ152552.1) from China (Fig. 2).

Previous studies conducted using 16SrRNA gene detected *Anaplasma* organisms in ruminants from China by PCR [21,22] is consistent with the findings of this study. However, most of the positive samples were detected by PCR for 16srRNA gene. This detection of DNA is an indication that *Anaplasma* parasite is present in cattle in Bauchi state. The result shows that cattle are exposed to ticks infected by the various *Anaplasma* species. This suggests that 16SrRNA can be said to be sensitive for the detection of *Anaplasma* species. [23] confirmed the presence of *Anaplasma* species in cattle by BLASTn analysis after the sequencing of the 16SrRNA gene. In this study, the BLASTn analysis of the 16SrRNA gene fragments revealed nucleotides homologous to those of *A. phagocytophilum* and *A. platys* from cattle samples. This finding indicates infection of cattle with *A. phagocytophilum* and *A. platys* in the study area and this was only identified based on the molecular study as microscopy showed only *Anaplasma marginale* in the blood samples examined. It is well known that *A. centrale* and *A. marginale* are the main *Anaplasma* species that infect cattle in Nigeria. However, the findings from this study have shown that cattle in Bauchi state can also contain various *Anaplasma* species including the zoonotic ones like *A. phagocytophilum*. This result is consistent with

the studies of [23] and [24] in other countries who found *A. phagocytophilum* to infect cattle.

*A. platys* was also detected in the cattle from this study and this implies that cattle could be potential hosts for various *Anaplasma* isolates different from the previous studies that cattle in the state are only infected with *A. marginale* and *A. centrale* and this is the first report of *A. phagocytophilum* and *A. platys* infecting cattle in Bauchi state even though there are other studies from Nigeria where they were also found. The discovery of *A. phagocytophilum* in cattle calls for further studies to determine its public health implications because it is a zoonotic pathogen and has public health implications.

## 5. CONCLUSION

The result from this study indicates that *Anaplasma* species from Bauchi state clustered with other *Anaplasma* sequences and based on multiple alignment, the sequence presented nucleotide differences between them suggesting that different strains are circulating in the state based on the analyzed data. DNA sequencing result of the 16SrRNA also revealed that *Anaplasma platys* and *Anaplasma phagocytophilum* are circulating among cattle herds in the state and this is of public health implication because *A. phagocytophilum* is zoonotic. Five nucleotide sequences were obtained from this study and deposited in Genbank under the following accession numbers: OQ538141 for *Anaplasma platys*, OQ538142 for *Anaplasma marginale*, OQ538143 for *Anaplasma phagocytophilum* and OQ538144 to OQ538145 for uncultured *Anaplasma* species

## ETHICAL APPROVAL

Ethical approval was obtained from the Animal use and care committee (AUCC) of the National veterinary Research institute Bauchi state (AEC/02/139/23). An introductory letter was collected from the Biological Sciences department of the ATBU and taken to the Abattoir before sample collection.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Alicja EL. Anaplasmosis. Merck Vet Manual;2020.

- Available:[http://www.merckvetmanual.com/mvm/circulatory system/blood parasites](http://www.merckvetmanual.com/mvm/circulatory_system/blood_parasites) 2015
2. Labruna MB. Ecology of rickettsia in South America. *Annals of the New York Acad of Sci.* 2009; 1166(1):156-166.
  3. Junsiri W, Watthanadirek A, Poolsawat N, Minsakorn S, Nooroong P, Jittapalapong S, Chawengkirttikul R, Anuracpreeda P. Molecular characterization of *Anaplasma marginale* based on the msp1a and msp1b genes. *Vet Microbiol* 2021;262:109236.
  4. Ramabu SS, Kgwatalala PM, Nsoso SJ, Gasebonwe S, Kgosiesele E. Anaplasma infection prevalence in beef and dairy cattle in the south east region of Botswana. *Vet Parasitol: Regional Studies and Reports.* 2018;12:4-8.
  5. Kocan KM, Cabezas-Cruz A. The genus *Anaplasma*: New challenges after reorganization. *Revue scientifique et technique –Office International de zootie.* 2015;34(2):577-586.
  6. Kocan KM, De la Fuente J, Guglielmone AA, Meléndez RD. Antigenes and alternatives for control of *Anaplasma marginale* infection in cattle. *Clinical Microbiol Reviews.* 2003;16(4):698-712.
  7. Belkahia H, Said MB, Ghribi R, Selmi R, Asker AB, Yahiaoui M, Bousrih M, Daaloul-Jedidi M, Messadi L. Molecular detection, genotyping and phylogeny of *Anaplasma* spp. in Rhipicephalus ticks from Tunisia. *Acta Tropica.* 2019;191:38-49.
  8. Kamani J, Schaer J, Umar AG, Pilarshimwi JY, Bukar L, González-Miguel J, Harrus S. Molecular detection and genetic characterization of *Anaplasma marginale* and *Anaplasma platys* in cattle in Nigeria. *Ticks and tick-borne diseases.* 2022;13(4):101955.
  9. Lack JB, Reichard MV, Van Den Bussche RA. Phylogeny and evolution of the Piroplasmida as inferred from 18S rRNA sequences. *Int Journal for Parasitol.* 2012;42(4):353-63.
  10. Van de Peer Y, Baldauf SL, Doolittle WF, Meyerid A. An updated and comprehensive rRNA phylogeny of (crown) eukaryotes based on rate-calibrated evolutionary distances. *Journal of Molecular Evolution.* 2000;51:565-76.
  11. de Echaide ST, Bono MF, Lugaresi C, Aguirre N, Mangold A, Moretta R, Farber M, Mondillo C. Detection of antibodies against *Anaplasma marginale* in milk using a recombinant MSP5 indirect ELISA.

- Veterinary microbiology. 2005;10;106(3-4):287-92.
12. Cheesbrough M. District laboratory practice in tropical countries. IJMS. 2018;1(1):65-8.
  13. Simuunza MC. Differential diagnosis of tick-borne diseases and population genetic analysis of *Babesia bovis* and *Babesia bigemina* (Doctoral dissertation, University of Glasgow); 2009.
  14. Anejo-Okopi JA, Okojokwu JO, Ebonyi AO, Ejeliogu EU, Isa SE, Audu O, Akpakpan EE, Nwachukwu EE, Ifokwe CK, Ali M, Lar P. Molecular characterization of *Cryptosporidium* in children aged 0-5 years with diarrhea in Jos, Nigeria. The Pan African Medical Journal. 2016;25.
  15. Hall TA. BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. In Nucleic acids symposium series 1999;41(41):95-98.
  16. Saitou N, Nei M. The neighbor-joining method: A new method for reconstructing phylogenetic trees. Molecular biology and evolution. 1987;4(4):406-25.
  17. Felsenstein J. Confidence limits on phylogenies: An approach using the bootstrap. Evolution. 1985;39(4):783-91.
  18. Tamura K, Stecher G, Kumar S. MEGA11: Molecular evolutionary genetics analysis version 11. Molecular Biology and Evolution. 2021;38(7):3022-7.
  19. Kimura M. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. Journal of Molecular Evolution. 1980;16:111-20.
  20. Kamani J, Schaer J, Umar AG, Pilarshimwi JY, Bukar L, González-Miguel J, Harrus S. Molecular detection and genetic characterization of *Anaplasma marginale* and *Anaplasma platys* in cattle in Nigeria. Ticks and Tick-Borne Diseases. 2022;13(4):101955.
  21. Zhang Y, Lv Y, Zhang F, Zhang W, Wang J, Cui Y, Wang R, Jian F, Zhang L, Ning C. Molecular and phylogenetic analysis of *Anaplasma spp.* in sheep and goats from six provinces of China. Journal of Veterinary Science. 2016;17(4):523-9.
  22. Yang J, Liu Z, Niu Q, Liu J, Xie J, Chen Q, Chen Z, Guan G, Liu G, Luo J, Yin H. Evaluation of different nested PCRs for detection of *Anaplasma phagocytophilum* in ruminants and ticks. BMC Veterinary Research. 2016;12(1):1-6.
  23. M'ghirbi Y, Bèji M, Oporto B, Khrouf F, Hurtado A, Bouattour A. *Anaplasma marginale* and *A. phagocytophilum* in cattle in Tunisia. Parasites & Vectors. 2016;9:1-8.
  24. Noaman V, Shayan P. A new PCR-RFLP method for detection of *Anaplasma marginale* based on 16S rRNA. Veterinary Research Communications. 2010;34:43-50.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/113886>